

SOME INTEGRAL INEQUALITIES VIA (p, q) -CALCULUS ON FINITE INTERVALS

MEVLÜT TUNÇ^α AND ESRA GÖV^β

ABSTRACT. The aim of this paper is to construct (p, q) -calculus on finite intervals. The (p_k, q_k) -derivative and (p_k, q_k) -integral are defined and some basic properties are given. Also, (p_k, q_k) -analogue of Hölder, Minkowski and Hermite-Hadamard inequality are proved.

1. INTRODUCTION

All of the scientific works deal with the ambition for giving the meaning of the universe in which we live. Every new discovery we made come up looking, feeling, living and transmitting in a different perspective. For understanding and transmitting these happenings, we all need different type of methods. As mathematicians, the main purpose of our studies is to *analyze* the nature and express in mathematical ways. In this sense, calculus which is the main well-known way become our alphabet while we are translating the universe into some notions.

Quantum calculus is a field that searches mathematical formulas which turn the original version when q tends to 1. The history of quantum analysis goes back to eighteenth century to when Euler introduced q in 'Introductio' in the tracks of Newton's infinite series. In nineteenth century, Jackson defined an integral which is called q -Jackson integral in 1910 and q -analysis has gone through a period of rapidly development. For more details, see [4, 5, 7, 10] and the references therein.

In recent years, as being one of the most desirable area, many authors are interested in quantum calculus. One can easily see new contributions to the field almost every day. This is due to the fact that quantum calculus has not also important applications in mathematics but also in particle physics, theoretical physics, analytic number theory, and computer science. In mathematics, q -analysis is closely linked with theory of ordinary fractional calculus, optimal control problems, q -difference and q -integral equations. In [17] and [18] Tariboon et al. define quantum calculus on finite intervals namely q_k -calculus, prove some of its properties and extend some of the important integral inequalities to quantum calculus.

In this paper, we give a generalization for the (p, q) -calculus which was first taken in [1] as (p, q) -integers for generalizing q -oscillator algebras which is well known in the earlier physics. Until then today, (p, q) -calculus become an appropriate workspace for both mathematicians and physicist, see [1, 2, 3, 6, 8, 9] and [11]-[15]. Our main goal is to open a new door for enlarging the field in which the studies

1991 *Mathematics Subject Classification.* 34A08, 26A33, 26D10, 26D20.

Key words and phrases. quantum calculus; (p, q) -derivative; (p, q) -integral; Hölder inequality; Minkowski inequality; Hermite-Hadamard inequality.

^β*Corresponding Author.*

This paper is in final form and no version of it will be submitted for publication elsewhere.

gain importance. The (p, q) -integers $[n]_{p,q}$ are defined by

$$(1.1) \quad [n]_{p,q} = \frac{p^n - q^n}{p - q}$$

where $0 < q < p \leq 1$. For each $k, n \in \mathbb{N}$, $n \geq k \geq 0$, the (p, q) -factorial and (p, q) -binomial are defined by

$$[n]_{p,q}! = \prod_{k=1}^n [k]_{p,q}, \quad n \geq 1, \quad [0]_{p,q}! = 1$$

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = \frac{[n]_{p,q}!}{[n-k]_{p,q}! [k]_{p,q}!}.$$

Let $f : \mathbb{R} \rightarrow \mathbb{R}$. The (p, q) -derivative of the function f is defined as

$$(1.2) \quad D_{p,q}f(x) = \frac{f(px) - f(qx)}{(p-q)x}, \quad x \neq 0$$

provided that $D_{p,q}f(0) = f'(0)$.

Let $f : C[0, a] \rightarrow \mathbb{R}$ ($a > 0$) then the (p, q) -integration of f defined by

$$(1.3) \quad \int_0^a f(t) d_{p,q}t = (q-p)a \sum_{n=0}^{\infty} \frac{p^n}{q^{n+1}} f\left(\frac{p^n}{q^{n+1}}a\right) \text{ if } \left|\frac{p}{q}\right| < 1$$

$$\int_0^a f(t) d_{p,q}t = (p-q)a \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} f\left(\frac{q^n}{p^{n+1}}a\right) \text{ if } \left|\frac{p}{q}\right| > 1.$$

The formula of (p, q) -integration by parts is given by

$$(1.4) \quad \int_a^b f(px) D_{p,q}g(x) d_{p,q}t = f(x)g(x)|_a^b - \int_a^b g(qx) D_{p,q}f(x) d_{p,q}t.$$

All notions written above reduce to the q -analogs when $p = 1$. For more details, see the references mentioned in above.

2. (p, q) -CALCULUS ON FINITE INTERVALS

In this section, we define (p_k, q_k) -derivative and (p_k, q_k) -integral on finite intervals. Let $I_k := [u_k, u_{k+1}]$ be an interval and $0 < q_k < p_k \leq 1$ be constants.

Definition 1. Let $f : I_k \rightarrow \mathbb{R}$ be a continuous function and assume that $u \in I_k$. Then the following equality

$$(2.1) \quad D_{p_k, q_k}f(u) = \frac{f(p_k u + (1-p_k)u_k) - f(q_k u + (1-q_k)u_k)}{(p_k - q_k)(u - u_k)}, \quad u \neq u_k$$

$$D_{p_k, q_k}f(u_k) = \lim_{u \rightarrow u_k} D_{p_k, q_k}f(u)$$

is called the (p_k, q_k) -derivative of a function f at u .

Obviously, f is (p_k, q_k) -differentiable on I_k provided $D_{p_k, q_k}f(u)$ exists for all $u \in I_k$. In (2.1), if $p_k = 1$, then $D_{p_k, q_k}f = D_{q_k}f$ which is the q_k -derivative of the function f and also if $q_k \rightarrow 1, u_k = 0$, (2.1) reduces to q -derivative of the function f , see [10, 18].

Example 1. For $u \in I_k$, if $f(u) = (u - u_k)^n$, then

$$\begin{aligned}
 D_{p_k, q_k} f(u) &= \frac{(p_k u + (1 - p_k) u_k - u_k)^n - (q_k u + (1 - q_k) u_k - u_k)^n}{(p_k - q_k)(u - u_k)} \\
 &= \frac{p_k^n (u - u_k)^n - q_k^n (u - u_k)^n}{(p_k - q_k)(u - u_k)} \\
 (2.2) \quad &= [n]_{p, k} (u - u_k)^{n-1}
 \end{aligned}$$

where $[n]_{p_k, q_k} = \frac{p_k^n - q_k^n}{p_k - q_k}$. If $p_k = 1$ in (2.2), then (2.2) reduces

$$D_{q_k} f(u) = [n]_{q_k} (u - u_k)^{n-1}$$

which is given in [18]. Also if $q_k \rightarrow 1, u_k = 0$, it reduces q -derivative of the given function, see [10].

Theorem 1. Suppose that $f, g : I_k \rightarrow \mathbb{R}$ is (p_k, q_k) -differentiable on I_k . Then:

(a) If $f + g : I_k \rightarrow \mathbb{R}$ is (p_k, q_k) -differentiable on I_k , then

$$(2.3) \quad D_{p_k, q_k} (f(u) + g(u)) = D_{p_k, q_k} f(u) + D_{p_k, q_k} g(u).$$

(b) If $\lambda f : I_k \rightarrow \mathbb{R}$ is (p_k, q_k) -differentiable on I_k for any constant λ , then

$$(2.4) \quad D_{p_k, q_k} f(u) = \lambda D_{p_k, q_k} f(u).$$

(c) If $fg : I_k \rightarrow \mathbb{R}$ is (p_k, q_k) -differentiable on I_k , then

$$\begin{aligned}
 (2.5) \quad D_{p_k, q_k} (fg)(u) &= g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(u) + f(q_k u + (1 - q_k) u_k) D_{p_k, q_k} g(u) \\
 &= f(p_k u + (1 - p_k) u_k) D_{p_k, q_k} g(u) + g(q_k u + (1 - q_k) u_k) D_{p_k, q_k} f(u)
 \end{aligned}$$

(d) If $g(p_k u) g(q_k u + (1 - \frac{q_k}{p_k}) u_k) \neq 0$, then $\frac{f}{g}$ is (p_k, q_k) -differentiable on I_k with

$$(2.6) \quad D_{p_k, q_k} \left(\frac{f}{g} \right) (u) = \frac{g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(u) - f(p_k u + (1 - p_k) u_k) D_{p_k, q_k} g(u)}{g(p_k u + (1 - p_k) u_k) g(q_k u + (1 - q_k) u_k)}.$$

Proof. The proofs of (a) and (b) are obvious.

(c) From Definition 1, we have

$$\begin{aligned}
 &D_{p_k, q_k} (fg)(u) \\
 &= \frac{f(p_k u + (1 - p_k) u_k) g(p_k u + (1 - p_k) u_k) - f(q_k u + (1 - q_k) u_k) g(p_k u + (1 - p_k) u_k)}{(p_k - q_k)(u - u_k)} \\
 &\quad + \frac{f(q_k u + (1 - q_k) u_k) g(p_k u + (1 - p_k) u_k) - f(q_k u + (1 - q_k) u_k) g(q_k u + (1 - q_k) u_k)}{(p_k - q_k)(u - u_k)} \\
 &= g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(u) + f(q_k u + (1 - q_k) u_k) D_{p_k, q_k} g(u)
 \end{aligned}$$

The second equation can be proved in similar way by interchanging the functions f and g .

(d) From Definition 1, we have

$$\begin{aligned}
& D_{p_k, q_k} \left(\frac{f}{g} \right) (u) \\
&= \frac{\left(\frac{f}{g} \right) (p_k u + (1 - p_k) u_k) - \left(\frac{f}{g} \right) (q_k u + (1 - q_k) u_k)}{(p_k - q_k) (u - u_k)} \\
&= \frac{f(p_k u + (1 - p_k) u_k) g(q_k u + (1 - q_k) u_k) - g(p_k u + (1 - p_k) u_k) f(q_k u + (1 - q_k) u_k)}{g(p_k u + (1 - p_k) u_k) g(q_k u + (1 - q_k) u_k) (p_k - q_k) (u - u_k)} \\
&= \frac{g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(u) - f(p_k u + (1 - p_k) u_k) D_{p_k, q_k} g(u)}{g(p_k u + (1 - p_k) u_k) g(q_k u + (1 - q_k) u_k)}
\end{aligned}$$

□

Definition 2. Let $f : I_k \rightarrow \mathbb{R}$ be a continuous function. If $D_{p_k, q_k} f$ is (p_k, q_k) -differentiable on I_k , the second-order derivative is defined as $D_{p_k, q_k}^2 f$ with $D_{p_k, q_k} (D_{p_k, q_k} f) : I_k \rightarrow \mathbb{R}$. By this way, we obtain n -th order (p_k, q_k) -derivative $D_{p_k, q_k}^n f : I_k \rightarrow \mathbb{R}$.

For instance, if $f : I_k \rightarrow \mathbb{R}$, then we have

$$\begin{aligned}
D_{p_k, q_k}^2 f(u) &= D_{p_k, q_k} (D_{p_k, q_k} f) (u) \\
&= \frac{D_{p_k, q_k} f(p_k u + (1 - p_k) u_k) - D_{p_k, q_k} f(q_k u + (1 - q_k) u_k)}{(p_k - q_k) (u - u_k)} \\
&= \frac{\frac{f(p_k(p_k u + (1 - p_k) u_k) + (1 - p_k) u_k) - f(q_k(p_k u + (1 - p_k) u_k) + (1 - q_k) u_k)}{(p_k - q_k)(p_k u + (1 - p_k) u_k - u_k)}}{(p_k - q_k) (u - u_k)} \\
&= \frac{\frac{f(p_k(q_k u + (1 - q_k) u_k) + (1 - p_k) u_k) - f(q_k(q_k u + (1 - q_k) u_k) + (1 - q_k) u_k)}{(p_k - q_k)(q_k u + (1 - q_k) u_k - u_k)}}{(p_k - q_k) (u - u_k)} \\
&= \frac{f(p_k(p_k u + (1 - p_k) u_k) + (1 - p_k) u_k) - f(q_k(p_k u + (1 - p_k) u_k) + (1 - q_k) u_k)}{p_k (p_k - q_k)^2 (u - u_k)^2} \\
&\quad - \frac{f(p_k(q_k u + (1 - q_k) u_k) + (1 - p_k) u_k) - f(q_k(q_k u + (1 - q_k) u_k) + (1 - q_k) u_k)}{q_k (p_k - q_k)^2 (u - u_k)^2} \\
&= \frac{f(p_k^2 u + (1 - p_k^2) u_k) - f(q_k p_k u + (1 - q_k p_k) u_k)}{p_k (p_k - q_k)^2 (u - u_k)^2} \\
&\quad - \frac{f(p_k q_k u + (1 - p_k q_k) u_k) - f(q_k^2 u + (1 - q_k^2) u_k)}{q_k (p_k - q_k)^2 (u - u_k)^2} \\
&= \frac{q_k f(p_k^2 u + (1 - p_k^2) u_k) - (p_k + q_k) f(p_k q_k u + (1 - p_k q_k) u_k) + p_k f(q_k^2 u + (1 - q_k^2) u_k)}{p_k q_k (p_k - q_k)^2 (u - u_k)^2}
\end{aligned}$$

and $D_{p_k, q_k}^2 f(u_k) = \lim_{u \rightarrow u_k} D_{p_k, q_k}^2 f(u)$.

We define the (p_k, q_k) -integration as the inverse (p_k, q_k) -differentiation. Assume that T_{p_k, q_k} is a shifting operator defined by

$$(2.7) \quad T_{p_k, q_k} F(u) = F\left(\frac{q_k}{p_k} u + \left(1 - \frac{q_k}{p_k}\right) u_k\right)$$

where $F(u)$ is the (p_k, q_k) -antiderivative of f . Applying mathematical induction to (2.7), we see

$$(2.8) \quad T_{p_k, q_k}^n F(u) = F\left(\frac{q_k^n}{p_k^n} u + \left(1 - \frac{q_k^n}{p_k^n}\right) u_k\right)$$

where $n = 1, 2, \dots$ and $T_{p_k, q_k}^0 F(u) = F(u)$. From Definition 1, we have

$$f(u) = \frac{F(p_k u + (1 - p_k) u_k) - F(q_k u + (1 - q_k) u_k)}{(p_k - q_k)(u - u_k)}.$$

Making a change of variable, $(p_k u + (1 - p_k) u_k) = t$, we have

$$\begin{aligned} f\left(\frac{t - (1 - p_k) u_k}{p_k}\right) &= \frac{F(t) - F\left(\frac{q_k}{p_k} t + \left(1 - \frac{q_k}{p_k}\right) u_k\right)}{\left(\frac{p_k - q_k}{p_k}\right)(t - u_k)} \\ &= \frac{1 - T_{p_k, q_k}}{\left(\frac{p_k - q_k}{p_k}\right)(u - u_k)} F(t). \end{aligned}$$

Thus, we obtain

$$F(t) = \frac{1}{1 - T_{p_k, q_k}} \left(1 - \frac{q_k}{p_k}\right) (t - u_k) f\left(\frac{t - (1 - p_k) u_k}{p_k}\right).$$

Therefore, applying the formula of expansion of geometric series to (2.8), we have the following formula

$$\begin{aligned} F(t) &= \left(1 - \frac{q_k}{p_k}\right) \sum_{n=0}^{\infty} T_{p_k, q_k}^n (t - u_k) f\left(\frac{t - (1 - p_k) u_k}{p_k}\right) \\ &= \left(1 - \frac{q_k}{p_k}\right) \sum_{n=0}^{\infty} \left(\frac{q_k^n}{p_k^n} t + \left(1 - \frac{q_k^n}{p_k^n}\right) u_k - u_k\right) f\left(\frac{1}{p_k} \left(\frac{q_k^n}{p_k^n} t + \left(1 - \frac{q_k^n}{p_k^n}\right) u_k\right) + \left(1 - \frac{1}{p_k}\right) u_k\right) \\ &= (p_k - q_k) (t - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}} t + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right) u_k\right). \end{aligned}$$

Thus, we get

$$F(u) = (p_k - q_k) (u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}} u + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right) u_k\right)$$

Now, we define the (p_k, q_k) -integral of f on a finite interval as follows:

Definition 3. Let $f : I_k \rightarrow \mathbb{R}$ is a continuous function. Then for $0 < q_k < p_k \leq 1$,

$$(2.9) \quad \int_{u_k}^u f(s) d_{p_k, q_k} s = (p_k - q_k) (u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}} u + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right) u_k\right)$$

is called (p_k, q_k) -integral of f for $u \in I_k$.

Moreover, if $a \in (u_k, u)$, then (p_k, q_k) -integral is defined by

$$\begin{aligned}
(2.10) \quad & \int_a^u f(s) d_{p_k, q_k} s \\
&= \int_{u_k}^u f(s) d_{p_k, q_k} s \int_{u_k}^a f(s) d_{p_k, q_k} s \\
&= (p_k - q_k)(u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}}u + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right)u_k\right) \\
&\quad - (p_k - q_k)(a - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}}a + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right)u_k\right).
\end{aligned}$$

Note that if $u_k = 0$ and $p = 1$, then (2.10) reduces to q_k -integral of the function. See, [18].

Remark 1. We assume $0 < q_k < p_k \leq 1$ for all of the above results. We shall mention that $0 < q_k < 1$, $0 < p_k \leq 1$ for interchanging p_k and q_k in the formulas. So, we have

$$\begin{aligned}
(2.11) \quad & \int_{u_k}^u f(s) d_{p_k, q_k} s = (p_k - q_k)(u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f\left(\frac{q_k^n}{p_k^{n+1}}u + \left(1 - \frac{q_k^n}{p_k^{n+1}}\right)u_k\right), \left|\frac{p}{q}\right| > 1 \\
& \int_{u_k}^u f(s) d_{p_k, q_k} s = (q_k - p_k)(u - u_k) \sum_{n=0}^{\infty} \frac{p_k^n}{q_k^{n+1}} f\left(\frac{p_k^n}{q_k^{n+1}}u + \left(1 - \frac{p_k^n}{q_k^{n+1}}\right)u_k\right), \left|\frac{p}{q}\right| < 1.
\end{aligned}$$

where $0 < q_k < 1$, $0 < p_k \leq 1$.

Remark 2. Note that, if we take $u_k = 0$ in (2.11), then (2.11) reduces to (1.3), [14, Definition 5]. Also, if $p_k = 1$ in (2.9), then (2.9) reduces to q_k -integral of a function f defined by

$$\int_{u_k}^u f(s) d_{q_k} s = (1 - q_k)(u - u_k) \sum_{n=0}^{\infty} q_k^n f(q_k^n u + (1 - q_k^n)u_k).$$

For more details, see [18].

Theorem 2. The following formulas hold for $u \in I_k$:

- (a) $D_{p_k, q_k} \int_{u_k}^u f(s) d_{p_k, q_k} s = f(u)$
- (b) $\int_{u_k}^u D_{p_k, q_k} f(s) d_{p_k, q_k} s = f(u)$
- (c) $\int_a^u D_{p_k, q_k} f(s) d_{p_k, q_k} s = f(u) - f(a)$, for $a \in (u_k, u)$.

Proof. (a) From Definition 1 and Definition 3, we obtain

$$\begin{aligned}
& D_{p_k, q_k} \int_{u_k}^u f(s) d_{p_k, q_k} s \\
&= D_{p_k, q_k} \left[(p_k - q_k) (u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f \left(\frac{q_k^n}{p_k^{n+1}} u + \left(1 - \frac{q_k^n}{p_k^{n+1}} \right) u_k \right) \right] \\
&= (p_k - q_k) \left\{ \frac{(p_k u + (1 - p_k) u_k - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f \left(\frac{q_k^n}{p_k^{n+1}} (p_k u + (1 - p_k) u_k) + \left(1 - \frac{q_k^n}{p_k^{n+1}} \right) u_k \right)}{(p_k - q_k) (u - u_k)} \right. \\
&\quad \left. - \frac{(q_k u + (1 - q_k) u_k - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} f \left(\frac{q_k^n}{p_k^{n+1}} (q_k u + (1 - q_k) u_k) + \left(1 - \frac{q_k^n}{p_k^{n+1}} \right) u_k \right)}{(p_k - q_k) (u - u_k)} \right\} \\
&= \left[\sum_{n=0}^{\infty} \frac{q_k^n}{p_k^n} f \left(\frac{q_k^n}{p_k^n} u + \left(1 - \frac{q_k^n}{p_k^n} \right) u_k \right) - \sum_{n=0}^{\infty} \frac{q_k^{n+1}}{p_k^{n+1}} f \left(\frac{q_k^{n+1}}{p_k^{n+1}} u + \left(1 - \frac{q_k^{n+1}}{p_k^{n+1}} \right) u_k \right) \right] \\
&= f(u).
\end{aligned}$$

(b) From Definition 1 and Definition 3, we get

$$\begin{aligned}
& \int_{u_k}^u D_{p_k, q_k} f(s) d_{p_k, q_k} s \\
&= \int_{u_k}^u \frac{f(p_k s + (1 - p_k) u_k) - f(q_k s + \left(1 - \frac{q_k}{p_k} \right) u_k)}{(p_k - q_k) (s - u_k)} d_{p_k, q_k} s \\
&= (p_k - q_k) (u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} \left[\frac{f \left(\frac{q_k^n}{p_k^{n+1}} (p_k u + (1 - p_k) u_k) + \left(1 - \frac{q_k^n}{p_k^{n+1}} \right) u_k \right)}{(p_k - q_k) (u - u_k)} \right. \\
&\quad \left. - \frac{f \left(\frac{q_k^n}{p_k^{n+1}} (q_k u + (1 - q_k) u_k) + \left(1 - \frac{q_k^n}{p_k^{n+1}} \right) u_k \right)}{(p_k - q_k) (u - u_k)} \right] \\
&= (u - u_k) \sum_{n=0}^{\infty} \frac{q_k^n}{p_k^{n+1}} \left[\frac{f \left(\frac{q_k^n}{p_k^n} u + \left(1 - \frac{q_k^n}{p_k^n} \right) u_k \right)}{(u - u_k)} \right. \\
&\quad \left. - \frac{f \left(\frac{q_k^{n+1}}{p_k^{n+1}} u + \left(1 - \frac{q_k^{n+1}}{p_k^{n+1}} \right) u_k \right)}{(u - u_k)} \right] \\
&= \sum_{n=0}^{\infty} f \left(\frac{q_k^n}{p_k^n} p_k u + \left(1 - \frac{q_k^n}{p_k^n} \right) u_k \right) - f \left(\frac{q_k^{n+1}}{p_k^{n+1}} p_k u + \left(1 - \frac{q_k^{n+1}}{p_k^{n+1}} \right) u_k \right) \\
&= f(u).
\end{aligned}$$

(c) The proof is carried on from the part of (b). \square

Theorem 3. Let $f, g : I_k \rightarrow \mathbb{R}$ are continuous functions. The following formulas hold:

$$(a) \int_{u_k}^u [f(s) + g(s)] d_{p_k, q_k} s = \int_{u_k}^u f(s) d_{p_k, q_k} s + \int_{u_k}^u g(s) d_{p_k, q_k} s;$$

$$\begin{aligned}
(b) \int_{u_k}^u \lambda f(s) d_{p_k, q_k} s &= \lambda \int_{u_k}^u f(s) d_{p_k, q_k} s; \\
(c) \int_{u_k}^u f(q_k s + (1 - q_k) u_k) D_{p_k, q_k} g(p_k s) d_{p_k, q_k} s &= (fg)(s)_{u_k}^u - \int_{u_k}^u g(p_k s + (1 - p_k) u_k) D_{p_k, q_k} f(s) d_{p_k, q_k} s \\
\text{or} \\
\int_{u_k}^u f(p_k s + (1 - p_k) u_k) D_{p_k, q_k} g(q_k s) d_{p_k, q_k} s &= (fg)(s)_{u_k}^u - \int_{u_k}^u g(q_k s + (1 - q_k) u_k) D_{p_k, q_k} f(s) d_{p_k, q_k} s
\end{aligned}$$

Theorem 4. where $u \in I_k, \lambda \in \mathbb{R}$.

Proof. The proofs of (a)-(b) are derived from Definition 3.

(c) From (2.5), we write

$$f(q_k u + (1 - q_k) u_k) D_{p_k, q_k} g(u) = D_{p_k, q_k} (fg)(u) - g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(u).$$

By integrating over $[u_k, u]$ and using Theorem 3 part (b), we get

$$\begin{aligned}
& \int_{u_k}^u f(q_k u + (1 - q_k) u_k) D_{p_k, q_k} g(p_k s) d_{p_k, q_k} s \\
&= (fg)(u) - \int_{u_k}^u g(p_k u + (1 - p_k) u_k) D_{p_k, q_k} f(s) d_{p_k, q_k} s.
\end{aligned}$$

□

3. INTEGRAL INEQUALITIES ON FINITE INTERVALS

Lets start with (p, q) -Hölder integral inequality on $I = [a, b]$:

Theorem 5. Let f and g be two functions defined on I , $0 < q < p \leq 1$ and

$s_1, s_2 > 1$ with $\frac{1}{s_1} + \frac{1}{s_2} = 1$. Then

$$(3.1) \quad \int_a^b |f(t)g(t)| {}_a d_{p, q} t \leq \left(\int_a^b |f(t)|^{s_1} {}_a d_{p, q} t \right)^{\frac{1}{s_1}} \left(\int_a^b |g(t)|^{s_2} {}_a d_{p, q} t \right)^{\frac{1}{s_2}}.$$

Proof. From Definition 3 and discrete Hölder inequality, we get

$$\begin{aligned}
\int_a^b |f(t)g(t)| {}_a d_{p, q} t &= (p - q)(b - a) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \left| f \left(\frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right) g \frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right| \\
&= (p - q)(b - a) \sum_{n=0}^{\infty} \left| f \left(\frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right) \left(\frac{q^n}{p^{n+1}} \right)^{\frac{1}{s_1}} \right| \\
&\quad \times \left| g \left(\frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right) \left(\frac{q^n}{p^{n+1}} \right)^{\frac{1}{s_2}} \right| \\
&\leq \left((p - q)(b - a) \sum_{n=0}^{\infty} \left| f \left(\frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right) \right|^{s_1} \left(\frac{q^n}{p^{n+1}} \right) \right)^{\frac{1}{s_1}} \\
&\quad \times \left((p - q)(b - a) \sum_{n=0}^{\infty} \left| g \left(\frac{q^n}{p^{n+1}} b + \left(1 - \frac{q^n}{p^{n+1}} \right) a \right) \right|^{s_2} \left(\frac{q^n}{p^{n+1}} \right) \right)^{\frac{1}{s_2}} \\
&= \left(\int_a^b |f(t)|^{s_1} {}_a d_{p, q} t \right)^{\frac{1}{s_1}} \left(\int_a^b |g(t)|^{s_2} {}_a d_{p, q} t \right)^{\frac{1}{s_2}}.
\end{aligned}$$

Thus, the proof is complete. □

It easy to show that we obtain the same result in the statement $p < q$.

Corollary 1. *Under the assumptions of Theorem 5, if we take $s_1 = s_2 = 2$, then we have the following formula,*

$$(3.2) \quad \int_a^b |f(t)g(t)| {}_a d_{p,q}t \leq \left(\int_a^b |f(t)|^2 {}_a d_{p,q}t \right)^{\frac{1}{2}} \left(\int_a^b |g(t)|^2 {}_a d_{p,q}t \right)^{\frac{1}{2}}$$

which we call (p, q)-Cauchy-Schwarz integral inequality.

Remark 3. *If $p = 1$, (3.1) and (3.2) reduces to q-Hölder integral inequality and q-Cauchy-Schwarz integral inequality respectively.*

Theorem 6. *Let f and g real-valued functions on $[a, b]$ such that $|f|^{s_1}$, $|g|^{s_1}$ and $|f + g|^{s_1}$ are (p, q)-integrable functions on $[a, b]$, $0 < q < p \leq 1$ and $s_1 > 1$. Then*

$$(3.3) \quad \left(\int_a^b |f(t) + g(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}} \leq \left(\int_a^b |f(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}} + \left(\int_a^b |g(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}}.$$

Equality holds if and only if $f(t) = 0$ almost everywhere or $g(t) = \mu f(t)$ almost everywhere with a constant $\mu \geq 0$.

Proof. Since $|f|^{s_1}$, $|g|^{s_1}$ and $|f + g|^{s_1}$ are (p, q)-integrable on $[a, b]$, by using the triangle inequality, we can write

$$\begin{aligned} \int_a^b |f(t) + g(t)|^{s_1} {}_a d_{p,q}t &= \int_a^b |f(t) + g(t)| |f(t) + g(t)| {}_a d_{p,q}t \\ &\leq \int_a^b |f(t)| |f(t) + g(t)|^{s_1-1} {}_a d_{p,q}t + \int_a^b |g(t)| |f(t) + g(t)|^{s_1-1} {}_a d_{p,q}t. \end{aligned}$$

Taking $s_1, s_2 > 1$ with $\frac{1}{s_1} + \frac{1}{s_2} = 1$ and using (p, q)–Hölder integral inequality, we have

$$(3.4) \quad \int_a^b |f(t)| |f(t) + g(t)|^{s_1-1} {}_a d_{p,q}t \leq \left(\int_a^b |f(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}} \left(\int_a^b |f(t) + g(t)|^{(s_1-1)s_2} {}_a d_{p,q}t \right)^{\frac{1}{s_2}}$$

and

$$(3.5) \quad \int_a^b |g(t)| |f(t) + g(t)|^{s_1-1} {}_a d_{p,q}t \leq \left(\int_a^b |g(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}} \left(\int_a^b |f(t) + g(t)|^{(s_1-1)s_2} {}_a d_{p,q}t \right)^{\frac{1}{s_2}}.$$

Since $(s_1 - 1)s_2 = s_1$, from (3.4) and (3.5), it easy to see that

$$\left(\int_a^b |f(t) + g(t)|^{s_1} {}_a d_{p,q}t \right)^{1-\frac{1}{s_2}} \leq \left(\int_a^b |f(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}} + \left(\int_a^b |g(t)|^{s_1} {}_a d_{p,q}t \right)^{\frac{1}{s_1}}$$

from which we obtain the required inequality. \square

Remark 4. *If $p = 1$, (3.3) reduces to*

$$\left(\int_a^b |f(t) + g(t)|^{s_1} d_qt \right)^{\frac{1}{s_1}} \leq \left(\int_a^b |f(t)|^{s_1} d_qt \right)^{\frac{1}{s_1}} + \left(\int_a^b |g(t)|^{s_1} d_qt \right)^{\frac{1}{s_1}}$$

which can be called q-Minkowski integral inequality.

Next, we present the (p, q) –Hermite-Hadamard integral inequality on $[a, b]$.

Theorem 7. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function. Then*

$$(3.6) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) {}_a d_{p,q} x \leq \frac{(p+q-1)f(a) + f(b)}{p+q}$$

Proof. Since f is convex on $[a, b]$, we know that

$$(3.7) \quad f((1-t)a + tb) \leq (1-t)f(a) + tf(b)$$

for all $t \in [0, 1]$. By taking (p, q) –integration for (3.7) over t on $[0, 1]$ for $p/q > 1$, we get

$$\begin{aligned} \int_0^1 f((1-t)a + tb) {}_0 d_{p,q} t &\leq \int_0^1 (1-t)f(a) {}_0 d_{p,q} t + \int_0^1 tf(b) {}_0 d_{p,q} t \\ &= f(a)(p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} - f(a)(p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \frac{q^n}{p^{n+1}} \\ &\quad + f(b)(p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} \frac{q^n}{p^{n+1}} \\ &= f(a) - \frac{f(a)}{p+q} + \frac{f(b)}{p+q} \\ &= \frac{(p+q-1)f(a) + f(b)}{p+q}. \end{aligned}$$

For $p/q < 1$; we get

$$\begin{aligned} \int_0^1 f((1-t)a + tb) {}_0 d_{p,q} t &\leq \int_0^1 (1-t)f(a) {}_0 d_{p,q} t + \int_0^1 tf(b) {}_0 d_{p,q} t \\ &= f(a)(q-p) \sum_{n=0}^{\infty} \frac{p^n}{q^{n+1}} - f(a)(q-p) \sum_{n=0}^{\infty} \frac{p^n}{q^{n+1}} \frac{p^n}{q^{n+1}} \\ &\quad + f(b)(q-p) \sum_{n=0}^{\infty} \frac{p^n}{q^{n+1}} \frac{p^n}{q^{n+1}} \\ &= f(a) - \frac{f(a)}{p+q} + \frac{f(b)}{p+q} \\ &= \frac{(p+q-1)f(a) + f(b)}{p+q} \end{aligned}$$

which gives the right hand side of (3.6). Also

$$\begin{aligned} \int_0^1 f((1-t)a + tb) d_{p,q} t &= (p-q) \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} f\left(\left(1 - \frac{q^n}{p^{n+1}}\right)a + \frac{q^n}{p^{n+1}}b\right) \\ &= \frac{(p-q)(b-a)}{b-a} \sum_{n=0}^{\infty} \frac{q^n}{p^{n+1}} f\left(\frac{q^n}{p^{n+1}}b + \left(1 - \frac{q^n}{p^{n+1}}\right)a\right) \\ &= \frac{1}{b-a} \int_a^b f(x) {}_a d_{p,q} x. \end{aligned}$$

To prove the left hand side, we write

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq f\left(\frac{(1-t)a+tb}{2} + \frac{ta+(1-t)b}{2}\right) \\ &\leq \frac{1}{2}[f((1-t)a+tb) + f(ta+(1-t)b)] \end{aligned}$$

and by integrating both side over $[0, 1]$ and making the change of variable, we get

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2} \left[\int_0^1 f((1-t)a+tb) {}_0d_{p,qt} + \int_0^1 f(ta+(1-t)b) {}_0d_{p,qt} \right] \\ &= \frac{1}{b-a} \int_a^b f(x) {}_ad_{p,q}x. \end{aligned}$$

□

Remark 5. If $p = 1$, (3.6) reduces to

$$(3.8) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) {}_ad_qx \leq \frac{qf(a) + f(b)}{1+q}$$

which is given in [17, Theorem 3.2]. One can easliy see that when $q \rightarrow 1$ in (3.8), the inequality turns classical Hermite-Hadamard integral inequality.

REFERENCES

- [1] R. Chakrabarti, R. Jagannathan, A (p, q) -oscillator realization of two-parameter quantum algebras, J. Phys. A: Math. Gen. 24 (1991), L711-L718.
- [2] I. Burban, Two-parameter deformation of the oscillator algebra and (p, q) analog of two dimensional conformal field theory. J. Nonlinear Math. Phys. 2(3-4), 384-391 (1995).
- [3] I. M. Burban and A. U. Klimyk, P, Q -differentiation, P, Q -integration and P, Q -hypergeometric functions related to quantum groups, Integral Transforms and Special Functions 2 (1994), 15-36.
- [4] T. Ernst, A Comprehensive Treatment of q -Calculus, Springer Basel, 2012.
- [5] T. Ernst, A method for q -calculus, J. Nonlinear Math. Phys. 10 (4) (2003) 487-525.
- [6] R. Floreanini, L. Lapointe and L. Vinet, A note on (p, q) -oscillators and bibasic hypergeometric functions, J. Phys. A: Math. Gen. 26 (1993), L611-L614.
- [7] H. Gauchman, Integral inequalities in q -calculus, Comput. Math. Appl. 47 (2004) 281-300.
- [8] M.N. Hounkonnou, J. Désiré, B. Kyemba, $R(p, q)$ -calculus: differentiation and integration, SUT Journal of Mathematics, Vol. 49, No. 2 (2013), 145-167.
- [9] R. Jagannathan, K.S. Rao, Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series. In: Proceedings of the International Conference on Number Theory and Mathematical Physics, pp. 20-21 (2005).
- [10] V. Kac, P. Cheung, Quantum Calculus, Universitext, Springer-Verlag, New York, 2002.
- [11] M. Mursaleen, K. J. Ansari and A. Khan, On (p, q) -analogue of Bernstein Operators, Appl. Math. Comput., 266(2015) 874-882.
- [12] M. Mursaleen, K. J. Ansari and A. Khan, Some Approximation Results by (p, q) -analogue of Bernstein-Stancu operators, Appl. Math. Comput., 264 (2015) 392-402 [Corrigendum: Appl. Math. Comput, 269 (2015) 744-746].
- [13] M. Mursaleen, Md. Nasiruzzaman and A. Nurgali, Some approximation results on Bernstein-Schurer operators defined by (p, q) -integers, J. Ineq. Appl., 2015 (2015): 249.
- [14] P.N. Sadjang, On the fundamental theorem of (p, q) -calculus and some (p, q) -Taylor formulas, arXiv:1309.3934v1 [math.QA], 2013.
- [15] V. Sahai and S. Yadav, Representations of two parameter quantum algebras and p, q -special functions, J. Math. Anal. Appl. 335(2007), 268-279.
- [16] W. Sudsutad, S.K. Ntouyas, J. Tariboon, Quantum integral inequalities for convex functions, J. Math. Inequal., 9 (2015), No. 3, 781-793.

- [17] J. Tariboon, S.K. Ntouyas, Quantum integral inequalities on finite intervals, *J. Inequal. Appl.* 2014 (2014) 121.
- [18] J. Tariboon, S.K. Ntouyas, Quantum calculus on finite intervals and applications to impulsive difference equations, *Adv. Differ. Equ.* 2013 (2013) 282.

E-mail address: ^αmevlutttunc@gmail.com

E-mail address: ^βesordulu@gmail.com

^{α,β}MUSTAFA KEMAL UNIVERSITY, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS, 31000, HATAY, TURKEY