

# A GENERAL DIVERGENCE MEASURE FOR MONOTONIC FUNCTIONS AND APPLICATIONS IN INFORMATION THEORY

S.S. DRAGOMIR

ABSTRACT. A general divergence measure for monotonic functions is introduced. Its connections with the  $f$ -divergence for convex functions are explored. The main properties are pointed out.

## 1. INTRODUCTION

Let  $(X, \mathcal{A})$  be a measurable space satisfying  $|\mathcal{A}| > 2$  and  $\mu$  be a  $\sigma$ -finite measure on  $(X, \mathcal{A})$ . Let  $\mathcal{P}$  be the set of all probability measures on  $(X, \mathcal{A})$  which are absolutely continuous with respect to  $\mu$ . For  $P, Q \in \mathcal{P}$ , let  $p = \frac{dP}{d\mu}$  and  $q = \frac{dQ}{d\mu}$  denote the *Radon-Nikodym* derivatives of  $P$  and  $Q$  with respect to  $\mu$ .

Two probability measures  $P, Q \in \mathcal{P}$  are said to be *orthogonal* and we denote this by  $Q \perp P$  if

$$P(\{q = 0\}) = Q(\{p = 0\}) = 1.$$

Let  $f : [0, \infty) \rightarrow (-\infty, \infty]$  be a convex function that is continuous at 0, i.e.,  $f(0) = \lim_{u \downarrow 0} f(u)$ .

In 1963, I. Csiszár [2] introduced the concept of  $f$ -divergence as follows.

**Definition 1.** Let  $P, Q \in \mathcal{P}$ . Then

$$(1.1) \quad I_f(Q, P) = \int_X p(x) f\left[\frac{q(x)}{p(x)}\right] d\mu(x),$$

is called the  $f$ -divergence of the probability distributions  $Q$  and  $P$ .

We now give some examples of  $f$ -divergences that are well-known and often used in the literature (see also [3]).

**1.1. The Class of  $\chi^\alpha$ -Divergences.** The  $f$ -divergences of this class, which is generated by the function  $\chi^\alpha$ ,  $\alpha \in [1, \infty)$ , defined by

$$\chi^\alpha(u) = |u - 1|^\alpha, \quad u \in [0, \infty)$$

have the form

$$(1.2) \quad I_f(Q, P) = \int_X p \left| \frac{q}{p} - 1 \right|^\alpha d\mu = \int_X p^{1-\alpha} |q - p|^\alpha d\mu.$$

From this class only the parameter  $\alpha = 1$  provides a distance in the topological sense, namely the *total variation distance*  $V(Q, P) = \int_X |q - p| d\mu$ . The most prominent special case of this class is, however, *Karl Pearson's  $\chi^2$ -divergence*.

---

*Date:* 13 January, 2004.

*2000 Mathematics Subject Classification.* 94Axx, 26D15, 26D10.

*Key words and phrases.*  $f$ -divergence, Convexity, Divergence measures, Monotonic functions.

1.2. **Dichotomy Class.** From this class, generated by the function  $f_\alpha : [0, \infty) \rightarrow \mathbb{R}$

$$f_\alpha(u) = \begin{cases} u - 1 - \ln u & \text{for } \alpha = 0; \\ \frac{1}{\alpha(1-\alpha)} [\alpha u + 1 - \alpha - u^\alpha] & \text{for } \alpha \in \mathbb{R} \setminus \{0, 1\}; \\ 1 - u + u \ln u & \text{for } \alpha = 1; \end{cases}$$

only the parameter  $\alpha = \frac{1}{2}$  ( $f_{\frac{1}{2}}(u) = 2(\sqrt{u} - 1)^2$ ) provides a distance, namely, the *Hellinger distance*

$$H(Q, P) = \left[ \int_X (\sqrt{q} - \sqrt{p})^2 d\mu \right]^{\frac{1}{2}}.$$

Another important divergence is the *Kullback-Leibler divergence* obtained for  $\alpha = 1$ ,

$$KL(Q, P) = \int_X q \ln \left( \frac{q}{p} \right) d\mu.$$

1.3. **Matsushita's Divergences.** The elements of this class, which is generated by the function  $\varphi_\alpha$ ,  $\alpha \in (0, 1]$  given by

$$\varphi_\alpha(u) := |1 - u^\alpha|^{\frac{1}{\alpha}}, \quad u \in [0, \infty),$$

are prototypes of metric divergences, providing the distances  $[I_{\varphi_\alpha}(Q, P)]^\alpha$ .

1.4. **Puri-Vineze Divergences.** This class is generated by the functions  $\Phi_\alpha$ ,  $\alpha \in [1, \infty)$  given by

$$\Phi_\alpha(u) := \frac{|1 - u|^\alpha}{(u + 1)^{\alpha-1}}, \quad u \in [0, \infty).$$

It has been shown in [4] that, this class provides the distances  $[I_{\Phi_\alpha}(Q, P)]^{\frac{1}{\alpha}}$ .

1.5. **Divergences of Arimoto-type.** This class is generated by the functions

$$\Psi_\alpha(u) := \begin{cases} \frac{\alpha}{\alpha-1} \left[ (1 + u^\alpha)^{\frac{1}{\alpha}} - 2^{\frac{1}{\alpha}-1} (1 + u) \right] & \text{for } \alpha \in (0, \infty) \setminus \{1\}; \\ (1 + u) \ln 2 + u \ln u - (1 + u) \ln (1 + u) & \text{for } \alpha = 1; \\ \frac{1}{2} |1 - u| & \text{for } \alpha = \infty. \end{cases}$$

It has been shown in [5] that, this class provides the distances  $[I_{\Psi_\alpha}(Q, P)]^{\min(\alpha, \frac{1}{\alpha})}$  for  $\alpha \in (0, \infty)$  and  $\frac{1}{2}V(Q, P)$  for  $\alpha = \infty$ .

## 2. SOME CLASSES OF NORMALISED FUNCTIONS

We denote by  $\mathcal{M}^\ddagger([0, \infty))$  the class of *monotonic nondecreasing functions* defined on  $[0, \infty)$  and by  $\mathcal{M}s([0, \infty))$  the class of *measurable functions* on  $[0, \infty)$ . We also consider  $\mathcal{L}e_1([0, \infty))$  the class of measurable functions  $g : [0, \infty) \rightarrow \mathbb{R}$  with the property that

$$(2.1) \quad g(t) \leq g(1) \leq g(s) \quad \text{for } 0 \leq t \leq 1 \leq s < \infty.$$

It is obvious that

$$(2.2) \quad \mathcal{M}^\ddagger([0, \infty)) \subsetneq \mathcal{L}e_1([0, \infty)),$$

and the inclusion (2.2) is strict.

We say that a function  $f : [0, \infty) \rightarrow \mathbb{R}$  is *normalised* if  $f(1) = 0$ . We denote by  $\mathcal{M}_{s_0}([0, \infty))$  the class of all normalised measurable functions defined on  $[0, \infty)$ . We also need the following classes of functions

$$\mathcal{C}o([0, \infty)) := \{f \in \mathcal{M}_{s_0}([0, \infty)) \mid f \text{ is continuous convex on } [0, \infty)\};$$

$$\mathcal{D}_0([0, \infty)) := \{f \in \mathcal{M}_{s_0}([0, \infty)) \mid f(t) = (t-1)g(t), \forall t \in [0, \infty), g \in \mathcal{M}^\#([0, \infty))\};$$

and

$$\mathcal{O}_0([0, \infty)) := \{f \in \mathcal{M}_{s_0}([0, \infty)) \mid f(t) = (t-1)g(t), \forall t \in [0, \infty), g \in \mathcal{L}e_1([0, \infty))\}.$$

From the definition of  $\mathcal{D}_0([0, \infty))$  and  $\mathcal{O}_0([0, \infty))$  and taking into account that the strict inclusion (2.2) holds, we deduce that

$$(2.3) \quad \mathcal{D}_0([0, \infty)) \subsetneq \mathcal{O}_0([0, \infty)),$$

and the inclusion is strict.

For the other two classes, we may state the following result.

**Lemma 1.** *We have the strict inclusion*

$$(2.4) \quad \mathcal{C}o([0, \infty)) \subsetneq \mathcal{D}_0([0, \infty)).$$

*Proof.* We will show that any continuous convex function  $f : [0, \infty) \rightarrow \mathbb{R}$  that is normalised may be represented as:

$$(2.5) \quad f(t) = (t-1)g(t) \text{ for any } t \in [0, \infty),$$

where  $g \in \mathcal{M}^\#([0, \infty))$ .

Now, let  $f \in \mathcal{C}o([0, \infty))$ . For  $\lambda \in [D_-f(1), D_+f(1)]$ , define

$$g_\lambda(t) := \begin{cases} \frac{f(t)}{t-1} & \text{if } t \in [0, 1) \cup (1, \infty), \\ \lambda & \text{if } t = 1. \end{cases}$$

We use the following well known result [1, p. 111]:

If  $\Psi$  is convex on  $(a, b)$  and  $a < s < t < u < b$ , then

$$(2.6) \quad \Psi(s, t) \leq \Psi(s, u) \leq \Psi(t, u),$$

where

$$\Psi(s, t) = \frac{\Psi(t) - \Psi(s)}{t - s}.$$

If  $\Psi$  is strictly convex on  $(a, b)$ , equality will not occur in (2.6).

If we apply the above result for  $0 < s < t < 1$ , then we can state

$$\frac{f(s)}{s-1} \leq \frac{f(t)}{t-1}.$$

Taking the limit over  $t \rightarrow 1, t < 1$ , we deduce

$$\frac{f(s)}{s-1} \leq D_-f(1)$$

showing that for  $0 < t < 1$ , we have  $g_\lambda(t) \leq \lambda$ .

Similarly, we may prove that for  $1 < t < \infty$ ,  $g_\lambda(t) \geq \lambda$ . If we use the same result for  $0 < t_1 < t_2 < 1$ , then we may write

$$\frac{f(t_1)}{t_1-1} \leq \frac{f(t_2)}{t_2-1},$$

which gives  $g_\lambda(t_1) \leq g_\lambda(t_2)$  for  $0 < t_1 < t_2 < 1$ .

In a similar fashion we can prove that for  $1 < t_1 < t_2 < \infty$ ,  $g_\lambda(t_1) \leq g_\lambda(t_2)$ , and thus we may conclude that the function  $g_\lambda$  is monotonic non-decreasing on the whole interval  $[0, \infty)$ .

If we consider now the function  $f(t) = (t-1)e^{\eta t}$ ,  $t \in [0, \infty)$ , we observe that  $f'(t) = (\eta t - 3)e^{\eta t}$ ,  $f''(t) = 8e^{\eta t}(2t-1)$  which shows that  $f$  is not convex on  $[0, \infty)$ . Obviously,  $f \in \mathcal{D}_0([0, \infty))$ , and thus the inclusion (2.4) is indeed strict. ■

**Remark 1.** If  $f \in \mathcal{D}_0([0, \infty))$  and  $g_1, g_2 \in \mathcal{M}^\#([0, \infty))$  are two functions with

$$f(t) = (t-1)g_1(t), \quad f(t) = (t-1)g_2(t)$$

for each  $t \in [0, \infty)$ , then we get

$$(t-1)[g_1(t) - g_2(t)] = 0$$

for any  $t \in [0, \infty)$  showing that  $g_1(t) = g_2(t)$  for each  $t \in [0, 1) \cup (1, \infty)$ . They may have different values in  $t = 1$ .

### 3. SOME FUNDAMENTAL PROPERTIES OF $f$ -DIVERGENCE FOR $f \in \mathcal{Co}([0, \infty))$

For  $f \in \mathcal{Co}([0, \infty))$  we obtain the  $*$ -conjugate function of  $f$  by

$$f^*(u) = uf\left(\frac{1}{u}\right), \quad u \in (0, \infty).$$

It is also known that if  $f \in \mathcal{Co}([0, \infty))$ , then  $f^* \in \mathcal{Co}([0, \infty))$ .

The following two theorems contain the most basic properties of  $f$ -divergences. For their proof we refer the reader to Chapter 1 of [6] (see also [3]).

**Theorem 1** (Uniqueness and Symmetry Theorem). *Let  $f, f_1$  be continuous convex on  $[0, \infty)$ .*

(i) *We have*

$$I_{f_1}(Q, P) = I_f(Q, P),$$

*for any  $P, Q \in \mathcal{P}$  if and only if there exists a constant  $c \in \mathbb{R}$  such that*

$$f_1(u) = f(u) + c(u-1),$$

*for any  $u \in [0, \infty)$ ;*

(ii) *We have*

$$I_{f^*}(Q, P) = I_f(Q, P),$$

*for any  $P, Q \in \mathcal{P}$  if and only if there exists a constant  $d \in \mathbb{R}$  such that*

$$f^*(u) = f(u) + d(c-1),$$

*for any  $u \in [0, \infty)$ .*

**Theorem 2** (Range of Values Theorem). *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a continuous convex function on  $[0, \infty)$ .*

*For any  $P, Q \in \mathcal{P}$ , we have the double inequality*

$$(3.1) \quad f(1) \leq I_f(Q, P) \leq f(0) + f^*(0).$$

(i) *If  $P = Q$ , then the equality holds in the first part of (3.1).*

*If  $f$  is strictly convex at 1, then the equality holds in the first part of (3.1) if and only if  $P = Q$ ;*

- (ii) If  $Q \perp P$ , then the equality holds in the second part of (3.1).  
 If  $f(0) + f^*(0) < \infty$ , then equality holds in the second part of (3.1) if and only if  $Q \perp P$ .

Define the function  $\tilde{f} : (0, \infty) \rightarrow \mathbb{R}$ ,  $\tilde{f}(u) = \frac{1}{2}(f(u) + f^*(u))$ . The following result is a refinement of the second inequality in Theorem 2 (see [3, Theorem 3]).

**Theorem 3.** Let  $f \in \mathcal{C}o([0, \infty))$  with  $f(0) + f^*(0) < \infty$ . Then

$$(3.2) \quad 0 \leq I_f(Q, P) \leq \tilde{f}(0) V(Q, P)$$

for any  $Q, P \in \mathcal{P}$ .

#### 4. A GENERAL DIVERGENCE MEASURE

If  $f : [0, \infty) \rightarrow \mathbb{R}$  is a general measurable function, then we may define the  $f$ -divergence in the same way, i.e., if  $P, Q \in \mathcal{P}$ , then

$$I_f(Q, P) = \int_X p(x) f \left[ \frac{q(x)}{p(x)} \right] d\mu(x).$$

For a measurable function  $g : [0, \infty) \rightarrow \mathbb{R}$ , we may also define the  $\delta$ -divergence by the formula

$$\delta_g(Q, P) = \int_X [q(x) - p(x)] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x).$$

It is obvious that the  $\delta$ -divergence of a function  $g$  may be seen as the  $f$ -divergence of the function  $f$ , where  $f(t) = (t-1)g(t)$  for  $t \in [0, \infty)$ .

If  $f \in \mathcal{C}o([0, \infty))$  and since  $f(t) = (t-1)g_\lambda(t)$ ,  $t \in [0, \infty)$ , we have

$$(4.1) \quad g_\lambda(t) := \begin{cases} \frac{f(t)}{t-1} & \text{if } t \in [0, 1) \cup (1, \infty), \\ \lambda & \text{if } t = 1; \end{cases}$$

and  $\lambda \in [D_-f(1), D_+f(1)]$ , shows that for any  $f \in \mathcal{C}o([0, \infty))$  we have

$$(4.2) \quad I_f(Q, P) = \delta_{g_\lambda}(Q, P) \quad \text{for any } P, Q \in \mathcal{P},$$

i.e., the  $f$ -divergence for any normalised continuous convex function  $f : [0, \infty) \rightarrow \mathbb{R}$  may be seen as the  $\delta$ -divergence of the function  $g_\lambda$  defined by (4.1).

In what follows, we point out some fundamental properties of the  $\delta$ -divergence.

**Theorem 4.** Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a measurable function on  $[0, \infty)$  and  $P, Q \in \mathcal{P}$ . If there exists the constants  $m, M$  with

$$(4.3) \quad -\infty < m \leq g \left[ \frac{q(x)}{p(x)} \right] \leq M < \infty$$

for  $\mu$ -a.e.  $x \in X$ , then we have the inequality

$$(4.4) \quad |\delta_g(Q, P)| \leq \frac{1}{2} (M - m) V(Q, P).$$

*Proof.* We observe that the following identity holds true

$$(4.5) \quad \delta_g(Q, P) = \int_X [q(x) - p(x)] \left[ g \left[ \frac{q(x)}{p(x)} \right] - \frac{m+M}{2} \right] d\mu(x)$$

By (4.3), we deduce that

$$\left| g \left[ \frac{q(x)}{p(x)} \right] - \frac{m+M}{2} \right| \leq \frac{1}{2} (M-m)$$

for  $\mu$ -a.e.  $x \in X$ .

Taking the modulus in (4.5) we deduce

$$\begin{aligned} |\delta_g(Q, P)| &\leq \int_X |q(x) - p(x)| \left| g \left[ \frac{q(x)}{p(x)} - \frac{m+M}{2} \right] \right| d\mu(x) \\ &\leq \frac{1}{2} (M-m) \int_X |q(x) - p(x)| d\mu(x) \\ &= \frac{1}{2} (M-m) V(Q, P) \end{aligned}$$

and the inequality (4.4) is proved. ■

The following corollary is a natural consequence of the above theorem.

**Corollary 1.** *Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a measurable function on  $[0, \infty)$ . If*

$$m := \operatorname{ess\,inf}_{t \in [0, \infty)} g(t) > -\infty, \quad M := \operatorname{ess\,sup}_{t \in [0, \infty)} g(t) < \infty,$$

then for any  $P, Q \in \mathcal{P}$ , we have the inequality

$$(4.6) \quad |\delta_g(Q, P)| \leq \frac{1}{2} (M-m) V(Q, P).$$

**Remark 2.** *We know that, if  $f : [0, \infty) \rightarrow \mathbb{R}$  is a normalised continuous convex function and if  $\lim_{t \downarrow 0} f^*(t) = \lim_{u \downarrow 0} [uf(\frac{1}{u})] =: f^*(0)$ , then we have the inequality [Theorem 2.3]*

$$(4.7) \quad I_f(Q, P) \leq \frac{f(0) + f^*(0)}{2} V(Q, P),$$

for any  $P, Q \in \mathcal{P}$ . We can prove this inequality by the use of Corollary 1 as follows.

We have

$$I_f(Q, P) = \delta_{g_\lambda}(Q, P),$$

where

$$g_\lambda(t) := \begin{cases} \frac{f(t)}{t-1} & \text{if } t \in [0, 1) \cup (1, \infty), \\ \lambda & \text{if } t = 1, \end{cases}$$

where  $\lambda \in [D_-f(1), D_+f(1)]$  and  $g_\lambda \in \mathcal{M}^\#([0, \infty))$ . We observe that for any  $t \in [0, \infty)$ , we have

$$g_\lambda(t) \geq \lim_{t \rightarrow 0^+} g_\lambda(t) = -f(0) = m > -\infty$$

and

$$\begin{aligned} g_\lambda(t) &\leq \lim_{t \rightarrow +\infty} g_\lambda(t) = \lim_{t \rightarrow +\infty} \frac{f(t)}{t-1} = \lim_{u \rightarrow 0^+} \left[ \frac{f(\frac{1}{u})}{\frac{1}{u}-1} \right] \\ &= \lim_{u \rightarrow 0^+} \left[ \frac{uf(\frac{1}{u})}{1-u} \right] = f^*(0) = M < \infty. \end{aligned}$$

Applying Corollary 1 for  $m = -f(0)$  and  $M = f^*(0)$ , we deduce the desired inequality (4.7).

The following result also holds.

**Theorem 5.** *Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a measurable function on  $[0, \infty)$  and  $P, Q \in \mathcal{P}$ . If there exists a constant  $K$  with  $K > 0$  such that*

$$(4.8) \quad \left| g\left(\frac{q(x)}{p(x)}\right) - g(1) \right| \leq K \left| \frac{q(x)}{p(x)} - 1 \right|^\alpha,$$

for  $\mu$ -a.e.  $x \in X$ , where  $\alpha \in (0, \infty)$  is a given number, then we have the inequality

$$(4.9) \quad |\delta_g(Q, P)| \leq K I_{\chi^{\alpha+1}}(Q, P).$$

*Proof.* We observe that the following identity holds true

$$(4.10) \quad \delta_g(Q, P) = \int_X [q(x) - p(x)] \left[ g\left[\frac{q(x)}{p(x)}\right] - g(1) \right] d\mu(x).$$

Taking the modulus in (4.10) and using the condition (4.8), we have successively

$$\begin{aligned} |\delta_g(Q, P)| &\leq \int_X |q(x) - p(x)| \left| g\left[\frac{q(x)}{p(x)}\right] - g(1) \right| d\mu(x) \\ &\leq K \int_X [p(x)]^{-\alpha} |q(x) - p(x)|^{\alpha+1} d\mu(x) \\ &\leq K I_{\chi^{\alpha+1}}(Q, P) \end{aligned}$$

and the inequality (4.9) is obtained. ■

The following corollary holds.

**Corollary 2.** *Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a measurable function on  $[0, \infty)$  with the property that there exists a constant  $K$  with the property that*

$$(4.11) \quad |g(t) - g(1)| \leq K |t - 1|^\alpha,$$

for a.e.  $t \in [0, \infty)$ , where  $\alpha > 0$  is a given number. Then for any  $P, Q \in \mathcal{P}$ , we have the inequality

$$(4.12) \quad |\delta_g(Q, P)| \leq K I_{\chi^{\alpha+1}}(Q, P).$$

**Remark 3.** *If the function  $g : [0, \infty) \rightarrow \mathbb{R}$  is Hölder continuous with a constant  $H > 0$  and  $\beta \in (0, 1]$ , i.e.,*

$$|g(t) - g(s)| \leq H |t - s|^\beta,$$

for any  $t, s \in [0, \infty)$ , then obviously (4.7) holds with  $K = H$  and  $\alpha = \beta$ .

*If  $g : [0, \infty) \rightarrow \mathbb{R}$  is Lipschitzian with the constant  $L > 0$ , i.e.,*

$$|g(t) - g(s)| \leq L |t - s|,$$

for any  $t, s \in [0, \infty)$ , then

$$(4.13) \quad |\delta_g(Q, P)| \leq K I_{\chi^2}(Q, P),$$

for any  $P, Q \in \mathcal{P}$ .

*Finally, if  $g$  is locally absolutely continuous and the derivative  $g' : [0, \infty) \rightarrow \mathbb{R}$  is essentially bounded, i.e.,  $\|g'\|_{[0, \infty), \infty} := \text{ess sup}_{t \in [0, \infty)} |g'(t)| < \infty$ , then we have the inequality*

$$(4.14) \quad |\delta_g(Q, P)| \leq \|g'\|_{[0, \infty), \infty} I_{\chi^2}(Q, P),$$

for any  $P, Q \in \mathcal{P}$ .

The following result concerning  $f$ -divergences for  $f$  convex functions holds.

**Theorem 6.** Let  $f : [0, \infty] \rightarrow \mathbb{R}$  be a continuous convex function on  $[0, \infty)$ . If  $\lambda \in [D_-f(1), D_+f(1)]$  ( $\lambda = f'(1)$  if  $f$  is differentiable at  $t = 1$ ), and there exists a constant  $K > 0$  and  $\alpha > 0$  such that

$$(4.15) \quad |f(t) - \lambda(t-1)| \leq K|t-1|^{\alpha+1},$$

for any  $t \in [0, \infty)$ , then we have the inequality

$$(4.16) \quad 0 \leq I_f(Q, P) \leq KI_{\chi^{\alpha+1}}(Q, P),$$

for any  $P, Q \in \mathcal{P}$ .

*Proof.* We have

$$I_f(Q, P) = \int_X [q(x) - p(x)] g_\lambda \left[ \frac{p(x)}{q(x)} \right] d\mu(x) = \delta_{g_\lambda}(Q, P),$$

where

$$g_\lambda(t) := \begin{cases} \frac{f(t)}{t-1} & \text{if } t \in [0, 1) \cup (1, \infty), \\ \lambda & \text{if } t = 1, \end{cases}$$

and  $\lambda \in [D_-f(1), D_+f(1)]$ .

Applying Corollary 2 for  $g_\lambda$ , we deduce the desired result. ■

## 5. THE POSITIVITY OF $\delta$ -DIVERGENCE FOR $g \in \mathcal{M}^\ddagger([0, \infty))$

The following result holds.

**Theorem 7.** If  $g \in \mathcal{M}^\ddagger([0, \infty))$ , then  $\delta_g(Q, P) \geq 0$  for any  $P, Q \in \mathcal{P}$ .

*Proof.* We use the identity

$$(5.1) \quad \begin{aligned} \delta_g(Q, P) &= \int_X [q(x) - p(x)] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\ &= \int_X p(x) \left[ \frac{q(x)}{p(x)} - 1 \right] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\ &= \frac{1}{2} \int_X \int_X p(x) p(y) \left[ \frac{q(x)}{p(x)} - \frac{q(y)}{p(y)} \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] d\mu(x) d\mu(y). \end{aligned}$$

Since  $g \in \mathcal{M}^\ddagger([0, \infty))$ , then for any  $t, s \in [0, \infty)$ , we have

$$(t-s)(g(t) - g(s)) \geq 0$$

giving that

$$\left[ \frac{q(x)}{p(x)} - \frac{q(y)}{p(y)} \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] \geq 0$$

for any  $x, y \in X$ .

Using the representation (5.1), we deduce the desired result. ■

The following corollary is a natural consequence of the above result.

**Corollary 3.** If  $f \in \mathcal{D}_0([0, \infty))$ , then  $I_f(Q, P) \geq 0$  for any  $P, Q \in \mathcal{P}$ .



*Proof.* If  $f \in \mathcal{D}_0([0, \infty))$ , then there exists a  $g \in \mathcal{M}^\ddagger([0, \infty))$  such that  $f(t) = (t-1)g(t)$  for any  $t \in [0, \infty)$ . Then

$$\begin{aligned} I_f(Q, P) &= \int_X p(x) f\left[\frac{q(x)}{p(x)}\right] d\mu(x) \\ &= \int_X p(x) \left[\frac{q(x)}{p(x)} - 1\right] g\left[\frac{q(x)}{p(x)}\right] d\mu(x) \\ &= \delta_g(Q, P) \geq 0, \end{aligned}$$

and the proof is completed. ■

In fact, the following improvement of Theorem 7 holds.

**Theorem 8.** *If  $g \in \mathcal{M}^\ddagger([0, \infty))$ , then*

$$(5.2) \quad \delta_g(Q, P) \geq |\delta_{|g|}(Q, P)| \geq 0,$$

for any  $P, Q \in \mathcal{P}$ .

*Proof.* Since  $g$  is monotonic nondecreasing, we have

$$(5.3) \quad \begin{aligned} &\left[\frac{q(x)}{p(x)} - \frac{q(y)}{p(y)}\right] \left[g\left[\frac{q(x)}{p(x)}\right] - g\left[\frac{q(y)}{p(y)}\right]\right] \\ &= \left|\left(\frac{q(x)}{p(x)} - \frac{q(y)}{p(y)}\right) \left(g\left[\frac{q(x)}{p(x)}\right] - g\left[\frac{q(y)}{p(y)}\right]\right)\right| \\ &\geq \left|\left(\frac{q(x)}{p(x)} - \frac{q(y)}{p(y)}\right) \left(\left|g\left[\frac{q(x)}{p(x)}\right]\right| - \left|g\left[\frac{q(y)}{p(y)}\right]\right|\right)\right| \end{aligned}$$

for any  $x, y \in X$ .

Multiplying (5.3) by  $p(x)p(y) \geq 0$  and integrating on  $X^2$ , we deduce

$$\begin{aligned} &\int_X \int_X p(x)p(y) \left(\frac{q(x)}{p(x)} - \frac{q(y)}{p(y)}\right) \left[g\left[\frac{q(x)}{p(x)}\right] - g\left[\frac{q(y)}{p(y)}\right]\right] d\mu(x) d\mu(y) \\ &\geq \left|\int_X \int_X p(x)p(y) \left(\frac{q(x)}{p(x)} - \frac{q(y)}{p(y)}\right) \left(g\left[\frac{q(x)}{p(x)}\right] - g\left[\frac{q(y)}{p(y)}\right]\right) d\mu(x) d\mu(y)\right|. \end{aligned}$$

Using the representation (5.1) and the same identity for  $|g|$ , we deduce the desired inequality (5.2). ■

Before we point out other possible refinements for the positivity inequality  $\delta_g(Q, P) \geq 0$ , where  $g \in \mathcal{M}^\ddagger([0, \infty))$ , we need the following divergence measure as well:

$$\bar{\delta}_h(Q, P) := \int_X |q(x) - p(x)| h\left[\frac{q(x)}{p(x)}\right] d\mu(x)$$

which will be called the *absolute  $\delta$ -divergence* generated by the function  $h : [0, \infty) \rightarrow \mathbb{R}$  that is assumed to be measurable on  $[0, \infty)$ .

The following result holds.

**Theorem 9.** *If  $g \in \mathcal{M}^\ddagger([0, \infty))$ , then*

$$(5.4) \quad \begin{aligned} &\delta_g(Q, P) \\ &\geq \max\left\{|\bar{\delta}_g(Q, P) - V(Q, P) I_g(Q, P)|, |\bar{\delta}_{|g|}(Q, P) - V(Q, P) I_{|g|}(Q, P)|\right\} \geq 0, \end{aligned}$$

for any  $P, Q \in \mathcal{P}$ .

*Proof.* Since  $g$  is monotonic, we have

$$(5.5) \quad \begin{aligned} & \left( \frac{q(x)}{p(x)} - \frac{q(y)}{p(y)} \right) \left( g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right) \\ &= \left| \left[ \left( \frac{q(x)}{p(x)} - 1 \right) - \left( \frac{q(y)}{p(y)} - 1 \right) \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] \right| \\ &\geq \begin{cases} \left| \left[ \left| \frac{q(x)}{p(x)} - 1 \right| - \left| \frac{q(y)}{p(y)} - 1 \right| \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] \right| \\ \left| \left[ \left| \frac{q(x)}{p(x)} - 1 \right| - \left| \frac{q(y)}{p(y)} - 1 \right| \right] \left[ \left| g \left[ \frac{q(x)}{p(x)} \right] \right| - \left| g \left[ \frac{q(y)}{p(y)} \right] \right| \right] \right| \end{cases} \end{aligned}$$

for any  $x, y \in X$ .

If we multiply (5.5) by  $p(x)p(y) \geq 0$  and integrate, we deduce

$$(5.6) \quad \begin{aligned} & \int_X \int_X p(x)p(y) \left( \frac{q(x)}{p(x)} - \frac{q(y)}{p(y)} \right) \left( g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right) d\mu(x) d\mu(y) \\ &\geq \begin{cases} \left| \int_X \int_X p(x)p(y) \left[ \left| \frac{q(x)}{p(x)} - 1 \right| - \left| \frac{q(y)}{p(y)} - 1 \right| \right] \right. \\ \quad \left. \times \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] d\mu(x) d\mu(y) \right| \\ \left| \int_X \int_X p(x)p(y) \left[ \left| \frac{q(x)}{p(x)} - 1 \right| - \left| \frac{q(y)}{p(y)} - 1 \right| \right] \right. \\ \quad \left. \times \left[ \left| g \left[ \frac{q(x)}{p(x)} \right] \right| - \left| g \left[ \frac{q(y)}{p(y)} \right] \right| \right] d\mu(x) d\mu(y) \right| \end{cases} \end{aligned}$$

for any  $x, y \in X$ .

Now, observe that

$$\begin{aligned} & \int_X \int_X p(x)p(y) \left[ \left| \frac{q(x)}{p(x)} - 1 \right| - \left| \frac{q(y)}{p(y)} - 1 \right| \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g \left[ \frac{q(y)}{p(y)} \right] \right] d\mu(x) d\mu(y) \\ &= \int_X \int_X p(x)p(y) \left[ \left| \frac{q(x)}{p(x)} - 1 \right| g \left[ \frac{q(x)}{p(x)} \right] + \left| \frac{q(y)}{p(y)} - 1 \right| g \left[ \frac{q(y)}{p(y)} \right] \right] d\mu(x) d\mu(y) \\ &\quad - \int_X \int_X p(x)p(y) \left[ \left| \frac{q(x)}{p(x)} - 1 \right| g \left[ \frac{q(y)}{p(y)} \right] + \left| \frac{q(y)}{p(y)} - 1 \right| g \left[ \frac{q(x)}{p(x)} \right] \right] d\mu(x) d\mu(y) \\ &= 2 \int_X p(y) d\mu(y) \int_X p(x) \left| \frac{q(x)}{p(x)} - 1 \right| g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\ &\quad - 2 \int_X p(x) \left| \frac{q(x)}{p(x)} - 1 \right| d\mu(x) \int_X p(y) g \left[ \frac{q(y)}{p(y)} \right] d\mu(y) \\ &= 2 [\bar{\delta}_g(Q, P) - V(Q, P) I_g(Q, P)], \end{aligned}$$

and a similar identity holds for the quantity in the second branch of (5.6).

Finally, using the representation (5.1), we deduce the desired inequality (5.4). ■

## 6. THE POSITIVITY OF $\delta$ -DIVERGENCE FOR $g \in \mathcal{L}e_1([0, \infty))$

The following result extending the positivity of  $\delta$ -divergence for monotonic functions, holds.

**Theorem 10.** *If  $g \in \mathcal{L}e_1([0, \infty))$ , then  $\delta_g(Q, P) \geq 0$  for any  $P, Q \in \mathcal{P}$ .*

*Proof.* We use the identity

$$\begin{aligned}
 (6.1) \quad \delta_g(Q, P) &= \int_X [q(x) - p(x)] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\
 &= \int_X p(x) \left[ \frac{q(x)}{p(x)} - 1 \right] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\
 &= \int_X p(x) \left[ \frac{q(x)}{p(x)} - 1 \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g(1) \right] d\mu(x).
 \end{aligned}$$

Since  $g \in \mathcal{L}e_1([0, \infty))$ , then for any  $t \in [0, \infty)$  we have

$$(t - 1)[g(t) - g(1)] \geq 0$$

giving that

$$\left( \frac{q(x)}{p(x)} - 1 \right) \left[ g \left[ \frac{q(x)}{p(x)} \right] - g(1) \right] \geq 0$$

for any  $x \in X$ .

Using the representation (6.1), we deduce the desired result. ■

**Corollary 4.** *If  $f \in \mathcal{O}_0([0, \infty))$ , then  $I_f(Q, P) \geq 0$  for any  $P, Q \in \mathcal{P}$ .*

*Proof.* If  $f \in \mathcal{O}_0([0, \infty))$ , then there exists a  $g \in \mathcal{L}e_1([0, \infty))$  such that  $f(t) = (t - 1)g(t)$  for any  $t \in [0, \infty)$ . Then

$$\begin{aligned}
 I_f(Q, P) &= \int_X p(x) f \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\
 &= \int_X p(x) \left[ \frac{q(x)}{p(x)} - 1 \right] g \left[ \frac{q(x)}{p(x)} \right] d\mu(x) \\
 &= \delta_g(Q, P) \geq 0,
 \end{aligned}$$

and the proof is completed. ■

The following improvement of Theorem 10 holds.

**Theorem 11.** *If  $g \in \mathcal{L}e_1([0, \infty))$ , then*

$$(6.2) \quad \delta_g(Q, P) \geq |\delta_{|g|}(Q, P)| \geq 0$$

for any  $P, Q \in \mathcal{P}$ .

*Proof.* Since  $g \in \mathcal{L}e_1([0, \infty))$ , we obviously have

$$\begin{aligned}
 (6.3) \quad &\left[ \frac{q(x)}{p(x)} - 1 \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g(1) \right] \\
 &= \left| \left( \frac{q(x)}{p(x)} - 1 \right) \left( g \left[ \frac{q(x)}{p(x)} \right] - g(1) \right) \right| \\
 &\geq \left| \left( \frac{q(x)}{p(x)} - 1 \right) \left( \left| g \left[ \frac{q(x)}{p(x)} \right] \right| - |g(1)| \right) \right|.
 \end{aligned}$$

Multiplying (6.3) by  $p(x) \geq 0$  and integrating on  $X$ , we have

$$\begin{aligned} & \int_X p(x) \left[ \frac{q(x)}{p(x)} - 1 \right] \left[ g \left[ \frac{q(x)}{p(x)} \right] - g(1) \right] d\mu(x) \\ &= \int_X p(x) \left| \left( \frac{q(x)}{p(x)} - 1 \right) \left( \left| g \left[ \frac{q(x)}{p(x)} \right] \right| - |g(1)| \right) \right| d\mu(x) \\ &\geq \left| \int_X p(x) \left( \frac{q(x)}{p(x)} - 1 \right) \left( \left| g \left[ \frac{q(x)}{p(x)} \right] \right| - |g(1)| \right) d\mu(x) \right| \\ &= |\delta_{|g|}(Q, P)|, \end{aligned}$$

and the inequality (6.2) is proved. ■

## 7. BOUNDS IN TERMS OF THE $\chi^2$ -DIVERGENCE

The following result may be stated.

**Theorem 12.** *Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a differentiable function such that there exists the constants  $\gamma, \Gamma \in \mathbb{R}$  with*

$$(7.1) \quad \gamma \leq g'(t) \leq \Gamma \quad \text{for any } t \in (0, \infty).$$

*Then we have the inequality*

$$(7.2) \quad \gamma D_{\chi^2}(Q, P) \leq \delta_g(Q, P) \leq \Gamma D_{\chi^2}(Q, P),$$

*for any  $P, Q \in \mathcal{P}$ .*

*Proof.* Consider the auxiliary function  $h_\gamma : [0, \infty) \rightarrow \mathbb{R}$ ,  $h_\gamma(t) := g(t) - \gamma(t-1)$ . Obviously,  $h_\gamma$  is differentiable on  $(0, \infty)$  and since, by (7.1),

$$h'_\gamma(t) = g'(t) - \gamma \geq 0$$

it follows that  $h_\gamma$  is monotonic nondecreasing on  $[0, \infty)$ .

Applying Theorem 7, we deduce

$$\delta_{h_\gamma}(Q, P) \geq 0 \quad \text{for any } P, Q \in \mathcal{P}$$

and since

$$\begin{aligned} \delta_{h_\gamma}(Q, P) &= \delta_{g-\gamma(\cdot-1)}(Q, P) \\ &= \int_X [q(x) - p(x)] \left[ g \left[ \frac{q(x)}{p(x)} \right] - \gamma \left[ \frac{q(x)}{p(x)} - 1 \right] \right] d\mu(x) \\ &= \delta_g(Q, P) - \gamma D_{\chi^2}(Q, P), \end{aligned}$$

then the first inequality in (7.2) is proved.

The second inequality may be proven in a similar manner by using the auxiliary function  $h_\Gamma : [0, \infty) \rightarrow \mathbb{R}$ ,  $h_\Gamma(t) := \Gamma(t-1) - g(t)$ . ■

The following corollary is a natural application of the above theorem.

**Corollary 5.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a differentiable convex function on  $(0, \infty)$  with  $f(1) = 0$ . If there exist the constants  $\gamma, \Gamma \in \mathbb{R}$  with the property that:*

$$(7.3) \quad \gamma(t-1)^2 + f(t) \leq f'(t)(t-1) \leq f(t) + \Gamma(t-1)^2$$

*for any  $t \in (0, \infty)$ , then we have the inequality:*

$$(7.4) \quad \gamma D_{\chi^2}(Q, P) \leq I_f(Q, P) \leq \Gamma D_{\chi^2}(Q, P)$$

*for any  $P, Q \in \mathcal{P}$ .*

*Proof.* We know that for any  $P, Q \in \mathcal{P}$ , we have (see for example (4.2)):

$$I_f(Q, P) = \delta_{g_{f'(1)}}(Q, P),$$

where

$$g_{f'(1)} = \begin{cases} \frac{f(t)}{t-1} & \text{if } t \in [0, 1) \cup (1, \infty), \\ f'(1) & \text{if } t = 1. \end{cases}$$

We observe that, by the hypothesis of the corollary,  $g_{f'(1)}$  is differentiable on  $(0, \infty)$  and

$$g'_{f'(1)}(t) = \frac{f'(t)(t-1) - f(t)}{(t-1)^2}$$

for any  $t \in (0, 1) \cup (1, \infty)$ .

Using (7.3), we deduce that

$$\gamma \leq g'_{f'(1)}(t) \leq \Gamma$$

for  $t \in (0, \infty)$ , and applying Theorem 12 above, for  $g = g_{f'(1)}$ , we deduce the desired inequality (7.4). ■

## 8. BOUNDS IN TERMS OF THE $J$ -DIVERGENCE

We recall that the *Jeffreys divergence* (or  $J$ -divergence for short) is defined as

$$(8.1) \quad J(Q, P) := \int_X [q(x) - p(x)] \ln \left[ \frac{q(x)}{p(x)} \right] d\mu(x),$$

where  $P, Q \in \mathcal{P}$ .

The following result holds.

**Theorem 13.** *Let  $g : [0, \infty) \rightarrow \mathbb{R}$  be a differentiable function such that there exists the constants  $\phi, \Phi \in \mathbb{R}$  with*

$$(8.2) \quad \phi \leq tg'(t) \leq \Phi \quad \text{for any } t \in (0, \infty).$$

*Then we have the inequality*

$$(8.3) \quad \phi J(Q, P) \leq \delta_g(Q, P) \leq \Phi J(Q, P),$$

*for any  $P, Q \in \mathcal{P}$ .*

*Proof.* Consider the auxiliary function  $h_\phi : [0, \infty) \rightarrow \mathbb{R}$ ,  $h_\phi(t) := g(t) - \phi \ln t$ . Obviously,  $h_\phi$  is differentiable on  $(0, \infty)$  and, by (8.2),

$$h'_\phi(t) = g'(t) - \frac{\phi}{t} = \frac{1}{t} [tg'(t) - \phi] \geq 0,$$

for any  $t \in (0, \infty)$ , showing that the function is monotonic nondecreasing on  $(0, \infty)$ .

Applying Theorem 7, we deduce

$$\delta_{h_\phi}(Q, P) \geq 0 \quad \text{for any } P, Q \in \mathcal{P}$$

and since

$$\begin{aligned} \delta_{h_\phi}(Q, P) &= \delta_{g-\phi \ln(\cdot)}(Q, P) \\ &= \int_X [q(x) - p(x)] \left[ g \left[ \frac{q(x)}{p(x)} \right] - \phi \ln \left[ \frac{q(x)}{p(x)} \right] \right] d\mu(x) \\ &= \delta_g(Q, P) - \phi J(Q, P), \end{aligned}$$

then the first inequality in (8.3) is proved.

The second inequality may be proven in a similar manner by using the auxiliary function  $h_\Phi : [0, \infty) \rightarrow \mathbb{R}$ ,  $h_\Phi(t) := \Phi \ln t - g(t)$ . ■

The following corollary is a natural application of the above theorem.

**Corollary 6.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a differentiable convex function on  $(0, \infty)$  with  $f(1) = 0$ . If there exist the constants  $\phi, \Phi \in \mathbb{R}$  with the property that:*

$$(8.4) \quad \phi(t-1)^2 + tf(t) \leq t(t-1)f'(t) \leq tf(t) + \Phi(t-1)^2$$

for any  $t \in (0, \infty)$ , then we have the inequality:

$$(8.5) \quad \phi J(Q, P) \leq I_f(Q, P) \leq \Phi J(Q, P)$$

for any  $P, Q \in \mathcal{P}$ .

The proof is similar to the one in Corollary 5 and we omit the details.

#### REFERENCES

- [1] G. de BARRA, *Measure Theory and Integration*, Ellis Horwood Ltd., 1981.
- [2] I. CSISZÁR, Eine informationstheoretische Ungleichung und ihre Anwendung auf den Beweis der Ergodizität von Markoffschen Ketten. (German) *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **8** (1963) 85–108.
- [3] P. CERONE, S.S. DRAGOMIR and F. ÖSTERREICHER, Bounds on extended  $f$ -divergences for a variety of classes, *RGMA Res. Rep. Coll.* **6**(2003), No.1, Article 5. [ONLINE: <http://rgmia.vu.edu.au/v6n1.html>].
- [4] P. KAFKA, F. ÖSTERREICHER and I. VINCZE, On powers of  $f$ -divergence defining a distance, *Studia Sci. Math. Hungar.*, **26** (1991), 415-422.
- [5] F. ÖSTERREICHER and I. VAJDA, A new class of metric divergences on probability spaces and its applicability in statistics. *Ann. Inst. Statist. Math.* **55** (2003), no. 3, 639–653.
- [6] F. LIESE and I. VAJDA, *Convex Statistical Distances*, Teubner – Texte zur Mathematik, Band **95**, Leipzig, 1987.

SCHOOL OF COMPUTER SCIENCE AND MATHEMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY,  
PO Box 14428, MCMC 8001, VICTORIA, AUSTRALIA.

*E-mail address:* sever.dragomir@vu.edu.au

*URL:* <http://rgmia.vu.edu.au/SSDragomirWeb.html>