

A REMARK ON THE SUM OF RECIPROCAL OF IMAGINARY PARTS OF ZETA ZEROS

MEHDI HASSANI

ABSTRACT. In this note, we give some explicit upper and lower bounds for the summation $\sum_{0 < \gamma \leq T} \frac{1}{\gamma}$, where γ is the imaginary part of nontrivial zeros $\rho = \beta + i\gamma$ of $\zeta(s)$, that is a zero with $0 \leq \beta \leq 1$. In a research report appeared in “*RGMIA Research Report Collection*, 9(2), Article 15, 2006”, we have given some bounds, but here we give the following more clean ones:

$$\frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \frac{15}{250} < \sum_{0 < \gamma \leq T} \frac{1}{\gamma} < \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \frac{109}{250},$$

where the left hand side holds for $T \geq 2$ and the right hand side holds for $T \geq 2.222$.

1. INTRODUCTION

The Riemann zeta function is defined for $\Re(s) > 1$ by $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$ and extended by analytic continuation to the complex plan with one singularity at $s = 1$; in fact a simple pole with residues 1. The functional equation for this function in symmetric form, is $\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \zeta(1-s)$, where $\Gamma(s) = \int_0^{\infty} e^{-t} t^{s-1} dt$ is a meromorphic function of the complex variable s , with simple poles at $s = 0, -1, -2, \dots$ (see [3]). By this equation, trivial zeros of $\zeta(s)$ are $s = -2, -4, -6, \dots$. Also, it implies symmetry of nontrivial zeros (other zeros $\rho = \beta + i\gamma$ which have the property $0 \leq \beta \leq 1$) according to the line $\Re(s) = \frac{1}{2}$. The summation

$$\mathcal{A}(T) = \sum_{0 < \gamma \leq T} \frac{1}{\gamma},$$

where γ is the imaginary part of nontrivial zeros appears in some explicit approximation of primes, and having some explicit approximations of it can be useful for careful computations. This is a summation over imaginary part of zeta zeros, and for approximating such summations we use Stieljes integral and integrating by parts; let $N(T)$ be the number of zeros ρ of $\zeta(s)$ with $0 < \Im(\rho) \leq T$ and $0 \leq \Re(\rho) \leq 1$. Then, supposing $1 < U \leq V$ and

2000 *Mathematics Subject Classification*. 11S40.

Key words and phrases. The Riemann zeta function.

$\Phi(t) \in C^1(U, V)$ to be non-negative, we have

$$(1.1) \quad \sum_{U < \gamma \leq V} \Phi(\gamma) = \int_U^V \Phi(t) dN(t) = - \int_U^V N(t) \Phi'(t) dt + N(V)\Phi(V) - N(U)\Phi(U).$$

About $N(T)$, Riemann [5] guessed that

$$(1.2) \quad N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + O(\log T).$$

This conjecture of Riemann proved by H. von Mangoldt more than 30 years later [1, 2]. An immediate corollary of above approximate formula, which is known as Riemann-van Mangoldt formula is $\mathcal{A}(T) = O(\log^2 T)$, which follows by partial summation from Riemann-van Mangoldt formula [2]. In 1941, Rosser [6] introduced the following approximation of $N(T)$:

$$(1.3) \quad |N(T) - F(T)| \leq R(T) \quad (T \geq 2),$$

where $F(T) = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + \frac{7}{8}$, and $R(T) = \frac{137}{1000} \log T + \frac{433}{1000} \log \log T + \frac{397}{250}$. This approximation allows us to make some explicit approximation of $\mathcal{A}(T)$.

2. APPROXIMATION OF $\mathcal{A}(T)$

2.1. Approximate Estimation of $\mathcal{A}(T)$. As we set above, γ_1 is the imaginary part of first nontrivial zero of the Riemann zeta function in the upper half plane and computations [4] give us $\gamma_1 = 14.13472514 \dots$. On using (1.1) with $\Phi(\gamma) = \frac{1}{\gamma}$, $0 < U < \gamma_1$ and $V = T$, we obtain

$$(2.1) \quad \mathcal{A}(T) = \int_U^T \frac{dN(t)}{t} = \int_U^T \frac{N(t)}{t^2} dt + \frac{N(T)}{T}.$$

Substituting $N(T)$ from (1.2), we obtain

$$\mathcal{A}(T) = \frac{1}{2\pi} \int_U^T \frac{\log\left(\frac{t}{2\pi}\right)}{t} dt - \frac{1}{2\pi} \int_U^T \frac{dt}{t} + \frac{1}{2\pi} \log \frac{T}{2\pi} - \frac{1}{2\pi} + O\left(\int_U^T \frac{\log(t)}{t^2} dt\right) + O\left(\frac{\log T}{T}\right).$$

Computing integrals and error terms, and then letting $U \rightarrow \gamma_1^-$, we get the following approximation

$$\mathcal{A}(T) = \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + O(1).$$

2.2. Explicit Estimation of $\mathcal{A}(T)$. Considering (1.3) and using (2.1) with $2 \leq U < \gamma_1$, for every $T \geq 2$ implies

$$\int_U^T \frac{F(t)}{t^2} dt - \int_U^T \frac{R(t)}{t^2} dt + \frac{F(T) - R(T)}{T} \leq \mathcal{A}(T) \leq \int_U^T \frac{F(t)}{t^2} dt + \int_U^T \frac{R(t)}{t^2} dt + \frac{F(T) + R(T)}{T}.$$

A simple calculation, yields $\frac{F(t)}{t^2} = \frac{d}{dt} \left\{ \frac{1}{4\pi} \log^2 t - \frac{1+\log(2\pi)}{2\pi} \log t + \frac{\log^2(2\pi) - 2\log(2\pi)}{4\pi} - \frac{7}{8t} \right\}$, and setting $\mathfrak{E}(t) = \int_1^\infty \frac{ds}{st^s}$, we also have $\frac{R(t)}{t^2} = \frac{d}{dt} \left\{ -\frac{433}{1000} \frac{\log \log t}{t} - \frac{137}{1000} \frac{\log t}{t} - \frac{69}{40t} - \frac{433}{1000} \mathfrak{E}(t) \right\}$. The integral of $\mathfrak{E}(t)$ converges for $t > 1$; in fact $\mathfrak{E}(t) \sim \frac{1}{t \log t}$ when $t \rightarrow \infty$. Using the fact $\frac{d}{dt} \mathfrak{E}(t) = -\frac{1}{t^2 \log t}$, we get $\frac{1}{t \log t} - \frac{1}{t \log^2 t} < \mathfrak{E}(t) < \frac{1}{t \log t} - \frac{31}{95t \log^2 t}$ for $t \geq 2$. Therefore, after letting $U \rightarrow \gamma_1^-$, we get the following explicit upper bound

$$\mathcal{A}(T) < \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \mathbf{c}_{\text{au}} - \frac{137 \log^2 T + 433 \log T - 433}{1000T \log^2 T} \quad (T \geq 2),$$

where $\mathbf{c}_{\text{au}} = 0.43596427 \dots < \frac{109}{250}$ and for $T \geq 2.222$ we have $-\frac{137 \log^2 T + 433 \log T - 433}{1000T \log^2 T} < 0$. Thus, we obtain $\mathcal{A}(T) < \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \frac{109}{250}$ for $T \geq 2.222$. Similarly, we get

$$\begin{aligned} \mathcal{A}(T) &> \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \mathbf{c}_{\text{al}} \\ &+ \frac{274 \log^3 T + 866(\log \log T) \log^2 T + 3313 \log^2 T + 433 \log T - 433}{1000T \log^2 T} \quad (T \geq 2), \end{aligned}$$

where $\mathbf{c}_{\text{al}} = 0.06058187 \dots > \frac{3}{50}$ and for $T \geq 2$ the last term in the above inequality is positive. So, we obtain $\mathcal{A}(T) > \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T + \frac{3}{50}$ for $T \geq 2$. Therefore we have proved the following result:

Proposition 2.1. *Letting $\mathcal{A}(T) = \sum_{0 < \gamma \leq T} \frac{1}{\gamma}$ with γ is imaginary part of zeta zeros, we have*

$$(2.2) \quad \frac{15}{250} < \mathcal{A}(T) - \left\{ \frac{1}{4\pi} \log^2 T - \frac{\log(2\pi)}{2\pi} \log T \right\} < \frac{109}{250},$$

where the left hand side holds for $T \geq 2$ and the right hand side holds for $T \geq 2.222$.

REFERENCES

- [1] H. Davenport, *Multiplicative Number Theory (Second Edition)*, Springer-Verlag, 1980.
- [2] Aleksandar Ivic, *The Riemann Zeta Function*, John Wiley & sons, 1985.
- [3] N.N. Lebedev, *Special Functions and their Applications*, Translated and edited by Richard A. Silverman, Dover Publications, New York, 1972.
- [4] Andrew Odlyzko, Tables of zeros of the Riemann zeta function:
http://www.dtc.umn.edu/~odlyzko/zeta_tables/
- [5] Bernhard Riemann, Ueber die Anzahl der Primzahlen unter einer gegebenen Grosse (On the Number of Prime Numbers less than a Given Quantity), *Monatsberichte der Berliner Akademie*, November 1859.

- [6] J. Barkley Rosser, Explicit bounds for some functions of prime numbers, *Amer. J. Math.*, Vol. 63, (1941) pp. 211-232.

MEHDI HASSANI,

DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES, P.O. BOX 45195-1159, ZANJAN, IRAN

E-mail address: mmhassany@yahoo.com, mmhassany@srttu.edu