

# OSTROWSKI'S INEQUALITIES FOR FUNCTIONS WHOSE DERIVATIVES ARE $s$ -CONVEX IN THE SECOND SENSE

M. ALOMARI<sup>A,\*</sup>, M. DARUS<sup>A</sup>, S.S. DRAGOMIR<sup>B</sup>, AND P. CERONE<sup>B</sup>

ABSTRACT. New inequalities of Ostrowski type for functions whose derivatives in absolute value are  $s$ -convex in the second sense are obtained.

## 1. INTRODUCTION

Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ , the interior of the interval  $I$ , such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'(x)| \leq M$ , then the following inequality,

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{b-a} \cdot \left[ \frac{(x-a)^2 + (b-x)^2}{2} \right]$$

holds. This result is known in the literature as the *Ostrowski inequality*. For recent results and generalizations concerning Ostrowski's inequality, see [1] – [5] and the references therein.

In [8], Hudzik and Maligranda considered among others the class of functions which are  *$s$ -convex in the second sense*. This class is defined in the following way: a function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ , where  $\mathbb{R}^+ = [0, \infty)$ , is said to be  $s$ -convex in the second sense if

$$f(\alpha x + \beta y) \leq \alpha^s f(x) + \beta^s f(y)$$

for all  $x, y \in [0, \infty)$ ,  $\alpha, \beta \geq 0$  with  $\alpha + \beta = 1$  and for some fixed  $s \in (0, 1]$ . This class of  $s$ -convex functions is usually denoted by  $K_s^2$ .

It can be easily seen that for  $s = 1$ ,  $s$ -convexity reduces to the ordinary convexity of functions defined on  $[0, \infty)$ .

In [7], Dragomir and Fitzpatrick proved a variant of Hadamard's inequality which holds for  $s$ -convex functions in the second sense:

**Theorem 1.** *Suppose that  $f : [0, \infty) \rightarrow [0, \infty)$  is an  $s$ -convex function in the second sense, where  $s \in (0, 1)$  and let  $a, b \in [0, \infty)$ ,  $a < b$ . If  $f \in L^1[a, b]$ , then the following inequalities hold:*

$$(1.2) \quad 2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1}.$$

*The constant  $k = \frac{1}{s+1}$  is the best possible in the second inequality in (1.2). The above inequalities are sharp.*

---

*Date:* July 08, 2009.

*2000 Mathematics Subject Classification.* 26D15.

*Key words and phrases.* Ostrowski's inequality, Convex function,  $s$ -convex function.

\*corresponding author.

For recent results and generalizations concerning  $s$ -convex functions see [7] – [9].

The aim of this paper is to establish some Ostrowski type inequalities for the class of functions whose derivatives in absolute value are  $s$ -convex functions in the second sense.

## 2. OSTROWSKI'S TYPE INEQUALITIES

In order to prove our main theorems, we need the following lemma that has been obtained in [1]:

**Lemma 1.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  where  $a, b \in I$  with  $a < b$ . If  $f' \in L[a, b]$ , then the following equality holds:*

$$f(x) - \frac{1}{b-a} \int_a^b f(u) du = \frac{(x-a)^2}{b-a} \int_0^1 t f'(tx + (1-t)a) dt - \frac{(b-x)^2}{b-a} \int_0^1 t f'(tx + (1-t)b) dt$$

for each  $x \in [a, b]$ .

A simple proof of the equality can be done by performing an integration by parts in the integrals from the right side and changing the variable. The details are left to the interested reader.

The following result may be stated:

**Theorem 2.** *Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|$  is  $s$ -convex in the second sense on  $[a, b]$  for some fixed  $s \in (0, 1]$  and  $|f'(x)| \leq M$ ,  $x \in [a, b]$ , then the following inequality holds:*

$$(2.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{b-a} \cdot \left[ \frac{(x-a)^2 + (b-x)^2}{s+1} \right],$$

for each  $x \in [a, b]$ .

*Proof.* By Lemma 1 and since  $|f'|$  is  $s$ -convex, then we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{(x-a)^2}{b-a} \int_0^1 t |f'(tx + (1-t)a)| dt \\ & \quad + \frac{(b-x)^2}{b-a} \int_0^1 t |f'(tx + (1-t)b)| dt \\ & \leq \frac{(x-a)^2}{b-a} \int_0^1 t [t^s |f'(x)| + (1-t)^s |f'(a)|] dt \\ & \quad + \frac{(b-x)^2}{b-a} \int_0^1 t [t^s |f'(x)| + (1-t)^s |f'(b)|] dt \end{aligned}$$

$$\begin{aligned}
&\leq \frac{M(x-a)^2}{b-a} \left( \frac{1}{s+2} + \frac{1}{(s+1)(s+2)} \right) \\
&\quad + \frac{M(b-x)^2}{b-a} \left( \frac{1}{s+2} + \frac{1}{(s+1)(s+2)} \right) \\
&= \frac{M}{b-a} \cdot \left[ \frac{(x-a)^2 + (b-x)^2}{s+1} \right],
\end{aligned}$$

where we have used the fact that

$$\int_0^1 t^{s+1} dt = \frac{1}{s+2} \quad \text{and} \quad \int_0^1 t(1-t)^s dt = \frac{1}{(s+1)(s+2)}.$$

This completes the proof. ■

The corresponding version for powers of the absolute value of the first derivative is incorporated in the following result:

**Theorem 3.** *Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is  $s$ -convex in the second sense on  $[a, b]$  for some fixed  $s \in (0, 1]$ ,  $p, q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$  and  $|f'(x)| \leq M$ ,  $x \in [a, b]$ , then the following inequality holds:*

$$(2.2) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{(1+p)^{\frac{1}{p}}} \left( \frac{2}{s+1} \right)^{\frac{1}{q}} \left[ \frac{(x-a)^2 + (b-x)^2}{b-a} \right],$$

for each  $x \in [a, b]$ .

*Proof.* Suppose that  $p > 1$ . From Lemma 1 and using the Hölder inequality, we have

$$\begin{aligned}
&\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \\
&\leq \frac{(x-a)^2}{b-a} \int_0^1 t |f'(tx + (1-t)a)| dt \\
&\quad + \frac{(b-x)^2}{b-a} \int_0^1 t |f'(tx + (1-t)b)| dt \\
&\leq \frac{(x-a)^2}{b-a} \left( \int_0^1 t^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\
&\quad + \frac{(b-x)^2}{b-a} \left( \int_0^1 t^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}}.
\end{aligned}$$

Since  $|f'|^q$  is  $s$ -convex in the second sense and  $|f'(x)| \leq M$ , then we have

$$\begin{aligned}
\int_0^1 |f'(tx + (1-t)a)|^q dt &\leq \int_0^1 [t^s |f'(x)|^q + (1-t)^s |f'(a)|^q] dt \\
&= \frac{|f'(x)|^q + |f'(a)|^q}{s+1} \leq \frac{2M^q}{s+1}
\end{aligned}$$

and

$$\begin{aligned} \int_0^1 |f'(tx + (1-t)b)|^q dt &\leq \int_0^1 [t^s |f'(x)|^q + (1-t)^s |f'(b)|^q] dt \\ &= \frac{|f'(x)|^q + |f'(b)|^q}{s+1} \leq \frac{2M^q}{s+1}. \end{aligned}$$

Therefore, we have

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{(1+p)^{\frac{1}{p}}} \left( \frac{2}{s+1} \right)^{\frac{1}{q}} \left[ \frac{(x-a)^2 + (b-x)^2}{b-a} \right],$$

where  $1/p + 1/q = 1$ , which is required. ■

**Corollary 1.** *Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^{p/(p-1)}$  is convex on  $[a, b]$ ,  $p > 1$ , and  $|f'(x)| \leq M$ ,  $x \in [a, b]$ , then the following inequality holds:*

$$(2.3) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{b-a} \left[ \frac{(x-a)^2 + (b-x)^2}{(1+p)^{\frac{1}{p}}} \right]$$

for each  $x \in [a, b]$ .

*Proof.* Setting  $s = 1$  in (2.2), we get the required result. ■

A different approach leads to the following result:

**Theorem 4.** *Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is  $s$ -convex in the second sense on  $[a, b]$  for some fixed  $s \in (0, 1]$  and  $q \geq 1$ , and  $|f'(x)| \leq M$ ,  $x \in [a, b]$ , then the following inequality holds:*

$$(2.4) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq M \left( \frac{2}{s+1} \right)^{\frac{1}{q}} \left[ \frac{(x-a)^2 + (b-x)^2}{2(b-a)} \right]$$

for each  $x \in [a, b]$ .

*Proof.* Suppose that  $q \geq 1$ . From Lemma 1 and using the well known power mean inequality, we have

$$\begin{aligned} &\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \\ &\leq \frac{(x-a)^2}{b-a} \int_0^1 t |f'(tx + (1-t)a)| dt \\ &\quad + \frac{(b-x)^2}{b-a} \int_0^1 t |f'(tx + (1-t)b)| dt \end{aligned}$$

$$\begin{aligned} &\leq \frac{(x-a)^2}{b-a} \left( \int_0^1 t dt \right)^{1-\frac{1}{q}} \left( \int_0^1 t |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\ &\quad + \frac{(b-x)^2}{b-a} \left( \int_0^1 t dt \right)^{1-\frac{1}{q}} \left( \int_0^1 t |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}}. \end{aligned}$$

Since  $|f'|^q$  is  $s$ -convex, we have

$$\begin{aligned} \int_0^1 t |f'(tx + (1-t)a)|^q dt &\leq \int_0^1 [t^{s+1} |f'(x)|^q + t(1-t)^s |f'(a)|^q] dt \\ &= \frac{|f'(x)|^q + (s+1) |f'(a)|^q}{(s+1)(s+2)} \leq \frac{M^q}{s+1} \end{aligned}$$

and

$$\begin{aligned} \int_0^1 t |f'(tx + (1-t)b)|^q dt &\leq \int_0^1 [t^{s+1} |f'(x)|^q + t(1-t)^s |f'(b)|^q] dt \\ &= \frac{|f'(x)|^q + (s+1) |f'(b)|^q}{(s+1)(s+2)} \leq \frac{M^q}{s+1}. \end{aligned}$$

Therefore, we have

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq M \left( \frac{1}{2} \right)^{1-\frac{1}{q}} \left( \frac{1}{s+1} \right)^{\frac{1}{q}} \left[ \frac{(x-a)^2 + (b-x)^2}{b-a} \right]$$

which is required. ■

**Corollary 2.** Let  $f : I \subset [0, \infty) \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$  such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is convex on  $[a, b]$ ,  $q \geq 1$ , then inequality (1.1) holds.

*Proof.* Setting  $s = 1$  in (1.2), we get the required result. ■

**Remark 1.** Since  $(1+p)^{\frac{1}{p}} < 2$  for any  $p > 1$ , then we observe that the inequality (2.4) is better than (2.2), meaning that the approach via the power mean inequality is better than the one through Hölder's inequality.

Midpoint type inequalities for functions whose derivatives in absolute value are  $s$ -convex in the second sense may be obtained from the previous results as follows:

**Corollary 3.** If in (2.4) we choose  $x = \frac{a+b}{2}$ , then we have

$$(2.5) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M(b-a)}{4} \left( \frac{2}{s+1} \right)^{\frac{1}{q}}, \quad q \geq 1,$$

where  $s \in (0, 1]$  and  $|f'|^q$  is  $s$ -convex in the second sense on  $[a, b]$ ,  $q \geq 1$ .

### 3. APPLICATIONS TO SPECIAL MEANS

We consider the means for arbitrary positive numbers  $\alpha, \beta$  ( $\alpha \neq \beta$ ) as follows:

(1) *Arithmetic mean:*

$$A(\alpha, \beta) = \frac{\alpha + \beta}{2},$$

(2) *Generalized log-mean*:

$$L_p(\alpha, \beta) = \left[ \frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{\frac{1}{p}}, \quad p \in \mathbb{R} \setminus \{-1, 0\}.$$

Now, using the results of Section 2, we give some applications to special means of real numbers.

In [8], the following example is given: Let  $s \in (0, 1)$  and  $a, b, c \in \mathbb{R}$ . We define a function  $f : [0, \infty) \rightarrow \mathbb{R}$ , as

$$f(t) = \begin{cases} a, & t = 0; \\ bt^s + c, & t > 0. \end{cases}$$

If  $b \geq 0$  and  $0 \leq c \leq a$ , then  $f \in K_s^2$ . Hence, for  $a = c = 0$ ,  $b = 1$ , we have  $f : [0, 1] \rightarrow [0, 1]$ ,  $f(t) = t^s$ ,  $f \in K_s^2$ .

**Proposition 1.** *Let  $0 < a < b$  and  $0 < s < 1$ . Then we have*

$$|A^s(a, b) - L_s^s(a, b)| \leq \frac{s(b-a)}{4} \left( \frac{2}{s+1} \right)^{\frac{1}{q}}, \quad q \geq 1.$$

*Proof.* The inequality follows from (2.5) applied to the  $s$ -convex function in the second sense  $f : [0, 1] \rightarrow [0, 1]$ ,  $f(x) = x^s$ . The details are omitted. ■

**Acknowledgement.** The first author acknowledges the financial support of Universiti Kebangsaan Malaysia, Faculty of Science and Technology, (UKM-GUP-TMK-07-02-107).

## REFERENCES

- [1] P. Cerone and S.S. Dragomir, Ostrowski type inequalities for functions whose derivatives satisfy certain convexity assumptions, *Demonstratio Math.*, 37 (2004), no. 2, 299-308
- [2] S.S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for  $s$ -convex functions in the second sense, *Demonstratio Math.*, 32 (4) (1999), 687-696.
- [3] N.S. Barnett, P. Cerone, S.S. Dragomir, M.R. Pinheiro and A. Sofo, Ostrowski type inequalities for functions whose modulus of derivatives are convex and applications, *RGMA Res. Rep. Coll.*, 5(2) (2002), Article 1. [ONLINE: <http://www.staff.vu.edu.au/RGMIA/v5n2.asp>]
- [4] S.S. Dragomir and A. Sofo, Ostrowski type inequalities for functions whose derivatives are convex, Proceedings of the 4th International Conference on Modelling and Simulation, November 11-13, 2002. Victoria University, Melbourne, Australia. *RGMA Res. Rep. Coll.*, 5(2002), Supplement, Article 30. [ONLINE: [http://www.staff.vu.edu.au/RGMIA/v5\(E\).asp](http://www.staff.vu.edu.au/RGMIA/v5(E).asp)]
- [5] S.S. Dragomir and Th. M. Rassias, (Eds) *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publishers, Dordrecht/Boston/London, 2002.
- [6] S.S. Dragomir and C.E.M. Pearce, Quasi-convex functions and Hadamard's inequality, *Bull. Australian Math. Soc.*, 57(1998), 377-385.
- [7] S.S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for  $s$ -convex functions in the second sense, *Demonstratio Math.*, 32 (4) (1999), 687-696.
- [8] H. Hudzik and L. Maligranda, Some remarks on  $s$ -convex functions, *Aequationes Math.*, 48 (1994), 100-111.
- [9] U.S. Kirmaci et al., Hadamard-type inequalities for  $s$ -convex functions, *Appl. Math. Comp.*, 193 (2007), 26-35.

<sup>a,\*</sup>SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITI KEBANGSAAN MALAYSIA, UKM, BANGI,  
43600, SELANGOR, MALAYSIA

*E-mail address:* [alomari@math.com](mailto:alomari@math.com)

*E-mail address:* [maslina@ukm.my](mailto:maslina@ukm.my)

<sup>b</sup>MATHEMATICS., SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428,  
MELBOURNE CITY, MC 8001, AUSTRALIA.

*E-mail address:* [sever.dragomir@vu.edu.au](mailto:sever.dragomir@vu.edu.au)

*URL:* <http://www.staff.vu.edu.au/rgmia/dragomir/>

*E-mail address:* [Pietro.Cerone@vu.edu.au](mailto:Pietro.Cerone@vu.edu.au)

*URL:* <http://www.staff.vu.edu.au/RGMIA/Cerone/>