

NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR FUNCTIONS WHOSE SECOND DERIVATIVES ABSOLUTE VALUES ARE QUASI-CONVEX

M. ALOMARI^{★,*}, M. DARUS[★], AND S.S. DRAGOMIR[◆]

ABSTRACT. In this note we obtain some inequalities of Hermite-Hadamard type for functions whose second derivatives have quasi-convex absolute values. Applications for special means are also provided.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping defined on the interval I of real numbers and $a, b \in I$, with $a < b$. The following two inequalities:

$$f\left(\frac{a+b}{2}\right) \leq \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

hold. This double inequality is known in the literature as the Hermite-Hadamard inequality for convex functions.

In recent years many authors have established several inequalities connected to this fact. For recent results, refinements, counterparts, generalizations and new Hermite-Hadamard-type inequalities see [1] – [5] and [7] – [11].

We recall that the notion of quasi-convex functions generalizes the notion of convex functions. More precisely, a function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *quasi-convex* on $[a, b]$ if

$$f(\lambda x + (1 - \lambda)y) \leq \max\{f(x), f(y)\}, \quad \forall x, y \in [a, b].$$

Clearly, any convex function is a quasi-convex function. Furthermore, there exist quasi-convex functions which are not convex, (see for instance [6]).

Recently, D.A. Ion [6] obtained two inequalities for the right hand side of the Hermite-Hadamard inequality for functions whose derivatives in absolute values are quasi-convex functions, as follows:

Theorem 1. *Let $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° , $a, b \in I^\circ$ with $a < b$. If $|f'|$ is quasi-convex on $[a, b]$, then the following inequality holds:*

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{4} \max\{|f'(a)|, |f'(b)|\}.$$

Key words and phrases. Convex function, s -Convex function, Hermite-Hadamard's inequality.

*The first author acknowledges the financial support of the Universiti Kebangsaan Malaysia, Faculty of Science and Technology, (UKM-GUP-TMK-07-02-107).

Theorem 2. Let $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° , $a, b \in I^\circ$ with $a < b$. If $|f'|^{p/(p-1)}$ is quasi-convex on $[a, b]$, then the following inequality holds:

$$\begin{aligned} \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)}{2(p+1)^{1/p}} \left(\max \left\{ |f'(a)|^{p/(p-1)}, |f'(b)|^{p/(p-1)} \right\} \right)^{(p-1)/p}. \end{aligned}$$

The main aim of this paper is to establish new refined inequalities of the right-hand side of the Hermite-Hadamard result for the class of functions whose second derivatives at certain powers are quasi-convex functions.

2. HERMITE-HADAMARD TYPE INEQUALITIES

In order to prove our main theorems, we need the following lemma [5], [10].

Lemma 1. Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I° , $a, b \in I$ with $a < b$ and f'' be integrable on $[a, b]$, then the following equality holds:

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx = \frac{(b-a)^2}{2} \int_0^1 t(1-t) f''(ta + (1-t)b) dt.$$

A simple proof of this equality can be also done by twice integrating by parts in the right hand side. The details are left to the interested reader.

The next theorem gives a new result of the upper Hermite-Hadamard inequality for quasi-convex functions.

Theorem 3. Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I° , $a, b \in I$ with $a < b$ and f'' be integrable on $[a, b]$. If $|f''|$ is quasi-convex on $[a, b]$, then the following inequality holds:

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{12} \max \{|f''(a)|, |f''(b)|\}.$$

Proof. From Lemma 1, we have

$$\begin{aligned} \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| &\leq \frac{(b-a)^2}{2} \int_0^1 t(1-t) |f''(ta + (1-t)b)| dt \\ &\leq \frac{(b-a)^2}{2} \int_0^1 t(1-t) \max \{|f''(a)|, |f''(b)|\} dt \\ &\leq \frac{(b-a)^2}{2} \max \{|f''(a)|, |f''(b)|\} \int_0^1 t(1-t) dt \\ &= \frac{(b-a)^2}{12} \max \{|f''(a)|, |f''(b)|\}, \end{aligned}$$

which completes the proof. ■

A similar result is embodied in the following theorem.

Theorem 4. Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I° , $a, b \in I$ with $a < b$ and f'' is integrable on $[a, b]$. If $|f''|^{p/(p-1)}$ is quasi-convex on $[a, b]$, for

$p > 1$, then the following inequality holds:

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{(b-a)^2}{8} \left(\frac{\sqrt{\pi}}{2} \right)^{\frac{1}{p}} \left(\frac{\Gamma(1+p)}{\Gamma(\frac{3}{2}+p)} \right)^{\frac{1}{p}} (\max\{|f''(a)|^q, |f''(b)|^q\})^{\frac{1}{q}} \end{aligned}$$

where $q = p/(p-1)$.

Proof. From Lemma 1 and using the well known Hölder integral inequality, we have successively

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{(b-a)^2}{2} \int_0^1 t(1-t) |f''(ta + (1-t)b)| dt \\ & \leq \frac{(b-a)^2}{2} \left(\int_0^1 (t-t^2)^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |f''(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{(b-a)^2}{2} \cdot \left(\frac{2^{-1-2p} \sqrt{\pi} \Gamma(1+p)}{\Gamma(\frac{3}{2}+p)} \right)^{\frac{1}{p}} \cdot (\max\{|f''(a)|^q, |f''(b)|^q\})^{\frac{1}{q}} \\ & = \frac{(b-a)^2}{8} \left(\frac{\sqrt{\pi}}{2} \right)^{\frac{1}{p}} \left(\frac{\Gamma(1+p)}{\Gamma(\frac{3}{2}+p)} \right)^{\frac{1}{p}} (\max\{|f''(a)|^q, |f''(b)|^q\})^{\frac{1}{q}}, \end{aligned}$$

where $1/p + 1/q = 1$, and we use the fact that

$$\int_0^1 (t-t^2)^p dt = \frac{2^{-1-2p} \sqrt{\pi} \Gamma(1+p)}{\Gamma(\frac{3}{2}+p)},$$

which completes the proof. ■

A more general inequality is given using Lemma 1, as follows:

Theorem 5. Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable mapping on I° , $a, b \in I$ with $a < b$ and f'' is integrable on $[a, b]$. If $|f''|^q$ is an quasi-convex on $[a, b]$, $q \geq 1$, then the following inequality holds:

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{12} (\max\{|f''(a)|^q, |f''(b)|^q\})^{1/q}$$

Proof. From Lemma 1 and using well known power mean inequality, we have

$$\begin{aligned}
& \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
& \leq \frac{(b-a)^2}{2} \int_0^1 t(1-t) |f''(ta + (1-t)b)| dt \\
& \leq \frac{(b-a)^2}{2} \left(\int_0^1 (t-t^2) dt \right)^{1-\frac{1}{q}} \left(\int_0^1 (t-t^2) |f''(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{(b-a)^2}{2} \cdot \left(\frac{1}{6} \right)^{1-\frac{1}{q}} \cdot \left(\frac{1}{6} \max \{ |f''(a)|^q, |f''(b)|^q \} \right)^{\frac{1}{q}} \\
& = \frac{(b-a)^2}{12} \left(\max \{ |f''(a)|^q, |f''(b)|^q \} \right)^{\frac{1}{q}},
\end{aligned}$$

which completes the proof. ■

3. APPLICATIONS TO SPECIAL MEANS

We now consider the means for arbitrary real numbers α, β ($\alpha \neq \beta$). We take

(1) *Arithmetic mean:*

$$A(\alpha, \beta) = \frac{\alpha + \beta}{2}, \quad \alpha, \beta \in \mathbb{R}.$$

(2) *Logarithmic mean:*

$$L(\alpha, \beta) = \frac{\alpha - \beta}{\ln |\alpha| - \ln |\beta|}, \quad |\alpha| \neq |\beta|, \alpha, \beta \neq 0, \alpha, \beta \in \mathbb{R}.$$

(3) *Generalized log-mean:*

$$L_n(\alpha, \beta) = \left[\frac{\beta^{n+1} - \alpha^{n+1}}{(n+1)(\beta - \alpha)} \right]^{\frac{1}{n}}, \quad n \in \mathbb{Z} \setminus \{-1, 0\}, \alpha, \beta \in \mathbb{R}, \alpha \neq \beta.$$

Now, using the results of Section 2, we give some applications for special means of real numbers.

Proposition 1. *Let $a, b \in \mathbb{R}$, $a < b$ and $n \in \mathbb{N}$, $n \geq 2$. Then, we have*

$$|L_n^n(a, b) - A(a^n, b^n)| \leq \frac{n(n-1)}{12} (b-a)^2 \max \{ |a|^{n-2}, |b|^{n-2} \}.$$

Proof. The assertion follows from Theorem 3 applied to the quasi-convex mapping $f(x) = x^n$, $x \in \mathbb{R}$. ■

Proposition 2. *Let $a, b \in \mathbb{R}$, $a < b$ and $0 \notin [a, b]$. Then, for all $p > 1$, we have*

$$\begin{aligned}
& |L^{-1}(a, b) - A(a^{-1}, b^{-1})| \\
& \leq \frac{(b-a)^2}{4} \left(\frac{\sqrt{\pi}}{2} \right)^{\frac{1}{p}} \left(\frac{\Gamma(1+p)}{\Gamma(\frac{3}{2}+p)} \right)^{\frac{1}{p}} \left(\max \{ |a|^{-3q}, |b|^{-3q} \} \right)^{\frac{1}{q}}.
\end{aligned}$$

Proof. The assertion follows from Theorem 4 applied to the quasi-convex mapping $f(x) = 1/x$, $x \in [a, b]$. ■

Proposition 3. *Let $a, b \in \mathbb{R}$, $a < b$ and $n \in \mathbb{N}$, $n \geq 2$. Then, for all $q \geq 1$, we have*

$$|L_n^n(a, b) - A^n(a, b)| \leq \frac{n(n-1)}{12} (b-a)^2 \left(\max \left\{ |a|^{(n-2)q}, |b|^{(n-2)q} \right\} \right)^{\frac{1}{q}}.$$

Proof. The assertion follows from Theorem 5 applied to the quasi-convex mapping $f(x) = x^n$, $x \in \mathbb{R}$. ■

REFERENCES

- [1] S.S. Dragomir, Two mappings in connection to Hadamard's inequalities, *J. Math. Anal. Appl.*, **167** (1992), 49–56.
- [2] S.S. Dragomir and R.P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Math. Lett.*, **11** (1998), 91–95.
- [3] S.S. Dragomir, Y.J. Cho and S.S. Kim, Inequalities of Hadamard's type for Lipschitzian mappings and their applications, *J. Math. Anal. Appl.*, **245** (2000), 489–501.
- [4] S.S. Dragomir and S. Wang, A new inequality of Ostrowski's type in L_1 norm and applications to some special means and to some numerical quadrature rule, *Tamkang J. Math.*, **28** (1997), 239–244.
- [5] S.S. Dragomir, On some inequalities for differentiable convex functions and applications, (submitted).
- [6] D.A. Ion, Some estimates on the Hermite-Hadamard inequality through quasi-convex functions, *Annals of University of Craiova Math. Comp. Sci. Ser.*, **34** (2007), 82–87.
- [7] U.S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers to midpoint formula, *Appl. Math. Comp.*, **147** (2004), 137–146.
- [8] U.S. Kirmaci and M.E. Özdemir, On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comp.*, **153** (2004), 361–368.
- [9] M.E. Özdemir, A theorem on mappings with bounded derivatives with applications to quadrature rules and means, *Appl. Math. Comp.*, **138** (2003), 425–434.
- [10] S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000. Online: [http://www.staff.vu.edu.au/RGMIA/monographs/hermite_hadamard.html].
- [11] C.E.M. Pearce and J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formula, *Appl. Math. Lett.*, **13** (2000), 51–55.
- [12] G.S. Yang, D.Y. Hwang and K.L. Tseng, Some inequalities for differentiable convex and concave mappings, *Appl. Math. Lett.*, **47** (2004), 207–216.

★SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITI KEBANGSAAN MALAYSIA,, UKM, BANGI, 43600, SELANGOR, MALAYSIA

E-mail address: mwomath@gmail.com

E-mail address: maslina@ukm.my

◆RESEARCH GROUP IN MATHEMATICAL INEQUALITIES & APPLICATIONS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://www.staff.vu.edu.au/rgmia/dragomir/>