

# INEQUALITIES IN TERMS OF THE GÂTEAUX DERIVATIVES FOR CONVEX FUNCTIONS IN LINEAR SPACES WITH APPLICATIONS

S.S. DRAGOMIR

ABSTRACT. Some inequalities in terms of the Gâteaux derivatives related to Jensen's inequality for convex functions defined on linear spaces are given. Applications for norms, mean  $f$ -deviations and  $f$ -divergence measures are provided as well.

## 1. INTRODUCTION

The Jensen inequality for convex functions plays a crucial role in the Theory of Inequalities due to the fact that other inequalities such as that arithmetic mean-geometric mean inequality, Hölder and Minkowski inequalities, Ky Fan's inequality etc. can be obtained as particular cases of it.

Let  $C$  be a convex subset of the linear space  $X$  and  $f$  a convex function on  $C$ . If  $\mathbf{p} = (p_1, \dots, p_n)$  is a probability sequence and  $\mathbf{x} = (x_1, \dots, x_n) \in C^n$ , then

$$(1.1) \quad f\left(\sum_{i=1}^n p_i x_i\right) \leq \sum_{i=1}^n p_i f(x_i),$$

is well known in the literature as Jensen's inequality.

Recently the author obtained the following refinement of Jensen's inequality (see [9])

$$(1.2) \quad \begin{aligned} f\left(\sum_{j=1}^n p_j x_j\right) &\leq \min_{k \in \{1, \dots, n\}} \left[ (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + p_k f(x_k) \right] \\ &\leq \frac{1}{n} \left[ \sum_{k=1}^n (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + \sum_{k=1}^n p_k f(x_k) \right] \\ &\leq \max_{k \in \{1, \dots, n\}} \left[ (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + p_k f(x_k) \right] \\ &\leq \sum_{j=1}^n p_j f(x_j), \end{aligned}$$

where  $f$ ,  $x_k$  and  $p_k$  are as above.

---

*Date:* February 07, 2009.

*1991 Mathematics Subject Classification.* 26D15; 94.

*Key words and phrases.* Convex functions, Gâteaux derivatives, Jensen's inequality, Norms, Mean  $f$ -deviations,  $f$ -Divergence measures.

The above result provides a different approach to the one that J. Pečarić and the author obtained in 1989, namely (see [14]):

$$\begin{aligned}
 (1.3) \quad f\left(\sum_{i=1}^n p_i x_i\right) &\leq \sum_{i_1, \dots, i_{k+1}=1}^n p_{i_1} \cdots p_{i_{k+1}} f\left(\frac{x_{i_1} + \cdots + x_{i_{k+1}}}{k+1}\right) \\
 &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \cdots p_{i_k} f\left(\frac{x_{i_1} + \cdots + x_{i_k}}{k}\right) \\
 &\leq \cdots \leq \sum_{i=1}^n p_i f(x_i),
 \end{aligned}$$

for  $k \geq 1$  and  $\mathbf{p}, \mathbf{x}$  as above.

If  $q_1, \dots, q_k \geq 0$  with  $\sum_{j=1}^k q_j = 1$ , then the following refinement obtained in 1994 by the author [6] also holds:

$$\begin{aligned}
 (1.4) \quad f\left(\sum_{i=1}^n p_i x_i\right) &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \cdots p_{i_k} f\left(\frac{x_{i_1} + \cdots + x_{i_k}}{k}\right) \\
 &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \cdots p_{i_k} f(q_1 x_{i_1} + \cdots + q_k x_{i_k}) \\
 &\leq \sum_{i=1}^n p_i f(x_i),
 \end{aligned}$$

where  $1 \leq k \leq n$  and  $\mathbf{p}, \mathbf{x}$  are as above.

For other refinements and applications related to Ky Fan's inequality, the arithmetic mean-geometric mean inequality, the generalised triangle inequality, the  $f$ -divergence measures etc. see [3]-[9].

In this paper, motivated by the above results, some new inequalities in terms of the Gâteaux derivatives related to Jensen's inequality for convex functions defined on linear spaces are given. Applications for norms, mean  $f$ -deviations and  $f$ -divergence measures are provided as well.

## 2. THE GÂTEAU DERIVATIVES OF CONVEX FUNCTIONS

Assume that  $f : X \rightarrow \mathbb{R}$  is a *convex function* on the real linear space  $X$ . Since for any vectors  $x, y \in X$  the function  $g_{x,y} : \mathbb{R} \rightarrow \mathbb{R}$ ,  $g_{x,y}(t) := f(x + ty)$  is convex it follows that the following limits exist

$$\nabla_{+(-)} f(x)(y) := \lim_{t \rightarrow 0+(-)} \frac{f(x + ty) - f(x)}{t}$$

and they are called the *right(left) Gâteaux derivatives* of the function  $f$  in the point  $x$  over the direction  $y$ .

It is obvious that for any  $t > 0 > s$  we have

$$\begin{aligned}
 (2.1) \quad \frac{f(x + ty) - f(x)}{t} &\geq \nabla_+ f(x)(y) = \inf_{t > 0} \left[ \frac{f(x + ty) - f(x)}{t} \right] \\
 &\geq \sup_{s < 0} \left[ \frac{f(x + sy) - f(x)}{s} \right] = \nabla_- f(x)(y) \geq \frac{f(x + sy) - f(x)}{s}
 \end{aligned}$$

for any  $x, y \in X$  and, in particular,

$$(2.2) \quad \nabla_- f(u)(u-v) \geq f(u) - f(v) \geq \nabla_+ f(v)(u-v)$$

for any  $u, v \in X$ . We call this *the gradient inequality* for the convex function  $f$ . It will be used frequently in the sequel in order to obtain various results related to Jensen's inequality.

The following properties are also of importance:

$$(2.3) \quad \nabla_+ f(x)(-y) = -\nabla_- f(x)(y),$$

and

$$(2.4) \quad \nabla_{+(-)} f(x)(\alpha y) = \alpha \nabla_{+(-)} f(x)(y)$$

for any  $x, y \in X$  and  $\alpha \geq 0$ .

The right Gâteaux derivative is *subadditive* while the left one is *superadditive*, i.e.,

$$(2.5) \quad \nabla_+ f(x)(y+z) \leq \nabla_+ f(x)(y) + \nabla_+ f(x)(z)$$

and

$$(2.6) \quad \nabla_- f(x)(y+z) \geq \nabla_- f(x)(y) + \nabla_- f(x)(z)$$

for any  $x, y, z \in X$ .

Some natural examples can be provided by the use of normed spaces.

Assume that  $(X, \|\cdot\|)$  is a real normed linear space. The function  $f : X \rightarrow \mathbb{R}$ ,  $f(x) := \frac{1}{2} \|x\|^2$  is a convex function which generates *the superior* and *the inferior semi-inner products*

$$\langle y, x \rangle_{s(i)} := \lim_{t \rightarrow 0+(-)} \frac{\|x + ty\|^2 - \|x\|^2}{t}.$$

For a comprehensive study of the properties of these mappings in the Geometry of Banach Spaces see the monograph [8].

For the convex function  $f_p : X \rightarrow \mathbb{R}$ ,  $f_p(x) := \|x\|^p$  with  $p > 1$ , we have

$$\nabla_{+(-)} f_p(x)(y) = \begin{cases} p \|x\|^{p-2} \langle y, x \rangle_{s(i)} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

for any  $y \in X$ .

If  $p = 1$ , then we have

$$\nabla_{+(-)} f_1(x)(y) = \begin{cases} \|x\|^{-1} \langle y, x \rangle_{s(i)} & \text{if } x \neq 0 \\ +(-) \|y\| & \text{if } x = 0 \end{cases}$$

for any  $y \in X$ .

This class of functions will be used to illustrate the inequalities obtained in the general case of convex functions defined on an entire linear space.

The following result holds:

**Theorem 1.** *Let  $f : X \rightarrow \mathbb{R}$  be a convex function. Then for any  $x, y \in X$  and  $t \in [0, 1]$  we have*

$$(2.7) \quad \begin{aligned} & t(1-t) [\nabla_- f(y)(y-x) - \nabla_+ f(x)(y-x)] \\ & \geq tf(x) + (1-t)f(y) - f(tx + (1-t)y) \\ & \geq t(1-t) [\nabla_+ f(tx + (1-t)y)(y-x) - \nabla_- f(tx + (1-t)y)(y-x)] \geq 0. \end{aligned}$$

*Proof.* Utilising the gradient inequality (2.2) we have

$$(2.8) \quad f(tx + (1-t)y) - f(x) \geq (1-t) \nabla_+ f(x)(y-x)$$

and

$$(2.9) \quad f(tx + (1-t)y) - f(y) \geq -t \nabla_- f(y)(y-x).$$

If we multiply (2.8) with  $t$  and (2.9) with  $1-t$  and add the resultant inequalities we obtain

$$\begin{aligned} & f(tx + (1-t)y) - tf(x) - (1-t)f(y) \\ & \geq (1-t)t \nabla_+ f(x)(y-x) - t(1-t) \nabla_- f(y)(y-x) \end{aligned}$$

which is clearly equivalent with the first part of (2.7).

By the gradient inequality we also have

$$(1-t) \nabla_- f(tx + (1-t)y)(y-x) \geq f(tx + (1-t)y) - f(x)$$

and

$$-t \nabla_+ f(tx + (1-t)y)(y-x) \geq f(tx + (1-t)y) - f(y)$$

which by the same procedure as above yields the second part of (2.7).  $\square$

The following particular case for norms may be stated:

**Corollary 1.** *If  $x$  and  $y$  are two vectors in the normed linear space  $(X, \|\cdot\|)$  such that  $0 \notin [x, y] := \{(1-s)x + sy, s \in [0, 1]\}$ , then for any  $p \geq 1$  we have the inequalities*

$$(2.10) \quad \begin{aligned} & pt(1-t) \left[ \|y\|^{p-2} \langle y-x, y \rangle_i - \|x\|^{p-2} \langle y-x, x \rangle_s \right] \\ & \geq t \|x\|^p + (1-t) \|y\|^p - \|tx + (1-t)y\|^p \\ & \geq pt(1-t) \|tx + (1-t)y\|^{p-2} [\langle y-x, tx + (1-t)y \rangle_s - \langle y-x, tx + (1-t)y \rangle_i] \geq 0 \end{aligned}$$

for any  $t \in [0, 1]$ . If  $p \geq 2$  the inequality holds for any  $x$  and  $y$ .

**Remark 1.** *We observe that for  $p = 1$  in (2.10) we derive the result*

$$(2.11) \quad \begin{aligned} & t(1-t) \left[ \left\langle y-x, \frac{y}{\|y\|} \right\rangle_i - \left\langle y-x, \frac{x}{\|x\|} \right\rangle_s \right] \\ & \geq t \|x\| + (1-t) \|y\| - \|tx + (1-t)y\| \\ & \geq t(1-t) \left[ \left\langle y-x, \frac{tx + (1-t)y}{\|tx + (1-t)y\|} \right\rangle_s - \left\langle y-x, \frac{tx + (1-t)y}{\|tx + (1-t)y\|} \right\rangle_i \right] \geq 0 \end{aligned}$$

while for  $p = 2$  we have

$$(2.12) \quad \begin{aligned} & 2t(1-t) [\langle y-x, y \rangle_i - \langle y-x, x \rangle_s] \\ & \geq t \|x\|^2 + (1-t) \|y\|^2 - \|tx + (1-t)y\|^2 \\ & \geq 2t(1-t) [\langle y-x, tx + (1-t)y \rangle_s - \langle y-x, tx + (1-t)y \rangle_i] \geq 0. \end{aligned}$$

We notice that the inequality (2.12) holds for any  $x, y \in X$  while in the inequality (2.11) we must assume that  $x, y$  and  $tx + (1-t)y$  are not zero.

**Remark 2.** If the normed space is smooth, i.e., the norm is Gâteaux differentiable in any nonzero point, then the superior and inferior semi-inner products coincide with the Lumer-Giles semi-inner product  $[\cdot, \cdot]$  that generates the norm and is linear in the first variable (see for instance [8]). In this situation the inequality (2.10) becomes

$$(2.13) \quad \begin{aligned} & pt(1-t) \left( \|y\|^{p-2} [y-x, y] - \|x\|^{p-2} [y-x, x] \right) \\ & \geq t \|x\|^p + (1-t) \|y\|^p - \|tx + (1-t)y\|^p \geq 0 \end{aligned}$$

and holds for any nonzero  $x$  and  $y$ .

Moreover, if  $(X, \langle \cdot, \cdot \rangle)$  is an inner product space, then (2.13) becomes

$$(2.14) \quad \begin{aligned} & pt(1-t) \left\langle y-x, \|y\|^{p-2} y - \|x\|^{p-2} x \right\rangle \\ & \geq t \|x\|^p + (1-t) \|y\|^p - \|tx + (1-t)y\|^p \geq 0. \end{aligned}$$

From (2.14) we deduce the particular inequalities of interest

$$(2.15) \quad t(1-t) \left\langle y-x, \frac{y}{\|y\|} - \frac{x}{\|x\|} \right\rangle \geq t \|x\| + (1-t) \|y\| - \|tx + (1-t)y\| \geq 0$$

and

$$(2.16) \quad 2t(1-t) \|y-x\|^2 \geq t \|x\|^2 + (1-t) \|y\|^2 - \|tx + (1-t)y\|^2 \geq 0.$$

Obviously, the inequality (2.16) can be proved directly on utilising the properties of the inner products.

**Problem 1.** It is an open question for the author whether the inequality (2.16) characterizes or not the class of inner product spaces within the class of normed spaces.

### 3. A REFINEMENT OF JENSEN'S INEQUALITY

For a convex function  $f : X \rightarrow \mathbb{R}$  defined on a linear space  $X$ , perhaps one of the most important result is the well known Jensen's inequality

$$(3.1) \quad f \left( \sum_{i=1}^n p_i x_i \right) \leq \sum_{i=1}^n p_i f(x_i),$$

which holds for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in X^n$  and any probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$ .

The following refinement of Jensen's inequality holds:

**Theorem 2.** *Let  $f : X \rightarrow \mathbb{R}$  be a convex function defined on a linear space  $X$ . Then for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in X^n$  and any probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$  we have the inequality*

$$(3.2) \quad \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right) \\ \geq \sum_{k=1}^n p_k \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)(x_k) - \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)\left(\sum_{i=1}^n p_i x_i\right) \geq 0.$$

In particular, for the uniform distribution, we have

$$(3.3) \quad \frac{1}{n} \sum_{i=1}^n f(x_i) - f\left(\frac{1}{n} \sum_{i=1}^n x_i\right) \\ \geq \frac{1}{n} \left[ \sum_{k=1}^n \nabla_+ f\left(\frac{1}{n} \sum_{i=1}^n x_i\right)(x_k) - \nabla_+ f\left(\frac{1}{n} \sum_{i=1}^n x_i\right)\left(\sum_{i=1}^n x_i\right) \right] \geq 0.$$

*Proof.* Utilising the gradient inequality (2.2) we have

$$(3.4) \quad f(x_k) - f\left(\sum_{i=1}^n p_i x_i\right) \geq \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)\left(x_k - \sum_{i=1}^n p_i x_i\right)$$

for any  $k \in \{1, \dots, n\}$ .

By the subadditivity of the functional  $\nabla_+ f(\cdot)(\cdot)$  in the second variable we also have

$$(3.5) \quad \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)\left(x_k - \sum_{i=1}^n p_i x_i\right) \\ \geq \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)(x_k) - \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)\left(\sum_{i=1}^n p_i x_i\right)$$

for any  $k \in \{1, \dots, n\}$ .

Utilising the inequalities (3.4) and (3.5) we get

$$(3.6) \quad f(x_k) - f\left(\sum_{i=1}^n p_i x_i\right) \\ \geq \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)(x_k) - \nabla_+ f\left(\sum_{i=1}^n p_i x_i\right)\left(\sum_{i=1}^n p_i x_i\right)$$

for any  $k \in \{1, \dots, n\}$ .

Now, if we multiply (3.6) with  $p_k \geq 0$  and sum over  $k$  from 1 to  $n$ , then we deduce the first inequality in (3.2). The second inequality is obvious by the subadditivity property of the functional  $\nabla_+ f(\cdot)(\cdot)$  in the second variable.  $\square$

The following particular case that provides a refinement for the generalised triangle inequality in normed linear spaces is of interest

**Corollary 2.** *Let  $(X, \|\cdot\|)$  be a normed linear space. Then for any  $p \geq 1$ , for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in X^n$  and any probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$  with  $\sum_{i=1}^n p_i x_i \neq 0$  we have the inequality*

$$(3.7) \quad \sum_{i=1}^n p_i \|x_i\|^p - \left\| \sum_{i=1}^n p_i x_i \right\|^p \\ \geq p \left\| \sum_{i=1}^n p_i x_i \right\|^{p-2} \left[ \sum_{k=1}^n p_k \left\langle x_k, \sum_{j=1}^n p_j x_j \right\rangle_s - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right] \geq 0.$$

If  $p \geq 2$  the inequality holds for any  $n$ -tuple of vectors and probability distribution.

In particular, we have the norm inequalities

$$(3.8) \quad \sum_{i=1}^n p_i \|x_i\| - \left\| \sum_{i=1}^n p_i x_i \right\| \\ \geq \left[ \sum_{k=1}^n p_k \left\langle x_k, \frac{\sum_{i=1}^n p_i x_i}{\left\| \sum_{i=1}^n p_i x_i \right\|} \right\rangle_s - \left\| \sum_{i=1}^n p_i x_i \right\| \right] \geq 0.$$

and

$$(3.9) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \\ \geq 2 \left[ \sum_{k=1}^n p_k \left\langle x_k, \sum_{i=1}^n p_i x_i \right\rangle_s - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right] \geq 0.$$

We notice that the first inequality in (3.9) is equivalent with

$$\sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{i=1}^n p_i x_i \right\|^2 \geq 2 \sum_{k=1}^n p_k \left\langle x_k, \sum_{i=1}^n p_i x_i \right\rangle_s$$

which provides the result

$$(3.10) \quad \frac{1}{2} \left[ \sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right] \geq \sum_{k=1}^n p_k \left\langle x_k, \sum_{i=1}^n p_i x_i \right\rangle_s \\ \left( \geq \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right)$$

for any  $n$ -tuple of vectors and probability distribution.

**Remark 3.** *If in the inequality (3.7) we consider the uniform distribution, then we get*

$$(3.11) \quad \sum_{i=1}^n \|x_i\|^p - n^{1-p} \left\| \sum_{i=1}^n x_i \right\|^p \\ \geq p n^{1-p} \left\| \sum_{i=1}^n x_i \right\|^{p-2} \left[ \sum_{k=1}^n \left\langle x_k, \sum_{i=1}^n x_i \right\rangle_s - \left\| \sum_{i=1}^n x_i \right\|^2 \right] \geq 0.$$

## 4. A REVERSE OF JENSEN'S INEQUALITY

The following result is of interest as well:

**Theorem 3.** *Let  $f : X \rightarrow \mathbb{R}$  be a convex function defined on a linear space  $X$ . Then for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in X^n$  and any probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$  we have the inequality*

$$(4.1) \quad \sum_{k=1}^n p_k \nabla_{-f}(x_k)(x_k) - \sum_{k=1}^n p_k \nabla_{-f}(x_k) \left( \sum_{i=1}^n p_i x_i \right) \\ \geq \sum_{i=1}^n p_i f(x_i) - f \left( \sum_{i=1}^n p_i x_i \right).$$

In particular, for the uniform distribution, we have

$$(4.2) \quad \frac{1}{n} \left[ \sum_{k=1}^n \nabla_{-f}(x_k)(x_k) - \sum_{k=1}^n \nabla_{-f}(x_k) \left( \frac{1}{n} \sum_{i=1}^n x_i \right) \right] \\ \geq \frac{1}{n} \sum_{i=1}^n f(x_i) - f \left( \frac{1}{n} \sum_{i=1}^n x_i \right).$$

*Proof.* Utilising the gradient inequality (2.2) we can state that

$$(4.3) \quad \nabla_{-f}(x_k) \left( x_k - \sum_{i=1}^n p_i x_i \right) \geq f(x_k) - f \left( \sum_{i=1}^n p_i x_i \right)$$

for any  $k \in \{1, \dots, n\}$ .

By the superadditivity of the functional  $\nabla_{-f}(\cdot)(\cdot)$  in the second variable we also have

$$(4.4) \quad \nabla_{-f}(x_k)(x_k) - \nabla_{-f}(x_k) \left( \sum_{i=1}^n p_i x_i \right) \geq \nabla_{-f}(x_k) \left( x_k - \sum_{i=1}^n p_i x_i \right)$$

for any  $k \in \{1, \dots, n\}$ .

Therefore, by (4.3) and (4.4) we get

$$(4.5) \quad \nabla_{-f}(x_k)(x_k) - \nabla_{-f}(x_k) \left( \sum_{i=1}^n p_i x_i \right) \geq f(x_k) - f \left( \sum_{i=1}^n p_i x_i \right)$$

for any  $k \in \{1, \dots, n\}$ .

Finally, by multiplying (4.5) with  $p_k \geq 0$  and summing over  $k$  from 1 to  $n$  we deduce the desired inequality (4.1).  $\square$

**Remark 4.** *If the function  $f$  is defined on the Euclidian space  $\mathbb{R}^n$  and is differentiable and convex, then from (4.1) we get the inequality*

$$(4.6) \quad \sum_{k=1}^n p_k \langle \nabla f(x_k), x_k \rangle - \left\langle \sum_{k=1}^n p_k \nabla f(x_k), \sum_{i=1}^n p_i x_i \right\rangle \\ \geq \sum_{i=1}^n p_i f(x_i) - f \left( \sum_{i=1}^n p_i x_i \right)$$



where, as usual, for  $x_k = (x_k^1, \dots, x_k^n)$ ,  $\nabla f(x_k) = \left( \frac{\partial f(x_k)}{\partial x_k^1}, \dots, \frac{\partial f(x_k)}{\partial x_k^n} \right)$ . This inequality was obtained firstly by Dragomir & Goh in 1996, see [12].

For one dimension we get the inequality

$$(4.7) \quad \sum_{k=1}^n p_k x_k f'(x_k) - \sum_{i=1}^n p_i x_i \sum_{k=1}^n p_k f'(x_k) \geq \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right)$$

that was discovered in 1994 by Dragomir and Ionescu, see [11].

The following reverse of the generalised triangle inequality holds:

**Corollary 3.** *Let  $(X, \|\cdot\|)$  be a normed linear space. Then for any  $p \geq 1$ , for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in X^n \setminus \{(0, \dots, 0)\}$  and any probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$  we have the inequality*

$$(4.8) \quad p \left[ \sum_{k=1}^n p_k \|x_k\|^p - \sum_{k=1}^n p_k \|x_k\|^{p-2} \left\langle \sum_{i=1}^n p_i x_i, x_k \right\rangle_i \right] \geq \sum_{i=1}^n p_i \|x_i\|^p - \left\| \sum_{i=1}^n p_i x_i \right\|^p.$$

In particular, we have the norm inequalities

$$(4.9) \quad \sum_{k=1}^n p_k \|x_k\| - \sum_{k=1}^n p_k \left\langle \sum_{i=1}^n p_i x_i, \frac{x_k}{\|x_k\|} \right\rangle_i \geq \sum_{i=1}^n p_i \|x_i\| - \left\| \sum_{i=1}^n p_i x_i \right\|$$

for  $x_k \neq 0, k \in \{1, \dots, n\}$  and

$$(4.10) \quad 2 \left[ \sum_{k=1}^n p_k \|x_k\|^2 - \sum_{k=1}^n p_k \left\langle \sum_{j=1}^n p_j x_j, x_k \right\rangle_i \right] \geq \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2,$$

for any  $x_k$ .

We observe that the inequality (4.10) is equivalent with

$$\sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{i=1}^n p_i x_i \right\|^2 \geq 2 \sum_{k=1}^n p_k \left\langle \sum_{j=1}^n p_j x_j, x_k \right\rangle_i$$

which provides the interesting result

$$(4.11) \quad \frac{1}{2} \left[ \sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right] \geq \sum_{k=1}^n p_k \left\langle \sum_{j=1}^n p_j x_j, x_k \right\rangle_i \\ \left( \geq \sum_{k=1}^n \sum_{j=1}^n p_j p_k \langle x_j, x_k \rangle_i \right)$$

holding for any  $n$ -tuple of vectors and probability distribution.

**Remark 5.** *If in the inequality (4.8) we consider the uniform distribution, then we get*

$$(4.12) \quad p \left[ \sum_{k=1}^n \|x_k\|^p - \frac{1}{n} \sum_{k=1}^n \|x_k\|^{p-2} \left\langle \sum_{j=1}^n x_j, x_k \right\rangle_i \right] \\ \geq \sum_{i=1}^n \|x_i\|^p - n^{1-p} \left\| \sum_{i=1}^n x_i \right\|^p.$$

For  $p \in [1, 2)$  all vectors  $x_k$  should not be zero.

## 5. BOUNDS FOR THE MEAN $f$ -DEVIATION

Let  $X$  be a real linear space. For a convex function  $f : X \rightarrow \mathbb{R}$  with the property that  $f(0) \geq 0$  we define the *mean  $f$ -deviation* of an  $n$ -tuple of vectors  $y = (y_1, \dots, y_n) \in X^n$  with the probability distribution  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{P}^n$  by the non-negative quantity

$$(5.1) \quad K_{f(\cdot)}(\mathbf{p}, \mathbf{y}) = K_f(\mathbf{p}, \mathbf{y}) := \sum_{i=1}^n p_i f \left( y_i - \sum_{k=1}^n p_k y_k \right).$$

The fact that  $K_f(\mathbf{p}, \mathbf{y})$  is non-negative follows by Jensen's inequality, namely

$$K_f(\mathbf{p}, \mathbf{y}) \geq f \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) = f(0) \geq 0.$$

Of course the concept can be extended for any function defined on  $X$ , however if the function is not convex or if it is convex but  $f(0) < 0$ , then we are not sure about the positivity of the quantity  $K_f(\mathbf{p}, \mathbf{y})$ .

A natural example of such deviations can be provided by the convex function  $f(y) := \|y\|^r$  with  $r \geq 1$  defined on a normed linear space  $(X, \|\cdot\|)$ . We denote this by

$$(5.2) \quad K_r(\mathbf{p}, \mathbf{y}) := \sum_{i=1}^n p_i \left\| y_i - \sum_{k=1}^n p_k y_k \right\|^r$$

and call it the *mean  $r$ -absolute deviation* of the  $n$ -tuple of vectors  $y = (y_1, \dots, y_n) \in X^n$  with the probability distribution  $\mathbf{p} = (p_1, \dots, p_n)$ .

Utilising the result from [9] we can state then the following result providing a non-trivial lower bound for the mean  $f$ -deviation:

**Theorem 4.** Let  $f : X \rightarrow [0, \infty)$  be a convex function with  $f(0) = 0$ . If  $y = (y_1, \dots, y_n) \in X^n$  and  $\mathbf{p} = (p_1, \dots, p_n)$  is a probability distribution with all  $p_i$  nonzero, then

$$(5.3) \quad K_f(\mathbf{p}, \mathbf{y}) \geq \max_{k \in \{1, \dots, n\}} \left\{ (1 - p_k) f \left[ \frac{p_k}{1 - p_k} \left( y_k - \sum_{l=1}^n p_l y_l \right) \right] + p_k f \left( y_k - \sum_{l=1}^n p_l y_l \right) \right\} (\geq 0).$$

The case for mean  $r$ -absolute deviation is incorporated in

**Corollary 4.** Let  $(X, \|\cdot\|)$  be a normed linear space. If  $y = (y_1, \dots, y_n) \in X^n$  and  $\mathbf{p} = (p_1, \dots, p_n)$  is a probability distribution with all  $p_i$  nonzero, then for  $r \geq 1$  we have

$$(5.4) \quad K_r(\mathbf{p}, \mathbf{y}) \geq \max_{k \in \{1, \dots, n\}} \left\{ \left[ (1 - p_k)^{1-r} p_k^r + p_k \right] \left\| y_k - \sum_{l=1}^n p_l y_l \right\|^r \right\}.$$

**Remark 6.** Since the function  $h_r(t) := (1 - t)^{1-r} t^r + t$ ,  $r \geq 1$ ,  $t \in [0, 1)$  is strictly increasing on  $[0, 1)$ , then

$$\min_{k \in \{1, \dots, n\}} \left\{ (1 - p_k)^{1-r} p_k^r + p_k \right\} = p_m + (1 - p_m)^{1-r} p_m^r,$$

where  $p_m := \min_{k \in \{1, \dots, n\}} p_k$ . By (5.4), we then obtain the following simpler inequality:

$$(5.5) \quad K_r(\mathbf{p}, \mathbf{y}) \geq \left[ p_m + (1 - p_m)^{1-r} \cdot p_m^r \right] \max_{k \in \{1, \dots, n\}} \left\| y_k - \sum_{l=1}^n p_l y_l \right\|^r,$$

which is perhaps more useful for applications.

We have the following double inequality for the mean  $f$ -mean deviation:

**Theorem 5.** Let  $f : X \rightarrow [0, \infty)$  be a convex function with  $f(0) = 0$ . If  $y = (y_1, \dots, y_n) \in X^n$  and  $\mathbf{p} = (p_1, \dots, p_n)$  is a probability distribution with all  $p_i$  nonzero, then

$$(5.6) \quad K_{\nabla_- f(\cdot)(\cdot)}(\mathbf{p}, \mathbf{y}) \geq K_{f(\cdot)}(\mathbf{p}, \mathbf{y}) \geq K_{\nabla_+ f(0)(\cdot)}(\mathbf{p}, \mathbf{y}) \geq 0.$$

*Proof.* If we use the inequality (3.2) for  $x_i = y_i - \sum_{k=1}^n p_k y_k$  we get

$$\begin{aligned} & \sum_{i=1}^n p_i f \left( y_i - \sum_{k=1}^n p_k y_k \right) - f \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) \\ & \geq \sum_{j=1}^n p_j \nabla_+ f \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) \left( y_j - \sum_{k=1}^n p_k y_k \right) \\ & \quad - \nabla_+ f \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) \geq 0 \end{aligned}$$

which is equivalent with the second part of (5.6).

Now, by utilising the inequality (4.1) for the same choice of  $x_i$  we get

$$\begin{aligned} & \sum_{j=1}^n p_j \nabla_- f \left( y_j - \sum_{k=1}^n p_k y_k \right) \left( y_j - \sum_{k=1}^n p_k y_k \right) \\ & \quad - \sum_{k=1}^n p_j \nabla_- f \left( y_j - \sum_{k=1}^n p_k y_k \right) \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right) \\ & \quad \geq \sum_{i=1}^n p_i f \left( y_i - \sum_{k=1}^n p_k y_k \right) - f \left( \sum_{i=1}^n p_i \left( y_i - \sum_{k=1}^n p_k y_k \right) \right), \end{aligned}$$

which in its turn is equivalent with the first inequality in (5.6).  $\square$

We observe that as examples of convex functions defined on the entire normed linear space  $(X, \|\cdot\|)$  that are convex and vanishes in 0 we can consider the functions

$$f(x) := g(\|x\|), \quad x \in X$$

where  $g : [0, \infty) \rightarrow [0, \infty)$  is a monotonic nondecreasing convex function with  $g(0) = 0$ .

For this kind of functions we have by direct computation that

$$\nabla_+ f(0)(u) = g'_+(0) \|u\| \quad \text{for any } u \in X$$

and

$$\nabla_- f(u)(u) = g'_-(\|u\|) \|u\| \quad \text{for any } u \in X.$$

We then have the following norm inequalities that are of interest:

**Corollary 5.** *Let  $(X, \|\cdot\|)$  be a normed linear space. If  $g : [0, \infty) \rightarrow [0, \infty)$  is a monotonic nondecreasing convex function with  $g(0) = 0$ , then for any  $y = (y_1, \dots, y_n) \in X^n$  and  $\mathbf{p} = (p_1, \dots, p_n)$  a probability distribution, we have*

$$(5.7) \quad \begin{aligned} & \sum_{i=1}^n p_i g'_- \left( \left\| y_i - \sum_{k=1}^n p_k y_k \right\| \right) \left\| y_i - \sum_{k=1}^n p_k y_k \right\| \\ & \quad \geq \sum_{i=1}^n p_i g \left( \left\| y_i - \sum_{k=1}^n p_k y_k \right\| \right) \geq g'_+(0) \sum_{i=1}^n p_i \left\| y_i - \sum_{k=1}^n p_k y_k \right\|. \end{aligned}$$

## 6. BOUNDS FOR $f$ -DIVERGENCE MEASURES

Given a convex function  $f : [0, \infty) \rightarrow \mathbb{R}$ , the  $f$ -divergence functional

$$(6.1) \quad I_f(\mathbf{p}, \mathbf{q}) := \sum_{i=1}^n q_i f \left( \frac{p_i}{q_i} \right),$$

where  $\mathbf{p} = (p_1, \dots, p_n)$ ,  $\mathbf{q} = (q_1, \dots, q_n)$  are positive sequences, was introduced by Csiszár in [1], as a generalized measure of information, a ‘‘distance function’’ on the set of probability distributions  $\mathbb{P}^n$ . As in [1], we interpret undefined expressions by

$$\begin{aligned} f(0) &= \lim_{t \rightarrow 0^+} f(t), & 0f\left(\frac{0}{0}\right) &= 0, \\ 0f\left(\frac{a}{0}\right) &= \lim_{q \rightarrow 0^+} qf\left(\frac{a}{q}\right) = a \lim_{t \rightarrow \infty} \frac{f(t)}{t}, & a &> 0. \end{aligned}$$

The following results were essentially given by Csiszár and Körner [2]:

- (i) If  $f$  is convex, then  $I_f(\mathbf{p}, \mathbf{q})$  is jointly convex in  $\mathbf{p}$  and  $\mathbf{q}$ ;  
 (ii) For every  $\mathbf{p}, \mathbf{q} \in R_+^n$ , we have

$$(6.2) \quad I_f(\mathbf{p}, \mathbf{q}) \geq \sum_{j=1}^n q_j f\left(\frac{\sum_{j=1}^n p_j}{\sum_{j=1}^n q_j}\right).$$

If  $f$  is strictly convex, equality holds in (6.2) iff

$$\frac{p_1}{q_1} = \frac{p_2}{q_2} = \dots = \frac{p_n}{q_n}.$$

If  $f$  is normalized, i.e.,  $f(1) = 0$ , then for every  $\mathbf{p}, \mathbf{q} \in R_+^n$  with  $\sum_{i=1}^n p_i = \sum_{i=1}^n q_i$ , we have the inequality

$$(6.3) \quad I_f(\mathbf{p}, \mathbf{q}) \geq 0.$$

In particular, if  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$ , then (6.3) holds. This is the well-known positivity property of the  $f$ -divergence.

We endeavour to extend this concept for functions defined on a cone in a linear space as follows (see also [10]).

Firstly, we recall that the subset  $K$  in a linear space  $X$  is a *cone* if the following two conditions are satisfied:

- (i) for any  $x, y \in K$  we have  $x + y \in K$ ;  
 (ii) for any  $x \in K$  and any  $\alpha \geq 0$  we have  $\alpha x \in K$ .

For a given  $n$ -tuple of vectors  $\mathbf{z} = (z_1, \dots, z_n) \in K^n$  and a probability distribution  $\mathbf{q} \in \mathbb{P}^n$  with all values nonzero, we can define, for the convex function  $f : K \rightarrow \mathbb{R}$ , the following  *$f$ -divergence of  $\mathbf{z}$  with the distribution  $\mathbf{q}$*

$$(6.4) \quad I_f(\mathbf{z}, \mathbf{q}) := \sum_{i=1}^n q_i f\left(\frac{z_i}{q_i}\right).$$

It is obvious that if  $X = \mathbb{R}$ ,  $K = [0, \infty)$  and  $\mathbf{x} = \mathbf{p} \in \mathbb{P}^n$  then we obtain the usual concept of the  $f$ -divergence associated with a function  $f : [0, \infty) \rightarrow \mathbb{R}$ .

Now, for a given  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in K^n$ , a probability distribution  $\mathbf{q} \in \mathbb{P}^n$  with all values nonzero and for any nonempty subset  $J$  of  $\{1, \dots, n\}$  we have

$$\mathbf{q}_J := (Q_J, \bar{Q}_J) \in \mathbb{P}^2$$

and

$$\mathbf{x}_J := (X_J, \bar{X}_J) \in K^2$$

where, as above

$$X_J := \sum_{i \in J} x_i, \quad \text{and} \quad \bar{X}_J := X_{\bar{J}}.$$

It is obvious that

$$I_f(\mathbf{x}_J, \mathbf{q}_J) = Q_J f\left(\frac{X_J}{Q_J}\right) + \bar{Q}_J f\left(\frac{\bar{X}_J}{\bar{Q}_J}\right).$$

The following inequality for the  $f$ -divergence of an  $n$ -tuple of vectors in a linear space holds [10]:

**Theorem 6.** *Let  $f : K \rightarrow \mathbb{R}$  be a convex function on the cone  $K$ . Then for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in K^n$ , a probability distribution  $\mathbf{q} \in \mathbb{P}^n$  with all values nonzero and for any nonempty subset  $J$  of  $\{1, \dots, n\}$  we have*

$$(6.5) \quad \begin{aligned} I_f(\mathbf{x}, \mathbf{q}) &\geq \max_{\emptyset \neq J \subset \{1, \dots, n\}} I_f(\mathbf{x}_J, \mathbf{q}_J) \geq I_f(\mathbf{x}_J, \mathbf{q}_J) \\ &\geq \min_{\emptyset \neq J \subset \{1, \dots, n\}} I_f(\mathbf{x}_J, \mathbf{q}_J) \geq f(X_n) \end{aligned}$$

where  $X_n := \sum_{i=1}^n x_i$ .

We observe that, for a given  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in K^n$ , a sufficient condition for the positivity of  $I_f(\mathbf{x}, \mathbf{q})$  for any probability distribution  $\mathbf{q} \in \mathbb{P}^n$  with all values nonzero is that  $f(X_n) \geq 0$ . In the scalar case and if  $\mathbf{x} = \mathbf{p} \in \mathbb{P}^n$ , then a sufficient condition for the positivity of the  $f$ -divergence  $I_f(\mathbf{p}, \mathbf{q})$  is that  $f(1) \geq 0$ .

The case of functions of a real variable that is of interest for applications is incorporated in [10]:

**Corollary 6.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a normalized convex function. Then for any  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$  we have*

$$(6.6) \quad I_f(\mathbf{p}, \mathbf{q}) \geq \max_{\emptyset \neq J \subset \{1, \dots, n\}} \left[ Q_J f\left(\frac{P_J}{Q_J}\right) + (1 - Q_J) f\left(\frac{1 - P_J}{1 - Q_J}\right) \right] (\geq 0).$$

In what follows, by the use of the results in Theorem 2 and Theorem 3 we can provide an upper and a lower bound for the positive difference  $I_f(\mathbf{x}, \mathbf{q}) - f(X_n)$ .

**Theorem 7.** *Let  $f : K \rightarrow \mathbb{R}$  be a convex function on the cone  $K$ . Then for any  $n$ -tuple of vectors  $\mathbf{x} = (x_1, \dots, x_n) \in K^n$  and a probability distribution  $\mathbf{q} \in \mathbb{P}^n$  with all values nonzero we have*

$$(6.7) \quad \begin{aligned} I_{\nabla - f(\cdot)(\cdot)}(\mathbf{x}, \mathbf{q}) - I_{\nabla - f(\cdot)(X_n)}(\mathbf{x}, \mathbf{q}) &\geq I_f(\mathbf{x}, \mathbf{q}) - f(X_n) \\ &\geq I_{\nabla + f(X_n)(\cdot)}(\mathbf{x}, \mathbf{q}) - \nabla + f(X_n)(X_n) \geq 0. \end{aligned}$$

The case of functions of a real variable that is useful for applications is as follows:

**Corollary 7.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a normalized convex function. Then for any  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$  we have*

$$(6.8) \quad I_{f'_-(\cdot)(\cdot)}(\mathbf{p}, \mathbf{q}) - I_{f'_-(\cdot)}(\mathbf{p}, \mathbf{q}) \geq I_f(\mathbf{p}, \mathbf{q}) \geq 0,$$

or, equivalently,

$$(6.9) \quad I_{f'_-(\cdot)[(\cdot)-1]}(\mathbf{p}, \mathbf{q}) \geq I_f(\mathbf{p}, \mathbf{q}) \geq 0.$$

The above corollary is useful to provide an upper bound in terms of the variational distance for the  $f$ -divergence  $I_f(\mathbf{p}, \mathbf{q})$  of normalized convex functions whose derivatives are bounded above and below.

**Proposition 1.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a normalized convex function and  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$ . If there exists the constants  $\gamma$  and  $\Gamma$  with*

$$-\infty < \gamma \leq f'_- \left( \frac{p_k}{q_k} \right) \leq \Gamma < \infty \text{ for all } k \in \{1, \dots, n\},$$

then we have the inequality

$$(6.10) \quad 0 \leq I_f(\mathbf{p}, \mathbf{q}) \leq \frac{1}{2} (\Gamma - \gamma) V(\mathbf{p}, \mathbf{q}),$$

where  $V(\mathbf{p}, \mathbf{q}) = \sum_{i=1}^n q_i \left| \frac{p_i}{q_i} - 1 \right| = \sum_{i=1}^n |p_i - q_i|$ .

*Proof.* By the inequality (6.9) we have successively that

$$\begin{aligned}
 0 &\leq I_f(\mathbf{p}, \mathbf{q}) \leq I_{f'_- (\cdot)[(\cdot)-1]}(\mathbf{p}, \mathbf{q}) \\
 &= \sum_{i=1}^n q_i \left( \frac{p_i}{q_i} - 1 \right) \left[ f'_- \left( \frac{p_i}{q_i} \right) - \frac{\Gamma + \gamma}{2} \right] \\
 &\leq \sum_{i=1}^n q_i \left| \frac{p_i}{q_i} - 1 \right| \left| f'_- \left( \frac{p_i}{q_i} \right) - \frac{\Gamma + \gamma}{2} \right| \\
 &\leq \frac{1}{2} (\Gamma - \gamma) \sum_{i=1}^n q_i \left| \frac{p_i}{q_i} - 1 \right|
 \end{aligned}$$

which proves the desired result (6.10).  $\square$

**Corollary 8.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a normalized convex function and  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$ . If there exist the constants  $r$  and  $R$  with*

$$0 < r \leq \frac{p_k}{q_k} \leq R < \infty \text{ for all } k \in \{1, \dots, n\},$$

*then we have the inequality*

$$(6.11) \quad 0 \leq I_f(\mathbf{p}, \mathbf{q}) \leq \frac{1}{2} [f'_-(R) - f'_-(r)] V(\mathbf{p}, \mathbf{q}).$$

The K. Pearson  $\chi^2$ -divergence is obtained for the convex function  $f(t) = (1-t)^2$ ,  $t \in \mathbb{R}$  and given by

$$\chi^2(p, q) := \sum_{j=1}^n q_j \left( \frac{p_j}{q_j} - 1 \right)^2 = \sum_{j=1}^n \frac{(p_j - q_j)^2}{q_j}.$$

Finally, the following proposition giving another upper bound in terms of the  $\chi^2$ -divergence can be stated:

**Proposition 2.** *Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be a normalized convex function and  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$ . If there exists the constant  $0 < \Delta < \infty$  with*

$$(6.12) \quad \left| \frac{f'_- \left( \frac{p_i}{q_i} \right) - f'_-(1)}{\frac{p_i}{q_i} - 1} \right| \leq \Delta \text{ for all } k \in \{1, \dots, n\},$$

*then we have the inequality*

$$(6.13) \quad 0 \leq I_f(\mathbf{p}, \mathbf{q}) \leq \Delta \chi^2(p, q).$$

*In particular, if  $f'_-(\cdot)$  satisfies the local Lipschitz condition*

$$(6.14) \quad |f'_-(x) - f'_-(1)| \leq \Delta |x - 1| \text{ for any } x \in (0, \infty)$$

*then (6.13) holds true for any  $\mathbf{p}, \mathbf{q} \in \mathbb{P}^n$ .*

*Proof.* We have from (6.9) that

$$\begin{aligned} 0 &\leq I_f(\mathbf{p}, \mathbf{q}) \leq I_{f'_- (\cdot)[(\cdot)-1]}(\mathbf{p}, \mathbf{q}) \\ &= \sum_{i=1}^n q_i \left( \frac{p_i}{q_i} - 1 \right) \left[ f'_- \left( \frac{p_i}{q_i} \right) - f'_-(1) \right] \\ &\leq \sum_{i=1}^n q_i \left( \frac{p_i}{q_i} - 1 \right)^2 \left| \frac{f'_- \left( \frac{p_i}{q_i} \right) - f'_-(1)}{\frac{p_i}{q_i} - 1} \right| \\ &\leq \Delta \sum_{i=1}^n q_i \left( \frac{p_i}{q_i} - 1 \right)^2 \end{aligned}$$

and the inequality (6.13) is obtained.  $\square$

**Remark 7.** *It is obvious that if one chooses in the above inequalities particular normalized convex functions that generates the Kullback-Leibler, Jeffreys, Hellinger or other divergence measures or discrepancies, that one can obtain some results of interest. However the details are not provided here.*

#### REFERENCES

- [1] I. Csiszár, Information-type measures of differences of probability distributions and indirect observations, *Studia Sci. Math. Hung.*, **2** (1967), 299-318.
- [2] I. Csiszár and J. Körner, *Information Theory: Coding Theorems for Discrete Memoryless Systems*, Academic Press, New York, 1981.
- [3] S.S. Dragomir, An improvement of Jensen's inequality, *Bull. Math. Soc. Sci. Math. Roumanie*, **34(82)** (1990), No. 4, 291-296.
- [4] S.S. Dragomir, Some refinements of Ky Fan's inequality, *J. Math. Anal. Appl.*, **163**(2) (1992), 317-321.
- [5] S.S. Dragomir, Some refinements of Jensen's inequality, *J. Math. Anal. Appl.*, **168**(2) (1992), 518-522.
- [6] S.S. Dragomir, A further improvement of Jensen's inequality, *Tamkang J. Math.*, **25**(1) (1994), 29-36.
- [7] S.S. Dragomir, A new improvement of Jensen's inequality, *Indian J. Pure and Appl. Math.*, **26**(10) (1995), 959-968.
- [8] S.S. Dragomir, *Semi-inner Products and Applications*, Nova Science Publishers Inc., NY, 2004.
- [9] S.S. Dragomir, A refinement of Jensen's inequality with applications for  $f$ -divergence measures, *Taiwanese J. Maths.* (in press).
- [10] S.S. Dragomir, A new refinement of Jensen's inequality in linear spaces with applications, Preprint *RGMA Res. Rep. Coll.* **12**(2009), Supplement, Article 6. [Online [http://www.staff.vu.edu.au/RGMA/v12\(E\).asp](http://www.staff.vu.edu.au/RGMA/v12(E).asp)].
- [11] S.S. Dragomir and N.M. Ionescu, Some converse of Jensen's inequality and applications. *Rev. Anal. Numér. Théor. Approx.* **23** (1994), no. 1, 71-78. MR1325895 (96c:26012).
- [12] S.S. Dragomir and C.J. Goh, A counterpart of Jensen's discrete inequality for differential-ble convex mappings and applications in information theory, *Mathl. Comput. Modelling*, **24**(1996), No. 2, 1-11.
- [13] S.S. Dragomir, J. Pečarić and L.E. Persson, Properties of some functionals related to Jensen's inequality, *Acta Math. Hung.*, **70**(1-2) (1996), 129-143.
- [14] J. Pečarić and S.S. Dragomir, A refinements of Jensen inequality and applications, *Studia Univ. Babeş-Bolyai, Mathematica*, **24**(1) (1989), 15-19.

RESEARCH GROUP IN MATHEMATICAL INEQUALITIES & APPLICATIONS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: [sever.dragomir@vu.edu.au](mailto:sever.dragomir@vu.edu.au)

URL: <http://www.staff.vu.edu.au/rgmia/dragomir/>