

**VECTOR AND OPERATOR TRAPEZOIDAL TYPE
INEQUALITIES FOR CONTINUOUS FUNCTIONS OF
SELFADJOINT OPERATORS IN HILBERT SPACES**

S.S. DRAGOMIR

ABSTRACT. Some vector and operator generalized trapezoidal inequalities for continuous functions of selfadjoint operators in Hilbert spaces are given. Applications for power and logarithmic functions of operators are provided as well.

1. INTRODUCTION

Let A be a selfadjoint linear operator on a complex Hilbert space $(H; \langle \cdot, \cdot \rangle)$. The *Gelfand map* establishes a $*$ -isometrically isomorphism Φ between the set $C(Sp(A))$ of all *continuous functions* defined on the *spectrum* of A , denoted $Sp(A)$, and the C^* -algebra $C^*(A)$ generated by A and the identity operator 1_H on H as follows (see for instance [10, p. 3]):

For any $f, g \in C(Sp(A))$ and any $\alpha, \beta \in \mathbb{C}$ we have

- (i) $\Phi(\alpha f + \beta g) = \alpha \Phi(f) + \beta \Phi(g)$;
- (ii) $\Phi(fg) = \Phi(f)\Phi(g)$ and $\Phi(\bar{f}) = \Phi(f)^*$;
- (iii) $\|\Phi(f)\| = \|f\| := \sup_{t \in Sp(A)} |f(t)|$;
- (iv) $\Phi(f_0) = 1_H$ and $\Phi(f_1) = A$, where $f_0(t) = 1$ and $f_1(t) = t$, for $t \in Sp(A)$.

With this notation we define

$$f(A) := \Phi(f) \text{ for all } f \in C(Sp(A))$$

and we call it the *continuous functional calculus* for a selfadjoint operator A .

If A is a selfadjoint operator and f is a real valued continuous function on $Sp(A)$, then $f(t) \geq 0$ for any $t \in Sp(A)$ implies that $f(A) \geq 0$, *i.e.* $f(A)$ is a *positive operator* on H . Moreover, if both f and g are real valued functions on $Sp(A)$ then the following important property holds:

$$(P) \quad f(t) \geq g(t) \text{ for any } t \in Sp(A) \text{ implies that } f(A) \geq g(A)$$

in the operator order of $B(H)$.

For a recent monograph devoted to various inequalities for continuous functions of selfadjoint operators, see [10] and the references therein.

For other recent results see [5], [6], [7], [12], [13], [14] and [15].

Let U be a selfadjoint operator on the complex Hilbert space $(H, \langle \cdot, \cdot \rangle)$ with the spectrum $Sp(U)$ included in the interval $[m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its *spectral family*. Then for any continuous function $f : [m, M] \rightarrow \mathbb{C}$,

1991 *Mathematics Subject Classification.* 47A63; 47A99.

Key words and phrases. Selfadjoint operators, Functions of Selfadjoint operators, Spectral representation, Inequalities for selfadjoint operators.

it is well known that we have the following *spectral representation in terms of the Riemann-Stieltjes integral*:

$$(1.1) \quad \langle f(U)x, y \rangle = \int_{m-0}^M f(\lambda) d(\langle E_\lambda x, y \rangle),$$

for any $x, y \in H$. The function $g_{x,y}(\lambda) := \langle E_\lambda x, y \rangle$ is of *bounded variation* on the interval $[m, M]$ and

$$g_{x,y}(m-0) = 0 \text{ and } g_{x,y}(M) = \langle x, y \rangle$$

for any $x, y \in H$. It is also well known that $g_x(\lambda) := \langle E_\lambda x, x \rangle$ is *monotonic nondecreasing* and *right continuous* on $[m, M]$.

With the notations introduced above, we have considered in the recent paper [8] the problem of bounding the error

$$\frac{f(M) + f(m)}{2} \cdot \langle x, y \rangle - \langle f(A)x, y \rangle$$

in approximating $\langle f(A)x, y \rangle$ by the trapezoidal type formula $\frac{f(M)+f(m)}{2} \cdot \langle x, y \rangle$, where x, y are vectors in the Hilbert space H and f is a continuous functions of the selfadjoint operator A with the spectrum in the compact interval of real numbers $[m, M]$.

We recall here only two such results. The first deals with the case of continuous functions of bounded variation and is incorporated in the following theorem [8]:

Theorem 1. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{C}$ is a continuous function of bounded variation on $[m, M]$, then we have the inequality*

$$(1.2) \quad \begin{aligned} & \left| \frac{f(M) + f(m)}{2} \cdot \langle x, y \rangle - \langle f(A)x, y \rangle \right| \\ & \leq \frac{1}{2} \max_{\lambda \in [m, M]} \left[\langle E_\lambda x, x \rangle^{1/2} \langle E_\lambda y, y \rangle^{1/2} \right. \\ & \quad \left. + \langle (1_H - E_\lambda)x, x \rangle^{1/2} \langle (1_H - E_\lambda)y, y \rangle^{1/2} \right] \bigvee_m^M(f) \\ & \leq \frac{1}{2} \|x\| \|y\| \bigvee_m^M(f) \end{aligned}$$

for any $x, y \in H$.

The case of Lipschitzian functions is as follows [8]:

Theorem 2. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{C}$ is Lipschitzian with the constant $L > 0$ on $[m, M]$, then*

we have the inequality

$$\begin{aligned}
(1.3) \quad & \left| \frac{f(M) + f(m)}{2} \cdot \langle x, y \rangle - \langle f(A)x, y \rangle \right| \\
& \leq \frac{1}{2} L \int_{m-0}^M \left[\langle E_\lambda x, x \rangle^{1/2} \langle E_\lambda y, y \rangle^{1/2} \right. \\
& \quad \left. + \langle (1_H - E_\lambda)x, x \rangle^{1/2} \langle (1_H - E_\lambda)y, y \rangle^{1/2} \right] d\lambda \\
& \leq \frac{1}{2} (M - m) L \|x\| \|y\|
\end{aligned}$$

for any $x, y \in H$.

In order to provide error bounds in approximating $\langle f(A)x, y \rangle$ with the quantity

$$\frac{1}{M - m} [f(m)(M \langle x, y \rangle - \langle Ax, y \rangle) + f(M)(\langle Ax, y \rangle - m \langle x, y \rangle)]$$

where $x, y \in H$, which is a generalized trapezoid formula, we obtained in the recent paper [9] the following results:

Theorem 3. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family.*

1. *If $f : [m, M] \rightarrow \mathbb{C}$ is of bounded variation on $[m, M]$, then*

$$\begin{aligned}
(1.4) \quad & \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\
& \leq \sup_{t \in [m, M]} \left[\frac{t - m}{M - m} \bigvee_{m-0}^t (\langle E_{(\cdot)} x, y \rangle) + \frac{M - t}{M - m} \bigvee_t^M (\langle E_{(\cdot)} x, y \rangle) \right] \bigvee_m^M (f) \\
& \leq \bigvee_{m-0}^M (\langle E_{(\cdot)} x, y \rangle) \bigvee_m^M (f) \leq \|x\| \|y\| \bigvee_m^M (f)
\end{aligned}$$

for any $x, y \in H$.

2. *If $f : [m, M] \rightarrow \mathbb{C}$ is Lipschitzian with the constant $L > 0$ on $[m, M]$, then*

$$\begin{aligned}
(1.5) \quad & \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\
& \leq L \int_{m-0}^M \left[\frac{t - m}{M - m} \bigvee_{m-0}^t (\langle E_{(\cdot)} x, y \rangle) + \frac{M - t}{M - m} \bigvee_t^M (\langle E_{(\cdot)} x, y \rangle) \right] dt \\
& \leq L (M - m) \bigvee_{m-0}^M (\langle E_{(\cdot)} x, y \rangle) \leq L (M - m) \|x\| \|y\|
\end{aligned}$$

for any $x, y \in H$.

3. If $f : [m, M] \rightarrow \mathbb{R}$ is monotonic nondecreasing on $[m, M]$, then

$$(1.6) \quad \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\ \leq \int_{m-0}^M \left[\frac{t-m}{M-m} \bigvee_{m-0}^t (\langle E_{(\cdot)} x, y \rangle) + \frac{M-t}{M-m} \bigvee_t^M (\langle E_{(\cdot)} x, y \rangle) \right] df(t) \\ \leq \bigvee_{m-0}^M (\langle E_{(\cdot)} x, y \rangle) [f(M) - f(m)] \leq \|x\| \|y\| [f(M) - f(m)]$$

for any $x, y \in H$.

For other results of this type see [9]. For scalar trapezoidal inequalities, see [1]-[4].

The main aim of this paper is to provide more results of the above type as well as bounds in the operator order for the selfadjoint linear operator

$$\left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \right|$$

where f is a continuous function of bounded variation, Lipschitzian or convex on $[m, M]$. Applications for certain functions of large interest such as the power and logarithmic functions are provided as well.

2. SOME NEW VECTOR INEQUALITIES

The following result for general continuous functions holds:

Theorem 4. Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{R}$ is continuous on $[m, M]$, then we have the inequalities:

$$(2.1) \quad \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\ \leq \left[\max_{t \in [m, M]} f(t) - \min_{t \in [m, M]} f(t) \right] \bigvee_m^M (\langle E_{(\cdot)} x, y \rangle) \\ \leq \left[\max_{t \in [m, M]} f(t) - \min_{t \in [m, M]} f(t) \right] \|x\| \|y\|$$

for any $x, y \in H$.

Proof. We observe that, by the spectral representation (1.1), we have the equality

$$(2.2) \quad \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \\ = \int_{m-0}^M \Phi_f(t) d(\langle E_t x, y \rangle)$$

for any $x, y \in H$, where $\Phi_f : [m, M] \rightarrow \mathbb{R}$ is given by

$$\Phi_f(t) = \frac{1}{M - m} [(M - t)f(m) + (t - m)f(M)] - f(t).$$

It is well known that if $p : [a, b] \rightarrow \mathbb{C}$ is a continuous function and $v : [a, b] \rightarrow \mathbb{C}$ is of bounded variation, then the Riemann-Stieltjes integral $\int_a^b p(t) dv(t)$ exists and the following inequality holds

$$(2.3) \quad \left| \int_a^b p(t) dv(t) \right| \leq \sup_{t \in [a, b]} |p(t)| \bigvee_a^b(v),$$

where $\bigvee_a^b(v)$ denotes the total variation of v on $[a, b]$.

Now, if we denote by $\gamma := \min_{t \in [m, M]} f(t)$ and by $\Gamma := \max_{t \in [m, M]} f(t)$ then we have

$$\begin{aligned} \gamma(M-t) &\leq (M-t)f(m) \leq \Gamma(M-t), \\ \gamma(t-m) &\leq (t-m)f(M) \leq \Gamma(t-m) \end{aligned}$$

and

$$-(M-m)\Gamma \leq -(M-m)f(t) \leq -\gamma(M-m)$$

for any $t \in [m, M]$. If we add these three inequalities, then we get

$$-(M-m)(\Gamma - \gamma) \leq (M-m)\Phi_f(t) \leq (M-m)(\Gamma - \gamma)$$

for any $t \in [m, M]$, which shows that

$$(2.4) \quad |\Phi_f(t)| \leq \Gamma - \gamma \text{ for any } t \in [m, M].$$

On applying the inequality (2.3) for the representation (2.2) we have from (2.4) that

$$\left| \int_{m-0}^M \Phi_f(t) d(\langle E_t x, y \rangle) \right| \leq (\Gamma - \gamma) \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle)$$

for any $x, y \in H$, which proves the first part of (2.1).

If P is a nonnegative operator on H , i.e., $\langle Px, x \rangle \geq 0$ for any $x \in H$, then the following inequality is a generalization of the Schwarz inequality in H

$$(2.5) \quad |\langle Px, y \rangle|^2 \leq \langle Px, x \rangle \langle Py, y \rangle$$

for any $x, y \in H$.

Now, if $d : m = t_0 < t_1 < \dots < t_{n-1} < t_n = M$ is an arbitrary partition of the interval $[m, M]$, then we have by Schwarz's inequality for nonnegative operators that

$$\begin{aligned} & \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle) \\ &= \sup_d \left\{ \sum_{i=0}^{n-1} |\langle (E_{t_{i+1}} - E_{t_i}) x, y \rangle| \right\} \\ &\leq \sup_d \left\{ \sum_{i=0}^{n-1} \left[\langle (E_{t_{i+1}} - E_{t_i}) x, x \rangle^{1/2} \langle (E_{t_{i+1}} - E_{t_i}) y, y \rangle^{1/2} \right] \right\} := I. \end{aligned}$$

By the Cauchy-Buniakovski-Schwarz inequality for sequences of real numbers we also have that

$$\begin{aligned}
I &\leq \sup_d \left\{ \left[\sum_{i=0}^{n-1} \langle (E_{t_{i+1}} - E_{t_i}) x, x \rangle \right]^{1/2} \left[\sum_{i=0}^{n-1} \langle (E_{t_{i+1}} - E_{t_i}) y, y \rangle \right]^{1/2} \right\} \\
&\leq \sup_d \left\{ \left[\sum_{i=0}^{n-1} \langle (E_{t_{i+1}} - E_{t_i}) x, x \rangle \right]^{1/2} \sup_d \left[\sum_{i=0}^{n-1} \langle (E_{t_{i+1}} - E_{t_i}) y, y \rangle \right]^{1/2} \right\} \\
&= \left[\bigvee_m^M (\langle E_{(\cdot)} x, x \rangle) \right]^{1/2} \left[\bigvee_m^M (\langle E_{(\cdot)} y, y \rangle) \right]^{1/2} = \|x\| \|y\|
\end{aligned}$$

for any $x, y \in H$.

These prove the last part of (2.1). \square

When the generating function is of bounded variation, we have the following result that complements Theorem 3:

Theorem 5. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{C}$ is continuous and of bounded variation on $[m, M]$, then we have the inequalities:*

$$\begin{aligned}
(2.6) \quad &\left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\
&\leq \max_{t \in [m, M]} \left[\frac{M - t}{M - m} \bigvee_m^t (f) + \frac{t - m}{M - m} \bigvee_t^M (f) \right] \bigvee_m^M (\langle E_{(\cdot)} x, y \rangle) \\
&\leq \bigvee_m^M (\langle E_{(\cdot)} x, y \rangle) \bigvee_m^M (f) \leq \bigvee_m^M (f) \|x\| \|y\|
\end{aligned}$$

for any $x, y \in H$.

Proof. First of all, observe that

$$\begin{aligned}
(2.7) \quad (M - m) \Phi_f(t) &= (t - M) [f(t) - f(m)] + (t - m) [f(M) - f(t)] \\
&= (t - M) \int_m^t df(s) + (t - m) \int_t^M df(s)
\end{aligned}$$

for any $t \in [m, M]$.

Therefore

$$\begin{aligned}
(2.8) \quad |\Phi_f(t)| &\leq \frac{M-t}{M-m} \left| \int_m^t df(s) \right| + \frac{t-m}{M-m} \left| \int_t^M df(s) \right| \\
&\leq \frac{M-t}{M-m} \bigvee_m^t(f) + \frac{t-m}{M-m} \bigvee_t^M(f) \\
&\leq \max \left\{ \frac{M-t}{M-m}, \frac{t-m}{M-m} \right\} \left[\bigvee_m^t(f) + \bigvee_t^M(f) \right] \\
&= \left[\frac{1}{2} + \frac{|t - \frac{m+M}{2}|}{M-m} \right] \bigvee_m^M(f)
\end{aligned}$$

for any $t \in [m, M]$, which implies that

$$\begin{aligned}
(2.9) \quad \max_{t \in [m, M]} |\Phi_f(t)| &\leq \max_{t \in [m, M]} \left[\frac{M-t}{M-m} \bigvee_m^t(f) + \frac{t-m}{M-m} \bigvee_t^M(f) \right] \\
&\leq \max_{t \in [m, M]} \left[\frac{1}{2} + \frac{|t - \frac{m+M}{2}|}{M-m} \right] \bigvee_m^M(f) = \bigvee_m^M(f).
\end{aligned}$$

On applying the inequality (2.3) for the representation (2.2) we have from (2.9) that

$$\begin{aligned}
&\left| \int_{m-0}^M \Phi_f(t) d(\langle E_t x, y \rangle) \right| \\
&\leq \max_{t \in [m, M]} \left[\frac{M-t}{M-m} \bigvee_m^t(f) + \frac{t-m}{M-m} \bigvee_t^M(f) \right] \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle) \\
&\leq \bigvee_m^M(f) \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle)
\end{aligned}$$

for any $x, y \in H$, which produces the desired result (2.6). \square

The case of Lipschitzian functions is as follows:

Theorem 6. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{C}$ is Lipschitzian with the constant $L > 0$ on $[m, M]$, then we have the inequalities:*

$$\begin{aligned}
(2.10) \quad &\left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M-m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\
&\leq \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle) \\
&\times \max_{t \in [m, M]} \left[\frac{M-t}{M-m} |f(t) - f(m)| + \frac{t-m}{M-m} |f(M) - f(t)| \right] \\
&\leq \frac{1}{2} (M-m) L \bigvee_m^M(\langle E_{(\cdot)} x, y \rangle) \leq \frac{1}{2} (M-m) L \|x\| \|y\|
\end{aligned}$$

for any $x, y \in H$.

Proof. We have from the first part of the equality (2.7) that

$$(2.11) \quad \begin{aligned} |\Phi_f(t)| &\leq \frac{M-t}{M-m} |f(t) - f(m)| + \frac{t-m}{M-m} |f(M) - f(t)| \\ &\leq \frac{2L}{M-m} (M-t)(t-m) \leq \frac{1}{2} (M-m)L \end{aligned}$$

for any $t \in [m, M]$, which, by a similar argument to the one from the above Theorem 5, produces the desired result (2.10). The details are omitted. \square

The following lemma may be stated.

Lemma 1. *Let $u : [a, b] \rightarrow \mathbb{R}$ and $\varphi, \Phi \in \mathbb{R}$ be such that $\Phi > \varphi$. The following statements are equivalent:*

- (i) *The function $u - \frac{\varphi + \Phi}{2} \cdot e$, where $e(t) = t$, $t \in [a, b]$, is $\frac{1}{2}(\Phi - \varphi)$ -Lipschitzian;*
- (ii) *We have the inequality:*

$$(2.12) \quad \varphi \leq \frac{u(t) - u(s)}{t - s} \leq \Phi \quad \text{for each } t, s \in [a, b] \quad \text{with } t \neq s;$$

- (iii) *We have the inequality:*

$$(2.13) \quad \varphi(t - s) \leq u(t) - u(s) \leq \Phi(t - s) \quad \text{for each } t, s \in [a, b] \quad \text{with } t > s.$$

Following [11], we can introduce the concept:

Definition 1. *The function $u : [a, b] \rightarrow \mathbb{R}$ which satisfies one of the equivalent conditions (i) – (iii) is said to be (φ, Φ) -Lipschitzian on $[a, b]$.*

Notice that in [11], the definition was introduced on utilizing the statement (iii) and only the equivalence (i) \Leftrightarrow (iii) was considered.

Utilising *Lagrange's mean value theorem*, we can state the following result that provides practical examples of (φ, Φ) -Lipschitzian functions.

Proposition 1. *Let $u : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) . If*

$$(2.14) \quad -\infty < \gamma := \inf_{t \in (a, b)} u'(t), \quad \sup_{t \in (a, b)} u'(t) =: \Gamma < \infty$$

then u is (γ, Γ) -Lipschitzian on $[a, b]$.

The following corollary holds:

Corollary 1. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $l, L \in \mathbb{R}$ are such that $L > l$ and $f : [m, M] \rightarrow \mathbb{R}$ is (l, L) -Lipschitzian on $[m, M]$, then we have the inequalities:*

$$(2.15) \quad \begin{aligned} &\left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\ &\leq \frac{1}{4} (M - m)(L - l) \bigvee_m^M (\langle E_{(\cdot)} x, y \rangle) \leq \frac{1}{4} (M - m)(L - l) \|x\| \|y\| \end{aligned}$$

for any $x, y \in H$.

Proof. Follows by applying the inequality (2.10) to the $\frac{1}{2}(L-l)$ -Lipschitzian function $f - \frac{1}{2}(l+L)e$, where $e(t) = t$, $t \in [m, M]$. The details are omitted. \square

When the generating function is continuous convex, we can state the following result as well:

Theorem 7. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$ and let $\{E_\lambda\}_\lambda$ be its spectral family. If $f : [m, M] \rightarrow \mathbb{R}$ is continuous convex on $[m, M]$ with finite lateral derivatives $f'_-(M)$ and $f'_+(m)$, then we have the inequalities:*

$$(2.16) \quad \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\ \leq \frac{1}{4}(M - m) [f'_-(M) - f'_+(m)] \bigvee_m^M (\langle E_{(\cdot)} x, y \rangle) \\ \leq \frac{1}{4}(M - m) [f'_-(M) - f'_+(m)] \|x\| \|y\|$$

for any $x, y \in H$.

Proof. By the convexity of f on $[m, M]$ we have

$$f(t) - f(M) \geq f'_-(M)(t - M)$$

for any $t \in [m, M]$. If we multiply this inequality with $t - m \geq 0$ we deduce

$$(2.17) \quad (t - m)f(t) - (t - m)f(M) \geq f'_-(M)(t - M)(t - m)$$

for any $t \in [m, M]$.

Similarly, we get

$$(2.18) \quad (M - t)f(t) - (M - t)f(m) \geq f'_+(m)(M - t)(t - m)$$

for any $t \in [m, M]$.

Summing the above inequalities and dividing by $M - m$ we deduce the inequality

$$(2.19) \quad \Phi_f(t) \leq \frac{(M - t)(t - m)}{M - m} [f'_-(M) - f'_+(m)] \\ \leq \frac{1}{4}(M - m) [f'_-(M) - f'_+(m)]$$

for any $t \in [m, M]$.

By the convexity of f we also have that

$$(2.20) \quad \frac{1}{M - m} [(M - t)f(m) + (t - m)f(M)] \geq f\left(\frac{(M - t)m + (t - m)M}{M - m}\right) \\ = f(t)$$

giving that

$$(2.21) \quad \Phi_f(t) \geq 0 \text{ for any } t \in [m, M].$$

Utilising (2.3) for the representation (2.2) we deduce from (2.19) and (2.21) the desired result (2.16). \square

3. INEQUALITIES IN THE OPERATOR ORDER

The following result providing some inequalities in the operator order may be stated:

Theorem 8. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$.*

1. *If $f : [m, M] \rightarrow \mathbb{R}$ is continuous on $[m, M]$, then*

$$(3.1) \quad \left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \right| \leq \left[\max_{t \in [m, M]} f(t) - \min_{t \in [m, M]} f(t) \right] 1_H.$$

2. *If $f : [m, M] \rightarrow \mathbb{C}$ is continuous and of bounded variation on $[m, M]$, then*

$$(3.2) \quad \left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \right| \leq \frac{M1_H - A}{M - m} \bigvee_m^A(f) + \frac{A - m1_H}{M - m} \bigvee_A^M(f) \leq \left[\frac{1}{2} + \frac{|A - \frac{m+M}{2}1_H|}{M - m} \right] \bigvee_m^M(f),$$

where $\bigvee_m^A(f)$ denotes the operator generated by the scalar function $[m, M] \ni t \mapsto$

$\bigvee_m^t(f) \in \mathbb{R}$. The same notation applies for $\bigvee_A^M(f)$.

3. *If $f : [m, M] \rightarrow \mathbb{C}$ is Lipschitzian with the constant $L > 0$ on $[m, M]$, then*

$$(3.3) \quad \left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \right| \leq \frac{M1_H - A}{M - m} |f(A) - f(m)1_H| + \frac{A - m1_H}{M - m} |f(M)1_H - f(A)| \leq \frac{1}{2}(M - m)L1_H.$$

4. *If $f : [m, M] \rightarrow \mathbb{R}$ is continuous convex on $[m, M]$ with finite lateral derivatives $f'_-(M)$ and $f'_+(m)$, then we have the inequalities:*

$$(3.4) \quad 0 \leq \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \leq \frac{(M1_H - A)(A - m1_H)}{M - m} [f'_-(M) - f'_+(m)] \leq \frac{1}{4}(M - m) [f'_-(M) - f'_+(m)] 1_H.$$

Proof. Follows by applying the property (P) to the scalar inequalities (2.4), (2.8), (2.11), (2.19) and (2.21). The details are omitted. \square

The following particular case is perhaps more useful for applications:

Corollary 2. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$. If $l, L \in \mathbb{R}$ with $L > l$ and $f : [m, M] \rightarrow \mathbb{R}$ is (l, L) -Lipschitzian on $[m, M]$, then we have the inequalities:*

$$(3.5) \quad \left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} - f(A) \right| \leq \frac{1}{4}(M - m)(L - l)1_H.$$

4. MORE INEQUALITIES FOR DIFFERENTIABLE FUNCTIONS

The following result holds:

Theorem 9. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$. Assume that the function $f : I \rightarrow \mathbb{C}$ with $[m, M] \subset \dot{I}$ (the interior of I) is differentiable on \dot{I} .*

1. *If the derivative f' is continuous and of bounded variation on $[m, M]$, then we have the inequality*

$$(4.1) \quad \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \leq \frac{1}{4}(M - m) \bigvee_m^M(f') \bigvee_m^M(\langle E_{(\cdot)}x, y \rangle) \leq \frac{1}{4}(M - m) \bigvee_m^M(f') \|x\| \|y\|$$

for any $x, y \in H$.

2. *If the derivative f' is Lipschitzian with the constant $K > 0$ on $[m, M]$, then we have the inequality*

$$(4.2) \quad \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \leq \frac{1}{8}(M - m)^2 K \bigvee_m^M(\langle E_{(\cdot)}x, y \rangle) \leq \frac{1}{8}(M - m)^2 K \|x\| \|y\|$$

for any $x, y \in H$.

Proof. First of all we notice that if $f : [m, M] \rightarrow \mathbb{C}$ is absolutely continuous on $[m, M]$ and such that the derivative f' is Riemann integrable on $[m, M]$, then we have the following representation in terms of the Riemann-Stieltjes integral:

$$(4.3) \quad \Phi_f(t) = \frac{1}{M - m} \int_m^M K(t, s) df'(s), \quad t \in [m, M],$$

where the kernel $K : [m, M]^2 \rightarrow \mathbb{R}$ is given by

$$(4.4) \quad K(t, s) := \begin{cases} (M - t)(s - m) & \text{if } m \leq s \leq t \\ (t - m)(M - s) & \text{if } t < s \leq M. \end{cases}$$

Indeed, since f' is Riemann integrable on $[m, M]$, it follows that the Riemann-Stieltjes integrals $\int_m^t (s - m) df'(s)$ and $\int_t^M (M - s) df'(s)$ exist for each $t \in [m, M]$. Now, integrating by parts in the Riemann-Stieltjes integral, we have:

$$\begin{aligned}
\int_m^M K(t, s) df'(s) &= (M - t) \int_m^t (s - m) df'(s) + (t - m) \int_t^M (M - s) df'(s) \\
&= (M - t) \left[(s - m) f'(s) \Big|_m^t - \int_m^t f'(s) ds \right] \\
&\quad + (t - m) \left[(M - s) f'(s) \Big|_t^M - \int_t^M f'(s) ds \right] \\
&= (M - t) [(t - m) f'(t) - (f(t) - f(m))] \\
&\quad + (t - m) [-(M - t) f'(t) + f(M) - f(t)] \\
&= (t - m) [f(M) - f(t)] - (M - t) [f(t) - f(m)] \\
&= (M - m) \Phi_f(t)
\end{aligned}$$

for any $t \in [m, M]$, which provides the desired representation (4.3).

Now, utilizing the representation (4.3) and the property (2.3), we have

$$\begin{aligned}
(4.5) \quad |\Phi_f(t)| &= \frac{1}{M - m} \left| (M - t) \int_m^t (s - m) df'(s) + (t - m) \int_t^M (M - s) df'(s) \right| \\
&\leq \frac{1}{M - m} \left[(M - t) \left| \int_m^t (s - m) df'(s) \right| + (t - m) \left| \int_t^M (M - s) df'(s) \right| \right] \\
&\leq \frac{1}{M - m} \\
&\quad \times \left[(M - t) \bigvee_m^t (f') \sup_{s \in [m, t]} (s - m) + (t - m) \bigvee_t^M (f') \sup_{s \in [t, M]} (M - s) \right] \\
&= \frac{(t - m)(M - t)}{M - m} \left[\bigvee_m^t (f') + \bigvee_t^M (f') \right] \\
&= \frac{(t - m)(M - t)}{M - m} \bigvee_m^M (f') \leq \frac{1}{4} (M - m) \bigvee_m^M (f')
\end{aligned}$$

for any $t \in [m, M]$.

On making use of the representation (2.2) we deduce the desired result (4.1).

Further, we utilize the fact that for an L -Lipschitzian function, $p : [\alpha, \beta] \rightarrow \mathbb{C}$ and a Riemann integrable function $v : [\alpha, \beta] \rightarrow \mathbb{C}$, the Riemann-Stieltjes integral $\int_\alpha^\beta p(s) dv(s)$ exists and

$$\left| \int_\alpha^\beta p(s) dv(s) \right| \leq L \int_\alpha^\beta |p(s)| ds.$$

Then, by utilizing (4.5) we have

$$\begin{aligned}
(4.6) \quad |\Phi_f(t)| &\leq \frac{1}{M-m} \left[(M-t) \left| \int_m^t (s-m) df'(s) \right| + (t-m) \left| \int_t^M (M-s) df'(s) \right| \right] \\
&\leq \frac{K}{M-m} \left[(M-t) \int_m^t (s-m) ds + (t-m) \int_t^M (M-s) ds \right] \\
&= \frac{K}{M-m} \left[\frac{(M-t)(t-m)^2}{2} + \frac{(t-m)(M-t)^2}{2} \right] \\
&= \frac{1}{2} (M-m)(t-m)(M-t) K \leq \frac{1}{8} (M-m)^2 K
\end{aligned}$$

for any $t \in [m, M]$.

On making use of the representation (2.2) we deduce the desired result (4.2). \square

The following inequalities in the operator order are of interest as well:

Theorem 10. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $m < M$. Assume that the function $f : I \rightarrow \mathbb{C}$ with $[m, M] \subset \dot{I}$ (the interior of I) is differentiable on \dot{I} .*

1. *If the derivative f' is continuous and of bounded variation on $[m, M]$, then we have the inequality*

$$\begin{aligned}
(4.7) \quad &\left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M-m} - f(A) \right| \\
&\leq \frac{(A - m1_H)(M1_H - A)}{M-m} \bigvee_m^M(f') \\
&\leq \frac{1}{4} (M-m) \bigvee_m^M(f') 1_H.
\end{aligned}$$

2. *If the derivative f' is Lipschitzian with the constant $K > 0$ on $[m, M]$, then we have the inequality*

$$\begin{aligned}
(4.8) \quad &\left| \frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M-m} - f(A) \right| \\
&\leq \frac{1}{2} (M-m)(A - m1_H)(M1_H - A) K \\
&\leq \frac{1}{8} (M-m)^2 K 1_H.
\end{aligned}$$

The proof follows by the property (P) applied for the scalar inequalities (4.5) and (4.6).

5. APPLICATIONS FOR PARTICULAR FUNCTIONS

It is obvious that the above results can be applied for various particular functions. However, we will restrict here only to the power and logarithmic functions.

1. Consider now the power function $f : (0, \infty) \rightarrow \mathbb{R}$, $f(t) = t^p$ with $p \neq 0$. On applying Theorem 7 we can state the following proposition:

Proposition 2. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $0 < m < M$. Then for any $x, y \in H$ we have the inequalities*

$$(5.1) \quad \left| \left\langle \left[\frac{m^p (M1_H - A) + M^p (A - m1_H)}{M - m} \right] x, y \right\rangle - \langle A^p x, y \rangle \right| \leq \frac{1}{2} (M - m) \Delta_p \|x\| \|y\|$$

where

$$\Delta_p = p \times \begin{cases} M^{p-1} - m^{p-1} & \text{if } p \in (-\infty, 0) \cup [1, \infty) \\ m^{p-1} - M^{p-1} & \text{if } 0 < p < 1. \end{cases}$$

In particular,

$$(5.2) \quad \left| \left\langle \left[\frac{M (M1_H - A) + m (A - m1_H)}{mM (M - m)} \right] x, y \right\rangle - \langle A^{-1} x, y \rangle \right| \leq \frac{1}{2} \frac{(M - m)^2 (M + m)}{m^2 M^2} \|x\| \|y\|$$

for any $x, y \in H$.

The following inequalities in the operator order also hold:

Proposition 3. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $0 < m < M$.*

If $p \in (-\infty, 0) \cup [1, \infty)$, then

$$(5.3) \quad \begin{aligned} 0 &\leq \frac{m^p (M1_H - A) + M^p (A - m1_H)}{M - m} - A^p \\ &\leq p \frac{(M1_H - A) (A - m1_H)}{M - m} (M^{p-1} - m^{p-1}) \\ &\leq \frac{1}{4} p (M - m) (M^{p-1} - m^{p-1}) 1_H. \end{aligned}$$

If $p \in (0, 1)$, then

$$(5.4) \quad \begin{aligned} 0 &\leq A^p - \frac{m^p (M1_H - A) + M^p (A - m1_H)}{M - m} \\ &\leq p \frac{(M1_H - A) (A - m1_H)}{M - m} (m^{p-1} - M^{p-1}) \\ &\leq \frac{1}{4} p (M - m) (m^{p-1} - M^{p-1}) 1_H. \end{aligned}$$

In particular, we have the inequalities

$$(5.5) \quad \begin{aligned} 0 &\leq \frac{M (M1_H - A) + m (A - m1_H)}{mM (M - m)} - A^{-1} \\ &\leq \frac{(M1_H - A) (A - m1_H)}{M - m} \cdot \frac{M^2 - m^2}{m^2 M^2} \\ &\leq \frac{1}{2} \frac{(M - m)^2 (M + m)}{m^2 M^2} 1_H. \end{aligned}$$

The proof follows from (3.4) and the details are omitted.

2. The case of logarithmic function is as follows:

Proposition 4. *Let A be a selfadjoint operator in the Hilbert space H with the spectrum $Sp(A) \subseteq [m, M]$ for some real numbers $0 < m < M$. Then for any $x, y \in H$ we have the inequalities*

$$(5.6) \quad \left| \left\langle \left[\frac{(M1_H - A) \ln m + (A - m1_H) \ln M}{M - m} \right] x, y \right\rangle - \langle \ln Ax, y \rangle \right| \leq \frac{1}{4} \frac{(M - m)^2}{mM} \|x\| \|y\|.$$

We also have the following inequality in the operator order

$$(5.7) \quad 0 \leq \ln A - \frac{(M1_H - A) \ln m + (A - m1_H) \ln M}{M - m} \leq \frac{(M1_H - A)(A - m1_H)}{Mm} \leq \frac{1}{4} \frac{(M - m)^2}{mM} 1_H.$$

Remark 1. *Similar results can be obtained if ones uses the inequalities from Theorem 9 and 10. However the details are left to the interested reader.*

REFERENCES

- [1] P. Cerone and S.S. Dragomir, Trapezoidal-type rules from an inequalities point of view, in *Handbook of Analytic-Computational Methods in Applied Mathematics*, G.A. Anastassiou (Ed), Chapman & Hall/CRC Press, New York, 2000, 65-134.
- [2] P. Cerone, S.S. Dragomir and C.E.M. Pearce, A generalised trapezoid inequality for functions of bounded variation, *Turkish J. of Math.*, **24**(2) (2000), 147-163.
- [3] S.S. Dragomir, On the trapezoid quadrature formula for Lipschitzian mappings and applications, *Tamkang J. of Math.*, **30**(2) (1999), 133-138.
- [4] S.S. Dragomir, An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products, *J Inequal. Pure & Appl. Math.*, **3**(2002), No. 3, Article 2002.
- [5] S.S. Dragomir, Čebyšev's type inequalities for functions of selfadjoint operators in Hilbert spaces, Preprint *RGMI Res. Rep. Coll.*, **11**(e) (2008), Art. 9. [ONLINE: [http://www.staff.vu.edu.au/RGMIA/v11\(E\).asp](http://www.staff.vu.edu.au/RGMIA/v11(E).asp)].
- [6] S.S. Dragomir, Grüss' type inequalities for functions of selfadjoint operators in Hilbert spaces, Preprint *RGMI Res. Rep. Coll.*, **11**(e) (2008), Art. 11. [ONLINE: [http://www.staff.vu.edu.au/RGMIA/v11\(E\).asp](http://www.staff.vu.edu.au/RGMIA/v11(E).asp)].
- [7] S.S. Dragomir, Inequalities for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces, Preprint *RGMI Res. Rep. Coll.*, **11**(e) (2008), Art. . [ONLINE: [http://www.staff.vu.edu.au/RGMIA/v11\(E\).asp](http://www.staff.vu.edu.au/RGMIA/v11(E).asp)].
- [8] S.S. Dragomir, Some trapezoidal vector inequalities for continuous functions of selfadjoint operators in Hilbert spaces, Preprint *RGMI Res. Rep. Coll.* **13** (2010), to appear.
- [9] S.S. Dragomir, Some generalized trapezoidal vector inequalities for continuous functions of selfadjoint operators in Hilbert spaces, Preprint *RGMI Res. Rep. Coll.* **13** (2010), to appear.
- [10] T. Furuta, J. Mičić Hot, J. Pečarić and Y. Seo, *Mond-Pečarić Method in Operator Inequalities. Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Element, Zagreb, 2005.
- [11] Z. Liu, Refinement of an inequality of Grüss type for Riemann-Stieltjes integral, *Soochow J. Math.*, **30**(4) (2004), 483-489.
- [12] A. Matković, J. Pečarić and I. Perić, A variant of Jensen's inequality of Mercer's type for operators with applications. *Linear Algebra Appl.* **418** (2006), No. 2-3, 551-564.
- [13] B. Mond and J. Pečarić, Convex inequalities in Hilbert spaces, *Houston J. Math.*, **19**(1993), 405-420.
- [14] B. Mond and J. Pečarić, Classical inequalities for matrix functions, *Utilitas Math.*, **46**(1994), 155-166.
- [15] J. Pečarić, J. Mičić and Y. Seo, Inequalities between operator means based on the Mond-Pečarić method. *Houston J. Math.* **30** (2004), no. 1, 191-207.

MATHEMATICS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428,
MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: `sever.dragomir@vu.edu.au`

URL: `http://www.staff.vu.edu.au/rgmia/dragomir/`