

Received 29/12/11

**NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR
FUNCTIONS WHOSE DERIVATIVES IN ABSOLUTE VALUE
ARE CONVEX WITH APPLICATIONS TO SPECIAL MEANS
AND TO GENERAL QUADRATURE FORMULA**

M. A. LATIF AND S. S. DRAGOMIR

ABSTRACT. In this paper some new Hadamard-type inequalities for functions whose derivatives in absolute values are convex are established. We also give some applications of our obtained results to get some error bounds for the general quadrature formula. Finally, Some applications to special means of real numbers are given as well.

1. INTRODUCTION

The following definition for convex functions is well known in the mathematical literature:

A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y),$$

holds for all $x, y \in I$ and $t \in [0, 1]$.

Many inequalities have been established for convex functions but the most famous is the Hermite-Hadamard's inequality, due to its rich geometrical significance and applications, which is stated as follow:

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping and $a, b \in I$ with $a < b$. Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}. \quad (1.1)$$

Both the inequalities hold in reversed direction if f is concave. Since its discovery in 1883, Hermite-Hadamard's inequality [4] has been considered the most useful inequality in mathematical analysis. Some of the classical inequalities for means can be derived from (1.1) for particular choices of the function f . A number of papers have been written on this inequality providing new proofs, noteworthy extensions, generalizations and numerous applications, see [1]-[14] and the references therein.

Date: November 18, 2011.

2000 Mathematics Subject Classification. Primary 26A51; 26D15.

Key words and phrases. Hermite-Hadamard's inequality, convex functions, Hölder inequality, power-mean inequality, special means.

This paper is in final form and no version of it will be submitted for publication elsewhere.

In a recent paper [12], K. L. Tseng et al., established the following result which gives a refinement of (1.1):

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} \leq \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \frac{1}{2} \left[f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right] \leq \frac{f(a) + f(b)}{2}, \end{aligned} \quad (1.2)$$

where $f : [a, b] \rightarrow \mathbb{R}$, is a convex function (see [12, Remark 2.11, page7.]).

The main aim of this paper is to establish some new Hermite-Hadamard type inequalities which give an estimate between $\frac{1}{b-a} \int_a^b f(x) dx$ and $\frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2}$ for functions whose derivatives in absolute value are convex and as a consequence we will get refinements of those results which have been established to estimate the difference between the middle and the leftmost terms in (1.1).

2. MAIN RESULTS

To prove our results we need the following lemma:

Lemma 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° , the interior of I , where $a, b \in I$ with $a < b$. If $f' \in L[a, b]$, then the following equality holds:*

$$\begin{aligned} &\frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \\ &= \frac{b-a}{16} \left[\int_0^1 t f' \left(t \frac{3a+b}{4} + (1-t)a \right) dt \right. \\ &\quad + \int_0^1 (t-1) f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) dt \\ &\quad + \int_0^1 t f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) dt \\ &\quad \left. + \int_0^1 (t-1) f' \left(tb + (1-t) \frac{a+3b}{4} \right) dt \right]. \end{aligned} \quad (2.1)$$

Proof. By integration by parts and by making use of the substitution $u = t \frac{3a+b}{4} + (1-t)a$, we have

$$\begin{aligned} &\frac{b-a}{16} \int_0^1 t f' \left(t \frac{3a+b}{4} + (1-t)a \right) dt \\ &= \frac{b-a}{16} \left[\frac{4t f' \left(t \frac{3a+b}{4} + (1-t)a \right)}{b-a} \Big|_0^1 - \frac{4}{b-a} \int_0^1 f \left(t \frac{3a+b}{4} + (1-t)a \right) dt \right] \\ &= \frac{1}{4} f \left(\frac{3a+b}{4} \right) - \frac{1}{b-a} \int_a^{\frac{3a+b}{4}} f(u) du. \end{aligned} \quad (2.2)$$

Analogously, we also have the following equalities:

$$\begin{aligned} \frac{b-a}{16} \int_0^1 (t-1) f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) dt \\ = \frac{1}{4} f \left(\frac{3a+b}{4} \right) - \frac{1}{b-a} \int_{\frac{3a+b}{4}}^{\frac{a+b}{2}} f(u) du, \end{aligned} \quad (2.3)$$

$$\begin{aligned} \frac{b-a}{16} \int_0^1 t f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) dt \\ = \frac{1}{4} f \left(\frac{a+3b}{4} \right) - \frac{1}{b-a} \int_{\frac{a+b}{2}}^{\frac{a+3b}{4}} f(u) du \end{aligned} \quad (2.4)$$

and

$$\begin{aligned} \frac{b-a}{16} \int_0^1 (t-1) f' \left(tb + (1-t) \frac{a+3b}{4} \right) dt \\ = \frac{1}{4} f \left(\frac{a+3b}{4} \right) - \frac{1}{b-a} \int_{\frac{a+3b}{4}}^b f(u) du. \end{aligned} \quad (2.5)$$

Adding (2.2)-(2.5), we get the desired equality. This completes the proof of the lemma. \square

Using the Lemma 1 the following results can be obtained:

Theorem 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then the following inequality holds:*

$$\begin{aligned} \left| \frac{f \left(\frac{3a+b}{4} \right) + f \left(\frac{a+3b}{4} \right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{b-a}{96} \right) \left[|f'(a)| + 4 \left| f' \left(\frac{3a+b}{4} \right) \right| \right. \\ \left. + 2 \left| f' \left(\frac{a+b}{2} \right) \right| + 4 \left| f' \left(\frac{a+3b}{4} \right) \right| + |f'(b)| \right]. \end{aligned} \quad (2.6)$$

Proof. Using Lemma 1 and taking the modulus, we have

$$\begin{aligned} \left| \frac{f \left(\frac{3a+b}{4} \right) + f \left(\frac{a+3b}{4} \right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ \leq \frac{b-a}{16} \left[\int_0^1 t \left| f' \left(t \frac{3a+b}{4} + (1-t) a \right) \right| dt \right. \\ \left. + \int_0^1 (1-t) \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right| dt \right. \\ \left. + \int_0^1 t \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right| dt \right. \\ \left. + \int_0^1 (1-t) \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right| dt \right]. \end{aligned} \quad (2.7)$$

Using the convexity of $|f'|$ on $[a, b]$, we observe that the following inequality holds:

$$\begin{aligned} \int_0^1 t \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right| dt \\ \leq \left| f' \left(\frac{3a+b}{4} \right) \right| \int_0^1 t^2 dt + |f'(a)| \int_0^1 t(1-t) dt \\ = \frac{1}{3} \left| f' \left(\frac{3a+b}{4} \right) \right| + \frac{1}{6} |f'(a)|. \end{aligned} \quad (2.8)$$

Similarly, we also have that the following inequalities hold:

$$\int_0^1 (1-t) \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right| dt \leq \frac{1}{6} \left| f' \left(\frac{a+b}{2} \right) \right| + \frac{1}{3} \left| f' \left(\frac{3a+b}{4} \right) \right|, \quad (2.9)$$

$$\int_0^1 t \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right| dt \leq \frac{1}{3} \left| f' \left(\frac{a+3b}{4} \right) \right| + \frac{1}{6} \left| f' \left(\frac{a+b}{2} \right) \right|, \quad (2.10)$$

and

$$\int_0^1 (1-t) \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right| dt \leq \frac{1}{6} |f'(b)| + \frac{1}{3} \left| f' \left(\frac{a+3b}{4} \right) \right|. \quad (2.11)$$

Utilizing the inequalities (2.8)-(2.11), we get (2.6).

This completes the proof of the theorem. \square

Corollary 1. *Suppose all the conditions of Theorem 1 are satisfied. Then*

$$\left| \frac{f \left(\frac{3a+b}{4} \right) + f \left(\frac{a+3b}{4} \right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{b-a}{16} \right) [|f'(a)| + |f'(b)|]. \quad (2.12)$$

Moreover, if $|f'(x)| \leq M$, for all $x \in [a, b]$, then we have also the following inequality:

$$\left| \frac{f \left(\frac{3a+b}{4} \right) + f \left(\frac{a+3b}{4} \right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{b-a}{8} \right) M. \quad (2.13)$$

Proof. It follows from Theorem 1 and using the convexity of $|f'|$. \square

Theorem 2. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is convex on $[a, b]$ for some fixed $q > 1$, then the following inequality holds:*

$$\begin{aligned} \left| \frac{f \left(\frac{3a+b}{4} \right) + f \left(\frac{a+3b}{4} \right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| &\leq \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{1}{q}} \left(\frac{b-a}{16} \right) \\ &\times \left\{ \left(\left| f' \left(\frac{3a+b}{4} \right) \right|^q + |f'(a)|^q \right)^{\frac{1}{q}} + \left(\left| f' \left(\frac{a+b}{2} \right) \right|^q + \left| f' \left(\frac{3a+b}{4} \right) \right|^q \right)^{\frac{1}{q}} \right. \\ &\left. + \left(\left| f' \left(\frac{a+3b}{4} \right) \right|^q + \left| f' \left(\frac{a+b}{2} \right) \right|^q \right)^{\frac{1}{q}} + \left(\left| f' \left(\frac{a+3b}{4} \right) \right|^q + |f'(b)|^q \right)^{\frac{1}{q}} \right\}, \end{aligned} \quad (2.14)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and using the well-known Hölder integral inequality, we have

$$\begin{aligned}
& \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\
& \leq \frac{b-a}{16} \left[\left(\int_0^1 t^p dt \right)^{\frac{1}{p}} \left(\int_0^1 \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right|^q dt \right)^{\frac{1}{q}} \right. \\
& \quad + \left(\int_0^1 (1-t)^p dt \right)^{\frac{1}{p}} \left(\int_0^1 \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right|^q dt \right)^{\frac{1}{q}} \\
& \quad + \left(\int_0^1 t^p dt \right)^{\frac{1}{p}} \left(\int_0^1 \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right|^q dt \right)^{\frac{1}{q}} \\
& \quad \left. + \left(\int_0^1 (1-t)^p dt \right)^{\frac{1}{p}} \left(\int_0^1 \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right|^q dt \right)^{\frac{1}{q}} \right]. \quad (2.15)
\end{aligned}$$

Since $|f'|^q$ is convex on $[a, b]$, we have

$$\begin{aligned}
& \int_0^1 \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right|^q dt \\
& \leq \left| f' \left(\frac{3a+b}{4} \right) \right|^q \int_0^1 t dt + |f'(a)|^q \int_0^1 (1-t) dt \\
& = \frac{1}{2} \left| f' \left(\frac{3a+b}{4} \right) \right|^q + \frac{1}{2} |f'(a)|^q.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \int_0^1 \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right|^q dt \leq \frac{1}{2} \left| f' \left(\frac{a+b}{2} \right) \right|^q + \frac{1}{2} \left| f' \left(\frac{3a+b}{4} \right) \right|^q, \\
& \int_0^1 \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right|^q dt \leq \frac{1}{2} \left| f' \left(\frac{a+3b}{4} \right) \right|^q + \frac{1}{2} \left| f' \left(\frac{a+b}{2} \right) \right|^q
\end{aligned}$$

and

$$\int_0^1 \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right|^q dt \leq \frac{1}{2} \left| f' \left(\frac{a+3b}{4} \right) \right|^q + \frac{1}{2} |f'(b)|^q.$$

Using the last four inequalities in (2.15), we get the inequality (2.14), which completes the proof of the theorem. \square

Corollary 2. *Suppose all the conditions of Theorem 2 are satisfied. Then*

$$\begin{aligned}
& \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\
& \leq \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{3}{q}} \left[1 + 3^{\frac{1}{q}} + 5^{\frac{1}{q}} + 7^{\frac{1}{q}} \right] \left(\frac{b-a}{16} \right) [|f'(a)| + |f'(b)|]. \quad (2.16)
\end{aligned}$$

Proof. It follows from Theorem 2 using the convexity of $|f'|^q$ and the fact

$$\sum_{k=1}^n (u_k + v_k)^s \leq \sum_{k=1}^n (u_k)^s + \sum_{k=1}^n (v_k)^s, \quad u_k, v_k \geq 0, 1 \leq k \leq n, 0 \leq s < 1.$$

\square

Theorem 3. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is convex on $[a, b]$ for some fixed $q \geq 1$, then the following inequality holds:

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{1}{2}\right) \left(\frac{1}{3}\right)^{\frac{1}{q}} \left(\frac{b-a}{16}\right) \\ & \times \left\{ \left(|f'(a)|^q + 2 \left| f'\left(\frac{3a+b}{4}\right) \right|^q \right)^{\frac{1}{q}} + \left(\left| f'\left(\frac{a+b}{2}\right) \right|^q + 2 \left| f'\left(\frac{3a+b}{4}\right) \right|^q \right)^{\frac{1}{q}} \right. \\ & \left. + \left(\left| f'\left(\frac{a+b}{2}\right) \right|^q + 2 \left| f'\left(\frac{a+3b}{4}\right) \right|^q \right)^{\frac{1}{q}} + \left(2 \left| f'\left(\frac{a+3b}{4}\right) \right|^q + |f'(b)|^q \right)^{\frac{1}{q}} \right\}. \end{aligned} \quad (2.17)$$

Proof. Suppose that $q \geq 1$. From Lemma 1 and using the well-known power-mean inequality, we have

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \frac{b-a}{16} \left[\left(\int_0^1 t dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t \left| f'\left(t\frac{3a+b}{4} + (1-t)a\right) \right|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad + \left(\int_0^1 (1-t) dt \right)^{1-\frac{1}{q}} \left(\int_0^1 (1-t) \left| f'\left(t\frac{a+b}{2} + (1-t)\frac{3a+b}{4}\right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad + \left(\int_0^1 t dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t \left| f'\left(t\frac{a+3b}{4} + (1-t)\frac{a+b}{2}\right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad \left. + \left(\int_0^1 (1-t) dt \right)^{1-\frac{1}{q}} \left(\int_0^1 (1-t) \left| f'\left(tb + (1-t)\frac{a+3b}{4}\right) \right|^q dt \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (2.18)$$

Since $|f'|^q$ is convex on $[a, b]$, we have

$$\begin{aligned} & \int_0^1 t \left| f'\left(t\frac{3a+b}{4} + (1-t)a\right) \right|^q dt \\ & \leq \left| f'\left(\frac{3a+b}{4}\right) \right|^q \int_0^1 t^2 dt + |f'(a)|^q \int_0^1 t(1-t) dt \\ & = \frac{1}{3} \left| f'\left(\frac{3a+b}{4}\right) \right|^q + \frac{1}{6} |f'(a)|^q. \end{aligned}$$

Analogously, we also have that the following inequalities:

$$\begin{aligned} & \int_0^1 (1-t) \left| f'\left(t\frac{a+b}{2} + (1-t)\frac{3a+b}{4}\right) \right|^q dt \\ & \leq \frac{1}{6} \left| f'\left(\frac{a+b}{2}\right) \right|^q + \frac{1}{3} \left| f'\left(\frac{3a+b}{4}\right) \right|^q, \\ & \int_0^1 t \left| f'\left(t\frac{a+3b}{4} + (1-t)\frac{a+b}{2}\right) \right|^q dt \leq \frac{1}{3} \left| f'\left(\frac{a+3b}{4}\right) \right|^q + \frac{1}{6} \left| f'\left(\frac{a+b}{2}\right) \right|^q \end{aligned}$$

and

$$\int_0^1 (1-t) \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right|^q \leq \frac{1}{3} \left| f' \left(\frac{a+3b}{4} \right) \right|^q + \frac{1}{6} |f'(b)|^q.$$

By making use of the last four inequalities in (2.18), we get (2.17). Hence the proof of the theorem is complete. \square

Corollary 3. *Suppose all the conditions of Theorem 2 are satisfied. Then using similar arguments as in Corollary 2, we get the following inequality:*

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \left(\frac{1}{3}\right)^{\frac{1}{q}} \left(\frac{1}{2}\right)^{\frac{1}{q}} \left[1 + 2^{\frac{1}{q}} + \left(\frac{1}{2}\right)^{\frac{1}{q}} + \left(\frac{5}{2}\right)^{\frac{1}{q}} \right] \left(\frac{b-a}{16}\right) [|f'(a)| + |f'(b)|]. \end{aligned} \quad (2.19)$$

Theorem 4. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is concave on $[a, b]$ for some fixed $q > 1$, then the following inequality holds:*

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{b-a}{16}\right) \left\{ \left| f' \left(\frac{7a+b}{8} \right) \right| + \left| f' \left(\frac{5a+3b}{8} \right) \right| \right. \\ & \quad \left. + \left| f' \left(\frac{3a+5b}{8} \right) \right| + \left| f' \left(\frac{a+7b}{8} \right) \right| \right\}, \end{aligned} \quad (2.20)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and using the well-known Hölder integral inequality for $q > 1$ and $p = \frac{q}{q-1}$, we have

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \left(\frac{b-a}{16}\right) \left[\left(\int_0^1 t^{\frac{q}{q-1}} dt \right)^{\frac{q-1}{q}} \left(\int_0^1 \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad + \left(\int_0^1 (1-t)^{\frac{q}{q-1}} dt \right)^{\frac{q-1}{q}} \left(\int_0^1 \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad + \left(\int_0^1 t^{\frac{q}{q-1}} dt \right)^{\frac{q-1}{q}} \left(\int_0^1 \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad \left. + \left(\int_0^1 (1-t)^{\frac{q}{q-1}} dt \right)^{\frac{q-1}{q}} \left(\int_0^1 \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right|^q dt \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (2.21)$$

Since $|f'|^q$ is concave on $[a, b]$ so by using the inequality (1.1), we obtain:

$$\int_0^1 \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right|^q dt \leq \left| f' \left(\frac{\frac{3a+b}{4} + a}{2} \right) \right|^q = \left| f' \left(\frac{7a+b}{8} \right) \right|^q$$

Analogously, we have that the following inequalities:

$$\int_0^1 \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right|^q dt \leq \left| f' \left(\frac{5a+3b}{8} \right) \right|^q,$$

$$\int_0^1 \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right|^q dt \leq \left| f' \left(\frac{3a+5b}{8} \right) \right|^q$$

and

$$\int_0^1 \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right|^q dt \leq \left| f' \left(\frac{a+7b}{8} \right) \right|^q.$$

Using the last four inequalities in (2.21), we get (2.20). This completes the proof of the theorem. \square

Corollary 4. *Suppose all the assumptions of Theorem 4 are satisfied and assume that $|f'|$ is a linear map, then we get the following inequality:*

$$\left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{q-1}{2q-1} \right)^{\frac{q-1}{q}} \left(\frac{b-a}{8} \right) |f'(a+b)|. \quad (2.22)$$

Proof. It is a direct consequence of Theorem 4 and using the linearity of $|f'|$. \square

Theorem 5. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is concave on $[a, b]$ for some fixed $q \geq 1$, then the following inequality holds:*

$$\begin{aligned} & \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \left(\frac{b-a}{32} \right) \left[\left| f' \left(\frac{13a+3b}{12} \right) \right| + \left| f' \left(\frac{11a+5b}{12} \right) \right| \right. \\ & \quad \left. + \left| f' \left(\frac{5a+13b}{12} \right) \right| + \left| f' \left(\frac{3a+13b}{12} \right) \right| \right]. \quad (2.23) \end{aligned}$$

Proof. First, by the concavity of $|f'|^q$ on $[a, b]$ and the power-mean inequality, we note that

$$\begin{aligned} |f(\lambda x + (1-\lambda)y)|^q & \geq \lambda |f(x)|^q + (1-\lambda) |f(y)|^q \\ & \geq (\lambda |f(x)| + (1-\lambda) |f(y)|)^q \end{aligned}$$

and hence

$$|f(\lambda x + (1-\lambda)y)| \geq \lambda |f(x)| + (1-\lambda) |f(y)|,$$

for all $\lambda \in [0, 1]$ and $x, y \in [a, b]$. This shows that $|f'|$ is also concave on $[a, b]$. Accordingly, using Lemma 1 and the Jensen's integral inequality, we have

$$\begin{aligned}
& \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\
& \leq \left(\frac{b-a}{16} \right) \left[\int_0^1 t \left| f' \left(t \frac{3a+b}{4} + (1-t)a \right) \right| dt \right. \\
& \quad + \int_0^1 (1-t) \left| f' \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) \right| dt \\
& \quad + \int_0^1 t \left| f' \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) \right| dt \\
& \quad \left. + \int_0^1 (1-t) \left| f' \left(tb + (1-t) \frac{a+3b}{4} \right) \right| dt \right] \\
& \leq \frac{b-a}{16} \left[\left(\int_0^1 t dt \right) \left| f' \left(\frac{\int_0^1 t \left(t \frac{3a+b}{4} + (1-t)a \right) dt}{\int_0^1 t dt} \right) \right| \right] \\
& \quad + \left(\int_0^1 (1-t) dt \right) \left| f' \left(\frac{\int_0^1 (1-t) \left(t \frac{a+b}{2} + (1-t) \frac{3a+b}{4} \right) dt}{\int_0^1 (1-t) dt} \right) \right| \\
& \quad + \left(\int_0^1 t dt \right) \left| f' \left(\frac{\int_0^1 t \left(t \frac{a+3b}{4} + (1-t) \frac{a+b}{2} \right) dt}{\int_0^1 t dt} \right) \right| \\
& \quad + \left(\int_0^1 (1-t) dt \right) \left| f' \left(\frac{\int_0^1 (1-t) \left(tb + (1-t) \frac{a+3b}{4} \right) dt}{\int_0^1 (1-t) dt} \right) \right|,
\end{aligned}$$

which is equivalent to (2.23) and the proof of the theorem is complete. \square

Corollary 5. *Suppose all the assumptions of Theorem 5 are satisfied and assume that $|f'|$ is a linear map, then we have the following inequality:*

$$\left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \left(\frac{b-a}{12} \right) |f'(a+b)|. \quad (2.24)$$

Proof. It follows from Theorem 5 and using the linearity of $|f'|$. \square

3. APPLICATION TO THE GENERAL QUADRATURE FORMULA

Let $d : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be a division of the interval $[a, b]$. Consider the general quadrature formula

$$\int_a^b f(x) dx = Q(f, d) + R(f, d), \quad (3.1)$$

where

$$Q(f, d) = \frac{1}{2} \sum_{i=0}^{n-1} \left[f \left(\frac{3x_i + x_{i+1}}{4} \right) + f \left(\frac{x_i + 3x_{i+1}}{4} \right) \right] (x_{i+1} - x_i)$$

and $R(f, d)$ is the associated error. Here, we derive some estimates for the error $R(f, d)$ given in (3.1).

Proposition 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then for every division d of $[a, b]$, we have:*

$$|R(f, d)| \leq \frac{1}{96} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left[|f'(x_i)| + 4 \left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right| \right. \\ \left. + 2 \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right| + 4 \left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right| + |f'(x_{i+1})| \right]. \quad (3.2)$$

Proof. By applying Theorem 1 on the subinterval $[x_i, x_{i+1}]$ ($i = 0, 1, \dots, n-1$) of the division d , we have

$$\left| \frac{1}{2} \left[f \left(\frac{3x_i + x_{i+1}}{4} \right) + f \left(\frac{x_i + 3x_{i+1}}{4} \right) \right] \right. \\ \left. - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right| \\ \leq \left(\frac{x_{i+1} - x_i}{96} \right) \left[|f'(x_i)| + 4 \left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right| \right. \\ \left. + 2 \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right| + 4 \left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right| + |f'(x_{i+1})| \right]. \quad (3.3)$$

Now

$$|R(f, d)| = \left| \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} f(x) dx \right. \\ \left. - \sum_{i=0}^{n-1} \frac{1}{2} \left[f \left(\frac{3x_i + x_{i+1}}{4} \right) + f \left(\frac{x_i + 3x_{i+1}}{4} \right) \right] (x_{i+1} - x_i) \right| \\ \leq \sum_{i=0}^{n-1} (x_{i+1} - x_i) \left| \frac{1}{2} \left[f \left(\frac{3x_i + x_{i+1}}{4} \right) + f \left(\frac{x_i + 3x_{i+1}}{4} \right) \right] \right. \\ \left. - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right|. \quad (3.4)$$

Using (3.3) in (3.4), we get (3.2). This completes the proof of the proposition. \square

Corollary 6. *Suppose all the assumptions of Proposition 1 are satisfied. Then*

$$|R(f, d)| \leq \frac{1}{16} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 [|f'(x_i)| + |f'(x_{i+1})|]. \quad (3.5)$$

Proof. It follows from Proposition 1 and using the convexity of $|f'|$. \square

Proposition 2. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is convex on $[a, b]$ for some fixed*

$q > 1$, then for every division d of $[a, b]$, we have

$$\begin{aligned}
& |R(f, d)| \\
& \leq \left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left(\frac{1}{2}\right)^{\frac{1}{q}+4} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left\{ \left(\left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right|^q + |f'(x_i)|^q \right)^{\frac{1}{q}} \right. \\
& \quad + \left(\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + \left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right|^q \right)^{\frac{1}{q}} \\
& \quad + \left(\left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right|^q + \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q \right)^{\frac{1}{q}} \\
& \quad \left. + \left(\left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right|^q + |f'(x_{i+1})|^q \right)^{\frac{1}{q}} \right\}, \quad (3.6)
\end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. The proof is similar to that of Proposition 1 and using Theorem 2. \square

Corollary 7. *Suppose all the conditions of Proposition 2 are satisfied. Then*

$$\begin{aligned}
|R(f, d)| & \leq \left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left(\frac{1}{2}\right)^{\frac{3}{q}+4} \left[1 + 3^{\frac{1}{q}} + 5^{\frac{1}{q}} + 7^{\frac{1}{q}}\right] \\
& \quad \times \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 [|f'(x_i)| + |f'(x_{i+1})|]. \quad (3.7)
\end{aligned}$$

Proposition 3. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is convex on $[a, b]$ for some fixed $q \geq 1$, then for every division d of $[a, b]$, we have*

$$\begin{aligned}
|R(f, d)| & \leq \left(\frac{1}{32}\right) \left(\frac{1}{3}\right)^{\frac{1}{q}} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left\{ \left(|f'(x_i)|^q + 2 \left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right|^q \right)^{\frac{1}{q}} \right. \\
& \quad + \left(\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + 2 \left| f' \left(\frac{3x_i + x_{i+1}}{4} \right) \right|^q \right)^{\frac{1}{q}} \\
& \quad + \left(\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + 2 \left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right|^q \right)^{\frac{1}{q}} \\
& \quad \left. + \left(2 \left| f' \left(\frac{x_i + 3x_{i+1}}{4} \right) \right|^q + |f'(x_{i+1})|^q \right)^{\frac{1}{q}} \right\}. \quad (3.8)
\end{aligned}$$

Proof. The proof is similar to that of Proposition 1 and using Theorem 3. \square

Corollary 8. *Suppose all the conditions of Proposition 3 are satisfied. Then*

$$\begin{aligned}
|R(f, d)| & \leq \left(\frac{1}{3}\right)^{\frac{1}{q}} \left(\frac{1}{32}\right) \left[1 + 2^{\frac{1}{q}} + \left(\frac{1}{2}\right)^{\frac{1}{q}} + \left(\frac{5}{2}\right)^{\frac{1}{q}}\right] \\
& \quad \times \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 [|f'(x_i)| + |f'(x_{i+1})|]. \quad (3.9)
\end{aligned}$$

Proposition 4. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is concave on $[a, b]$ for some fixed $q > 1$, then for every division d of $[a, b]$, we have

$$|R(f, d)| \leq \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{16}\right) \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 \left\{ \left| f' \left(\frac{7x_i + x_{i+1}}{8} \right) \right| \right. \\ \left. + \left| f' \left(\frac{5x_i + 3x_{i+1}}{8} \right) \right| + \left| f' \left(\frac{3x_i + 5x_{i+1}}{8} \right) \right| + \left| f' \left(\frac{x_i + 7x_{i+1}}{8} \right) \right| \right\}, \quad (3.10)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. The proof is similar to that of Proposition 1 and it follows from Theorem 4. \square

Corollary 9. Suppose all the conditions of Proposition 4 are satisfied. If $|f'|$ is a linear mapping, then we have the following inequality:

$$|R(f, d)| \leq \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{8}\right) \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 |f'(x_{i+1} + x_i)|. \quad (3.11)$$

Proposition 5. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is concave on $[a, b]$ for some fixed $q \geq 1$ and $|f'|^q$ is a linear mapping, then for every division d of $[a, b]$, then the following inequality holds:

$$|R(f, d)| \leq \left(\frac{1}{32}\right) \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 |f'(x_{i+1} + x_i)|. \quad (3.12)$$

Proof. The proof is similar to that of Proposition 1 and it follows from Theorem 5. \square

4. APPLICATIONS TO SPECIAL MEANS

Now, we consider the applications of our Theorems to the special means. We consider the means for arbitrary real numbers $a, b \in \mathbb{R}$. We take

(1) The arithmetic mean:

$$A(a, b) = \frac{a+b}{2}; \quad a, b \in \mathbb{R}.$$

(2) The harmonic mean:

$$H(a, b) = \frac{2}{\frac{1}{a} + \frac{1}{b}}; \quad a, b \in \mathbb{R} \setminus \{0\}.$$

(3) The logarithmic mean:

$$L(a, b) = \frac{\ln|b| - \ln|a|}{b-a}; \quad a, b \in \mathbb{R}, a \neq b, a, b \neq 0.$$

(4) Generalized log-mean:

$$L_n(a, b) = \left[\frac{b^{n+1} - a^{n+1}}{(n+1)(b-a)} \right]^{\frac{1}{n}}; \quad a, b \in \mathbb{R}, n \in \mathbb{Z} \setminus \{-1, 0\}, a \neq b, a, b \neq 0.$$

Now using the results of Section 2, we give some applications to special means of real numbers.

Proposition 6. Let $a, b \in \mathbb{R}$, $a < b$, $0 \notin [a, b]$ and $n \in \mathbb{Z}$, $|n| \geq 2$. Then

$$\left| A \left(\left(\frac{3a+b}{4} \right)^n, \left(\frac{a+3b}{4} \right)^n \right) - L_n^n(a, b) \right| \leq |n| \left(\frac{b-a}{8} \right) A \left(|a|^{n-1}, |b|^{n-1} \right). \quad (4.1)$$

Proof. The assertion follows from Corollary 1 when applied to the function $f(x) = x^n$, $x \in [a, b]$, $n \in \mathbb{Z}$, $|n| \geq 2$. \square

Proposition 7. Let $a, b \in \mathbb{R}$, $a < b$, $0 \notin [a, b]$. Then

$$\left| H^{-1} \left(\frac{3a+b}{4}, \frac{a+3b}{4} \right) - L(a, b) \right| \leq \left(\frac{b-a}{8} \right) A \left(|a|^{-2}, |b|^{-2} \right). \quad (4.2)$$

Proof. It is a direct consequence of Corollary 1 when applied to the function, $f(x) = \frac{1}{x}$, $x \in [a, b]$. \square

Remark 1. One can get several inequalities for means by using the other results of this paper but we omit the details to the interested reader.

REFERENCES

- [1] Dragomir S S and Pearce C E M. Selected Topic on Hermite- Hadamard Inequalities and Applications, Melbourne and Adelaide, December, 2000.
- [2] Dragomir S S and Fitzpatrick S. The Hadamard's inequality for s -convex functions in the second sense, Demonstratio Math. 32 (4) (1999), 687-696.
- [3] Dragomir S S and Agarwal R P. Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, Appl. Math. Lett., 11(5) (1998) 91-95.
- [4] Hadamard J. Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, J. Math Pures Appl., 58 (1893), 171–215.
- [5] Hudzik H and Maligranda L. Some remarks on s -convex functions, Aequationes Math. 48 (1994) 100–111.
- [6] Kirmaci U S, Klaričić Bakula K, Özdemir M E and Pečarić J. Hadamard-type inequalities for s -convex functions, Appl. Math. Comput., 193(1) (2007) 26-35
- [7] Kavurmaci H, Avcı M and Özdemir M E. New inequalities of Hermite–Hadamard type for convex functions with applications arXiv: 1006.1593v1[math.CA]
- [8] Pachpatte B G. On some inequalities for convex functions, RGMIA Research Report Collection, 6(E) (2003).
- [9] Pearce C E M. and Pečarić J E. Inequalities for differentiable mappings with application to special means and quadrature formula, Appl. Math. Lett., 13(2) (2000) 51-55.
- [10] Pečarić J E, Proschan F and Tong Y L. Convex Functions, Partial Orderings, and Statistical Applications, Academic Press Inc., 1992.
- [11] Mitrinović D S. Analytic Inequalities, Springer Verlag, Berlin/New York, 1970.
- [12] Tseng K L, Hwang S R and Dragomir S S. Fejér-type inequalities (I). Journal of Inequalities and Applications Volume 2010, Article ID 531976, 7 pages.
- [13] Tunç M. On some new inequalities for convex functions, Turk J Math 35 (2011) , 1 – 7.
- [14] Tunç M. New integral inequalities for s -convex functions, RGMIA Research Report Collection Volume 13, Issue 2, 2010 Preprint Available Online: <http://ajmaa.org/RGMIA/v13n2.php>.

COLLEGE OF SCIENCE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIL, HAIL 2440, SAUDI ARABIA

E-mail address: m_amer_latif@hotmail.com

SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au