

Inequalities for the Seiffert's means in terms of the identric mean

Shaoqin Gao

College of Mathematics and Computer Science, Hebei University, Baoding, 071002, China

Abstract. Some inequalities for certain bivariate means are obtained. In particular, inequalities for those introduced by Seiffert.

§1 Introduction

For $a, b > 0$ with $a \neq b$ the first and second Seiffert's means $P(a, b)$ and $T(a, b)$ were introduced by Seiffert [1,2] as follows:

$$P(a, b) = \frac{a - b}{4 \arctan(\sqrt{a/b}) - \pi}. \quad (1.1)$$

$$T(a, b) = \frac{a - b}{2 \arctan \frac{a-b}{a+b}}. \quad (1.2)$$

Recently, the inequalities for means have been the subject of intensive research [3-21]. In particular, many remarkable inequalities for the Seiffert's means can be found in the literature [4,15-20].

Let $H(a, b) = 2ab/(a + b)$, $A(a, b) = (a+b)/2$, $G(a, b) = \sqrt{ab}$, $I(a, b) = 1/e(b^b/a^a)^{1/(b-a)}$ and $L(a, b) = (b-a)/(\log b - \log a)$ be the harmonic, arithmetic, geometric, identric and logarithmic means of two positive real numbers a and b with $a \neq b$. Then

$$\min\{a, b\} < H(a, b) < G(a, b) < L(a, b) < I(a, b) < A(a, b) < \max\{a, b\}. \quad (1.3)$$

In [1], Seiffert proved

$$L(a, b) < P(a, b) < I(a, b)$$

for all $a, b > 0$ with $a \neq b$.

The following bounds for the first Seiffert's mean $P(a, b)$ in terms of the power mean $M_r(a, b) = ((a^r + b^r)/2)^{1/r}$ ($r \neq 0$) were presented by Jagers in [18]:

$$M_{1/2} < P(a, b) < M_{2/3}(a, b) \quad (1.4)$$

for all $a, b > 0$ with $a \neq b$.

Hästö [20] found the sharp lower bound for the first Seiffert's mean as follow:

$$M_{\log 2 / \log \pi}(a, b) < P(a, b) \quad (1.5)$$

for all $a, b > 0$ with $a \neq b$.

In [3], Seiffert proved

$$P(a, b) > \frac{3A(a, b)G(a, b)}{A(a, b) + 2G(a, b)} \quad \text{and} \quad P(a, b) > \frac{2}{\pi}A(a, b) \quad (1.6)$$

This research is partly supported by the Education Department of Hebei province(2009107).

MR Subject Classification: 26D15.

Keywords: Inequalities, Bivariate means, the Seiffert's means.

for all $a, b > 0$ with $a \neq b$.

In [4], the authors found the greatest value α and the least value β such that the double inequality $\alpha A(a, b) + (1 - \alpha)H(a, b) < P(a, b) < \beta A(a, b) + (1 - \beta)H(a, b)$ holds for all $a, b > 0$ with $a \neq b$.

In [21], the author proved

$$I < A < \frac{e}{2}I, \quad A_{2/3} < I < \frac{2\sqrt{2}}{e}A_{2/3}. \quad (1.7)$$

The purpose of the present paper is to obtain the inequalities of type (1.7) for the Seiffert's means in terms of the identric mean.

§2 Main Results

In what follows we will assume, without loss of generality, that $a > b > 0$.

Theorem 2.1. *For the first Seiffert's mean, the double inequality*

$$\frac{e}{\pi}I(a, b) < P(a, b) < I(a, b)$$

holds, where the constant $\frac{e}{\pi}$ and 1 are best possible.

Proof. Let $t^2 = a/b > 1$. Consider the function

$$f(t) = \frac{P(t^2, 1)}{I(t^2, 1)} = \frac{e(t^2 - 1)t^{\frac{2t^2}{1-t^2}}}{4 \arctan t - \pi}. \quad (2.1)$$

Its logarithmic derivative is

$$\frac{f'(t)}{f(t)} = \frac{4t \ln t}{(t^2 - 1)^2(4 \arctan t - \pi)}g(t), \quad (2.2)$$

where

$$g(t) = 4 \arctan t - \pi - \frac{(t^2 - 1)^2}{t(t^2 + 1) \ln t}. \quad (2.3)$$

Simple computations lead to

$$\lim_{t \rightarrow 1^+} g(t) = 0, \quad (2.4)$$

$$g'(t) = \frac{g_1(t)}{t^2(t^2 + 1)^2 \ln^2 t}, \quad (2.5)$$

where

$$g_1(t) = 4t^2(t^2 + 1) \ln^2 t - (t^6 + 5t^4 - 5t^2 - 1) \ln t + (t^2 - 1)^2(t^2 + 1). \quad (2.6)$$

$$\lim_{t \rightarrow 1^+} g_1(t) = 0, \quad (2.7)$$

$$g'_1(t) = 8(2t^3 + t) \ln^2 t + 6(-t^5 - 2t^3 + 3t) \ln t + 5t^5 - 9t^3 + 3t + \frac{1}{t}, \quad (2.8)$$

$$\lim_{t \rightarrow 1^+} g'_1(t) = 0, \quad (2.9)$$

$$g''_1(t) = 8(6t^2 + 1) \ln^2 t + (-30t^4 - 4t^2 + 34) \ln t + 19t^4 - 39t^2 + 21 - \frac{1}{t^2}, \quad (2.10)$$

$$\lim_{t \rightarrow 1^+} g''_1(t) = 0, \quad (2.11)$$

$$g'''_1(t) = 2tg_2(t), \quad (2.12)$$

where

$$g_2(t) = 48 \ln^2 t + (-60t^2 + 44 + \frac{8}{t^2}) \ln t + 23t^2 - 41 + \frac{17}{t^2} + \frac{1}{t^4}. \quad (2.13)$$

$$\lim_{t \rightarrow 1^+} g_2(t) = 0, \quad (2.14)$$

$$g_2'(t) = \frac{2}{t} g_3(t), \quad (2.15)$$

$$g_3(t) = 48 \ln t + \left(-60t^2 - \frac{8}{t^2}\right) \ln t - 7t^2 + 22 - \frac{13}{t^2} - \frac{2}{t^4}, \quad (2.16)$$

$$\lim_{t \rightarrow 1^+} g_3(t) = 0, \quad (2.17)$$

$$g_3'(t) = 2t g_4(t), \quad (2.18)$$

where

$$g_4(t) = \left(-60 + \frac{8}{t^4}\right) \ln t - 37 + \frac{24}{t^2} + \frac{9}{t^4} + \frac{4}{t^6}. \quad (2.19)$$

$$\lim_{t \rightarrow 1^+} g_4(t) = 0, \quad (2.20)$$

$$g_4'(t) = \frac{4}{t^5} \left(-8 \ln t - 15t^4 - 12t^2 - 7 - \frac{6}{t^2}\right) < 0 \quad (2.21)$$

for $t > 1$, hence $g_4(t)$ is strictly decreasing in $[1, +\infty)$. It follows from (2.20) and (2.18) together with the monotonicity of $g_4(t)$ that $g_3'(t) < 0$, hence $g_3(t)$ is strictly decreasing in $[1, +\infty)$. From (2.17) and (2.15) together with the monotonicity of $g_3(t)$ we know that $g_2'(t) < 0$, hence $g_2(t)$ is strictly decreasing in $[1, +\infty)$.

Repeating the above procedures, we can get $g'(t) < 0$, hence $g(t)$ is strictly decreasing in $[1, +\infty)$.

From (2.4) and (2.2) together with the monotonicity of $g(t)$ we know that $f'(t) < 0$, hence $f(t)$ is strictly decreasing in $[1, +\infty)$.

Hence

$$f(t) < \lim_{t \rightarrow 1^+} f(t) = 1$$

and

$$f(t) > \lim_{t \rightarrow +\infty} f(t) = \frac{e}{\pi}.$$

The proof of the inequality $\frac{e}{\pi} I(a, b) < P(a, b) < I(a, b)$ is complete. Since $f(t)$ is continuous for $t > 1$, it follows that the constant $\frac{e}{\pi}$ and 1 are the best possible. \square

Theorem 2.2. For the second Seiffert's mean, the double inequality

$$I(a, b) < T(a, b) < \frac{2e}{\pi} I(a, b)$$

holds, where the constant 1 and $\frac{2e}{\pi}$ are best possible.

Proof. Let $t = a/b > 1$. Consider the function

$$f(t) = \frac{T(t, 1)}{I(t, 1)} = \frac{e(t-1)}{2t^{\frac{t}{t-1}} \arctan \frac{t-1}{t+1}}. \quad (2.22)$$

Its logarithmic derivative is

$$\frac{f'(t)}{f(t)} = \frac{\ln t}{(t-1)^2 \arctan \frac{t-1}{t+1}} g(t), \quad (2.23)$$

where

$$g(t) = \arctan \frac{t-1}{t+1} - \frac{(t-1)^2}{(t^2+1) \ln t}. \quad (2.24)$$

Simple computations lead to

$$\lim_{t \rightarrow 1^+} g(t) = 0, \quad (2.25)$$

$$g'(t) = \frac{g_1(t)}{(t^2 + 1)^2 \ln^2 t}, \quad (2.26)$$

where

$$g_1(t) = (t^2 + 1) \ln^2 t - 2(t^2 - 1) \ln t + (t - 1)^2 \left(t + \frac{1}{t}\right). \quad (2.27)$$

$$\lim_{t \rightarrow 1^+} g_1(t) = 0, \quad (2.28)$$

$$g_1'(t) = 2t \ln^2 t + \left(-2t + \frac{2}{t}\right) \ln t + 3t^2 - 6t + 2 + \frac{2}{t} - \frac{1}{t^2}, \quad (2.29)$$

$$\lim_{t \rightarrow 1^+} g_1'(t) = 0, \quad (2.30)$$

$$g_1''(t) = 2 \ln^2 t + \left(2 - \frac{2}{t^2}\right) \ln t + 6t - 8 + \frac{2}{t^3}, \quad (2.31)$$

$$\lim_{t \rightarrow 1^+} g_1''(t) = 0, \quad (2.32)$$

$$g_1'''(t) = \frac{1}{t} g_2(t), \quad (2.33)$$

where

$$g_2(t) = 4 \ln t + \frac{4}{t^2} \ln t + 6t - 2 - \frac{2}{t^2} - \frac{6}{t^3}. \quad (2.34)$$

$$\lim_{t \rightarrow 1^+} g_2(t) = 0, \quad (2.35)$$

$$g_2'(t) = \frac{1}{t^3} g_3(t), \quad (2.36)$$

$$g_3(t) = 4t^2 - 8 \ln t + 8 + 6t^3 + \frac{18}{t}, \quad (2.37)$$

$$\lim_{t \rightarrow 1^+} g_3(t) > 0, \quad (2.38)$$

$$g_3'(t) = 8t - \frac{8}{t} + 18t^2 - \frac{18}{t^2}, \quad (2.39)$$

$$\lim_{t \rightarrow 1^+} g_3'(t) = 0. \quad (2.40)$$

$$g_3''(t) = 8 + \frac{8}{t^2} + 36t + \frac{36}{t^3}.$$

We can see clearly that $g_3''(t) > 0$ for $t > 1$, hence $g_3'(t)$ is strictly increasing in $[1, +\infty)$. From (2.40) we have $g_3'(t) > 0$ for $t > 1$, hence $g_3(t)$ is strictly increasing in $[1, +\infty)$.

It follows from (2.38) and (2.36) together with the monotonicity of $g_3(t)$ that $g_2'(t) > 0$, hence $g_2(t)$ is strictly increasing in $[1, +\infty)$. From (2.35) and (2.33) together with the monotonicity of $g_2(t)$ we know that $g_1''(t) > 0$, hence $g_1'(t)$ is strictly increasing in $[1, +\infty)$.

Repeating the above procedures, we can get $g'(t) > 0$, hence $g(t)$ is strictly increasing in $[1, +\infty)$.

From (2.25) and (2.23) together with the monotonicity of $g(t)$ we know that $f'(t) > 0$, hence $f(t)$ is strictly increasing in $[1, +\infty)$.

Hence

$$f(t) > \lim_{t \rightarrow 1^+} f(t) = 1$$

and

$$f(t) < \lim_{t \rightarrow +\infty} f(t) = \frac{2e}{\pi}.$$

The proof of the inequality $I(a, b) < P(a, b) < \frac{2e}{\pi} I(a, b)$ is complete. Since $f(t)$ is continuous

for $t > 1$, it follows that the constant 1 and $\frac{2e}{\pi}$ are the best possible. \square

References

- [1] H.J.Seiffert, Problem 887, *Nieuw Archief voor Wiskunde*, vol 11, no. 2, pp.176-176, 1993.
- [2] H.J.Seiffert, Aufgabe β 16, *Die Wurzel*, vol 29, pp.221-222, 1995.
- [3] H.J.Seiffert, Ungleichungen für einen bestimmten mittelwert, *Nieuw Archief voor Wiskunde*, vol 13, no. 2, pp.195-198, 1995.
- [4] Y.M.Chu, Y.F.Qiu, M.K.Wang and G.D.Wang, The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean, *Journal of Inequalities and Applications*, Article ID 436457, doi: 10.1155/436457, 7 pages, 2010.
- [5] M.K.Wang, Y.M.Chu and Y.F.Qiu, Some comparison inequalities for generalized Muirhead and identric means," *Journal of Inequalities and Applications*, vol.2010, Article ID 295620,10 pages, 2010.
- [6] B.Y.Long and Y.M.Chu, Optimal inequalities for generalized logarithmic, arithmetic and geometric means, *Journal of Inequalities and Applications*, vol.2010, Article ID 806825, 10 pages, 2010.
- [7] B.Y.Long and Y.M.Chu, Optimal power mean bounds for the weighted geometric mean of classical means, *Journal of Inequalities and Applications*, vol.2010, Article ID 905679, 6 pages, 2010.
- [8] W.F.Xia, Y.M.Chu and G.D.Wang, The optimal upper and lower power mean bounds for a convex combination of the arithmetic and logarithmic means, *Abstract and Applied Analysis*, vol.2010, Article ID604804, 9 pages, 2010.
- [9] Y.M.Chu and B.Y.Long, Best possible inequalities between generalized logarithmic mean and classical means, *Abstract and Applied Analysis*, vol.2010, Article ID 303286, 13 pages, 2010.
- [10] M.Y.Shi, Y.M.Chu and Y.P.Jiang, Optimal inequalities among various means of two arguments, *Abstract and Applied Analysis*, vol.2009, Article ID 694394, 10 pages, 2009.
- [11] Y.M.Chu and W.F.Xia, Two sharp inequalities for power mean, geometric mean and harmonic mean, *Journal of Inequalities and Applications*, vol.2009, Article ID 741923, 6 pages, 2009.
- [12] Y.M.Chu and W.F.Xia, Inequalities for generalized logarithmic means, *Journal of Inequalities and Applications*, vol.2009, Article ID 763252, 7 pages, 2009.
- [13] J.J.Wen and W.L.Wang, The optimization for the inequalities of power means, *Journal of Inequalities and Applications*, vol.2006, Article ID 46782, 25 pages, 2006.
- [14] T.Hara, M.Uchiyama and S.E.Takahasi, A refinement of various mean inequalities, *Journal of Inequalities and Applications*, vol.2, no.4, pp.387-395, 1998.
- [15] E.Neuman and J.Sándor, On the Schwab-Borchardt mean, *Mathematica Pannonica*, vol.17, no.1, pp.49-59, 2006.
- [16] E.Neuman and J.Sándor, On the Schwab-Borchardt mean, *Mathematica Pannonica*, vol.14, no.2, pp.253-266, 2003.
- [17] E.Neuman and J.Sándor, On certain means of two arguments and their extensions, *International Journal of Mathematics and Mathematical Sciences*, no.16, pp.981-993, 2003.
- [18] A.A.Jagers, Solution of problem 887, *Nieuw Archief voor Wiskunde*, vol.12, pp.230-231, 1994.
- [19] P.A.Hästö, A monotonicity property of ratios of symmetric homogeneous means, *Journal of Inequalities in Pure and Applied Mathematics*, 2002, 3(5): article 71, 1-54.
- [20] P.A.Hästö, Optimal inequalities between Seiffert's mean and power mean, *Mathematical Inequalities and Applications*, Vol 7, no.1, pp. 47-53, 2004.
- [21] Edward Neuman and József Sándor, Copanion inequalities for certain bivariate means, *Applicable Analysis and Discrete Mathematics*, vol.3, 46-51, 2009.