

**ON COMPANION OF OSTROWSKI INEQUALITY FOR  
MAPPINGS WHOSE FIRST DERIVATIVES ABSOLUTE VALUE  
ARE CONVEX WITH APPLICATIONS**

MOHAMMAD W. ALOMARI\*<sup>▲</sup>, M. EMIN ÖZDEMİR<sup>♣</sup>, AND HAVVA KAVURMACI<sup>♣</sup>

ABSTRACT. Several inequalities for a companion of Ostrowski inequality for absolutely continuous mappings whose first derivatives absolute value are convex (resp. concave) are established. Applications to a composite quadrature rule, to p.d.f.'s, and to special means are provided.

1. INTRODUCTION

In 1938, Ostrowski established a very interesting inequality for differentiable mappings with bounded derivatives, as follows [6]:

**Theorem 1.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ , interior of the interval  $I$ , such that  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'(x)| \leq M$ , then the following inequality,*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq M(b-a) \left[ \frac{1}{4} + \frac{\left(x - \frac{a+b}{2}\right)^2}{(b-a)^2} \right]$$

holds for all  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is the best possible in the sense that it cannot be replaced by a smaller constant.

For recent results concerning Ostrowski inequality see [1], [2] and [3]. Also, the reader may refer to the monograph [6] where various inequalities of Ostrowski type are discussed.

In [5], Guessab and Schmeisser have proved among others, the following companion of Ostrowski inequality:

**Theorem 2.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  satisfy the Lipschitz condition, i.e.,  $|f(t) - f(s)| \leq M|t - s|$ . Then for each  $x \in [a, \frac{a+b}{2}]$ , we have the inequality,*

$$(1.1) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{8} + 2 \left( \frac{x - \frac{3a+b}{4}}{b-a} \right)^2 \right] (b-a) M.$$

---

*Date:* 26/06/2011.

*1991 Mathematics Subject Classification.* 26D10, 26A15, 26A16, 26A51.

*Key words and phrases.* Convex mappings, Hermite-Hadamard inequality, Ostrowski inequality.

<sup>▲</sup>Corresponding Author.

The constant  $1/8$  is the best possible in the sense that it cannot be replaced by a smaller constant.

We may also note that the best inequality in (1.1) is obtained for  $x = \frac{3a+b}{4}$ , giving the trapezoid type inequality,

$$(1.2) \quad \left| \frac{f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{8} M$$

The constant  $1/8$  is sharp in (1.2) in the sense mentioned above.

Companions of Ostrowski integral inequality for absolutely continuous functions was considered by Dragomir in [[7], pp.228], as follows :

**Theorem 3.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous function on  $[a, b]$ . Then we have the inequalities,

$$(1.3) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \begin{cases} \left[ \frac{1}{8} + 2 \left( \frac{x - \frac{3a+b}{4}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty, & f' \in L_\infty[a, b] \\ \left( \frac{2}{q+1} \right)^{1/q} \left[ \left( \frac{x-a}{b-a} \right)^{q+1} + \left( \frac{a+b-x}{b-a} \right)^{q+1} \right]^{1/q} (b-a)^{1/q} \|f'\|_{[a,b],p}, & p > 1, \frac{1}{p} + \frac{1}{q} = 1, \text{ and } f' \in L_p[a, b] \\ \left[ \frac{1}{4} + \left| \frac{x - \frac{3a+b}{4}}{b-a} \right| \right] \|f'\|_{[a,b],1} \end{cases}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

In [9], the following theorem which was obtained by Dragomir and Agarwal contains the Hermite-Hadamard type integral inequality:

**Theorem 4.** Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ . If  $|f'|$  is convex on  $[a, b]$ , then the following inequality holds:

$$(1.4) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{(b-a)(|f'(a)| + |f'(b)|)}{8}.$$

In [8], S.S. Dragomir established some inequalities for this companion for mappings of bounded variation. Also, Z. Liu introduced some companions of an Ostrowski type integral inequality for functions whose derivatives are absolutely continuous in [10]. Recently, N.S. Barnett et al. have proved some companions for the Ostrowski inequality and the generalized trapezoid inequality in [4].

The aim of this paper is to study a companion of Ostrowski inequality Theorem 2 for the class of functions whose derivatives in absolute value are convex (concave) functions.

## 2. RESULTS

In order to prove our results, we need the following lemma [7]:

**Lemma 1.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous mapping on  $I^\circ$ , where  $a, b \in I$  with  $a < b$ , such that  $f' \in L_1[a, b]$ . Then, the following equality holds*

$$\frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt = \frac{1}{b-a} \int_a^b p(x,t) f'(t) dt,$$

where,

$$p(x,t) = \begin{cases} t-a, & t \in [a, x] \\ t - \frac{a+b}{2}, & t \in (x, a+b-x] \\ t-b, & t \in (a+b-x, b] \end{cases},$$

for all  $x \in [a, \frac{a+b}{2}]$ .

A simple proof of the equality can be done by performing integration by parts. The details are left to the interested reader (see [7]).

Let us begin with the following result:

**Theorem 5.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous mapping on  $I^\circ$ , where  $a, b \in I$  with  $a < b$ , such that  $f' \in L_1[a, b]$ . If  $|f'|$  is convex on  $[a, b]$ , then we have the following inequality:*

$$(2.1) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{(x-a)^2}{6(b-a)} (|f'(a)| + |f'(b)|) \\ + \frac{8(x-a)^2 + 3(a+b-2x)^2}{24(b-a)} (|f'(x)| + |f'(a+b-x)|)$$

for all  $x \in [a, \frac{a+b}{2}]$ .

*Proof.* Using Lemma 1 and the modulus, we have

$$\left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{b-a} \int_a^b |p(x,t)| |f'(t)| dt \\ = \frac{1}{b-a} \left[ \int_a^x |p(x,t)| |f'(t)| dt + \int_x^{a+b-x} |p(x,t)| |f'(t)| dt + \int_{a+b-x}^b |p(x,t)| |f'(t)| dt \right]$$

Since  $|f'|$  is convex on  $[a, b] = [a, x] \cup (x, a+b-x) \cup (a+b-x, b]$ , therefore we have

$$|f'(t)| \leq \frac{t-a}{x-a} |f'(x)| + \frac{x-t}{x-a} |f'(a)|, \quad t \in [a, x];$$

$$|f'(t)| \leq \frac{t-x}{a+b-2x} |f'(a+b-x)| + \frac{a+b-x-t}{a+b-2x} |f'(x)|, \quad t \in (x, a+b-x];$$

and

$$|f'(t)| \leq \frac{t-a-b+x}{x-a} |f'(b)| + \frac{b-t}{x-a} |f'(a+b-x)|, \quad t \in (a+b-x, b];$$

which follows that,

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{b-a} \int_a^x |t-a| \left[ \frac{t-a}{x-a} |f'(x)| + \frac{x-t}{x-a} |f'(a)| \right] dt \\ & \quad + \frac{1}{b-a} \int_x^{a+b-x} \left| t - \frac{a+b}{2} \right| \left[ \frac{t-x}{a+b-2x} |f'(a+b-x)| + \frac{a+b-x-t}{a+b-2x} |f'(x)| \right] dt \\ & \quad + \frac{1}{b-a} \int_{a+b-x}^b |t-b| \left[ \frac{t-a-b+x}{x-a} |f'(b)| + \frac{b-t}{x-a} |f'(a+b-x)| \right] dt \\ & = \frac{1}{(b-a)(x-a)} \left[ |f'(x)| \frac{(x-a)^3}{3} + |f'(a)| \frac{(x-a)^3}{6} \right] \\ & \quad + \frac{1}{(b-a)(a+b-2x)} \frac{(a+b-2x)^3}{8} [|f'(a+b-x)| + |f'(x)|] \\ & \quad + \frac{1}{(b-a)(x-a)} \left[ |f'(b)| \frac{(x-a)^3}{6} + |f'(a+b-x)| \frac{(x-a)^3}{3} \right] \\ & = \frac{(x-a)^2}{6(b-a)} [|f'(a)| + 2|f'(x)| + 2|f'(a+b-x)| + |f'(b)|] \\ & \quad + \frac{(a+b-2x)^2}{8(b-a)} [|f'(a+b-x)| + |f'(x)|] \\ & = \frac{(x-a)^2}{6(b-a)} (|f'(a)| + |f'(b)|) \\ & \quad + \frac{8(x-a)^2 + 3(a+b-2x)^2}{24(b-a)} (|f'(x)| + |f'(a+b-x)|), \end{aligned}$$

where

$$\int_x^{a+b-x} (t-x) \left| t - \frac{a+b}{2} \right| dt = \int_x^{a+b-x} (a+b-x-t) \left| t - \frac{a+b}{2} \right| dt = \frac{(a+b-2x)^3}{8},$$

which completes the proof.  $\square$

An Ostrowski type inequality may be deduced as follows:

**Corollary 1.** *Let  $f$  as in Theorem 5. Additionally, if  $f$  is symmetric about the  $x$ -axis, i.e.,  $f(a+b-x) = f(x)$ , we have*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(x-a)^2}{6(b-a)} (|f'(a)| + |f'(b)|) \\ & \quad + \frac{8(x-a)^2 + 3(a+b-2x)^2}{24(b-a)} (|f'(x)| + |f'(a+b-x)|) \end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

**Remark 1.** In Theorem 5, if we choose  $x = a$ , then we get

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{b-a}{8} (|f'(a)| + |f'(b)|).$$

which is the inequality in (1.4).

**Corollary 2.** In Theorem 5, if we choose

(1)  $x = \frac{3a+b}{4}$ , then we get

$$(2.2) \quad \left| \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{(b-a)}{96} \left[ |f'(a)| + 5 \left| f'\left(\frac{3a+b}{4}\right) \right| + 5 \left| f'\left(\frac{a+3b}{4}\right) \right| + |f'(b)| \right]$$

(2)  $x = \frac{a+b}{2}$ , then we get

(2.3)

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{24} \left[ |f'(a)| + 4 \left| f'\left(\frac{a+b}{2}\right) \right| + |f'(b)| \right].$$

Another result may be considered using the Hölder inequality, as follows:

**Theorem 6.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous mapping on  $I^\circ$ , where  $a, b \in I$  with  $a < b$ , such that  $f' \in L_1[a, b]$ . If  $|f'|^q$ ,  $q > 1$  is convex on  $[a, b]$ , then we have the following inequality:

$$(2.4) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{2^{1/q} (b-a)^{1/p}} \left\{ (x-a)^2 [|f'(a)|^q + |f'(x)|^q]^{1/q} \right. \\ \left. + \frac{(a+b-2x)^2}{2} [|f'(x)|^q + |f'(a+b-x)|^q]^{1/q} \right. \\ \left. + (x-a)^2 [|f'(a+b-x)|^q + |f'(b)|^q]^{1/q} \right\},$$

for all  $x \in [a, \frac{a+b}{2}]$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* Using Lemma 1 and the Hölder inequality, we have

$$\begin{aligned}
(2.5) \quad & \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
& \leq \frac{1}{b-a} \int_a^b |p(x,t)| |f'(t)| dt \\
& = \frac{1}{b-a} \left[ \int_a^x |t-a| |f'(t)| dt + \int_x^{a+b-x} \left| t - \frac{a+b}{2} \right| |f'(t)| dt + \int_{a+b-x}^b |t-b| |f'(t)| dt \right] \\
& \leq \frac{1}{b-a} \left[ \left( \int_a^x |t-a|^p dt \right)^{1/p} \left( \int_a^x |f'(t)|^q dt \right)^{1/q} \right. \\
& \quad + \left( \int_x^{a+b-x} \left| t - \frac{a+b}{2} \right|^p dt \right)^{1/p} \left( \int_x^{a+b-x} |f'(t)|^q dt \right)^{1/q} \\
& \quad \left. + \left( \int_{a+b-x}^b |t-b|^p dt \right)^{1/p} \left( \int_{a+b-x}^b |f'(t)|^q dt \right)^{1/q} \right]
\end{aligned}$$

Since  $|f'|^q$  is convex on  $[a, b] = [a, x] \cup (x, a+b-x) \cup (a+b-x, b]$ , therefore we have

$$\begin{aligned}
|f'(t)|^q & \leq \frac{t-a}{x-a} |f'(x)|^q + \frac{x-t}{x-a} |f'(a)|^q, \quad t \in [a, x]; \\
|f'(t)|^q & \leq \frac{t-x}{a+b-2x} |f'(a+b-x)|^q + \frac{a+b-x-t}{a+b-2x} |f'(x)|^q, \quad t \in (x, a+b-x);
\end{aligned}$$

and

$$|f'(t)|^q \leq \frac{t-a-b+x}{x-a} |f'(b)|^q + \frac{b-t}{x-a} |f'(a+b-x)|^q, \quad t \in (a+b-x, b];$$

which follows that,

$$\begin{aligned}
& \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
& \leq \frac{1}{b-a} \left[ \left( \int_a^x |t-a|^p dt \right)^{1/p} \left( \int_a^x \left[ \frac{t-a}{x-a} |f'(x)|^q + \frac{x-t}{x-a} |f'(a)|^q \right] dt \right)^{1/q} \right. \\
& \quad + \left( \int_x^{a+b-x} \left| t - \frac{a+b}{2} \right|^p dt \right)^{1/p} \\
& \quad \times \left( \int_x^{a+b-x} \left[ \frac{t-x}{a+b-2x} |f'(a+b-x)|^q + \frac{a+b-x-t}{a+b-2x} |f'(x)|^q \right] dt \right)^{1/q} \\
& \quad + \frac{1}{b-a} \left( \int_{a+b-x}^b |t-b|^p dt \right)^{1/p} \\
& \quad \left. \times \left( \int_{a+b-x}^b \left[ \frac{t-a-b+x}{x-a} |f'(b)|^q + \frac{b-t}{x-a} |f'(a+b-x)|^q \right] dt \right)^{1/q} \right]
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{b-a} \left[ \left( \frac{(x-a)^{p+1}}{(p+1)} \right)^{1/p} \left( \frac{x-a}{2} \right)^{1/q} (|f'(a)|^q + |f'(x)|^q)^{1/q} \right. \\
&\quad + \left( \frac{2}{(p+1)} \left( \frac{a+b}{2} - x \right)^{p+1} \right)^{1/p} \left( \frac{a+b}{2} - x \right)^{1/q} (|f'(x)|^q + |f'(a+b-x)|^q)^{1/q} \\
&\quad \left. + \left( \frac{(x-a)^{p+1}}{(p+1)} \right)^{1/p} \left( \frac{x-a}{2} \right)^{1/q} (|f'(a+b-x)|^q + |f'(b)|^q)^{1/q} \right] \\
&= \frac{1}{2^{1/q} (b-a) (p+1)^{1/p}} \left[ (x-a)^2 (|f'(a)|^q + |f'(x)|^q)^{1/q} \right. \\
&\quad + \frac{(a+b-2x)^2}{2} (|f'(x)|^q + |f'(a+b-x)|^q)^{1/q} \\
&\quad \left. + (x-a)^2 (|f'(a+b-x)|^q + |f'(b)|^q)^{1/q} \right],
\end{aligned}$$

since  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $q > 1$ , which completes the proof.  $\square$

**Corollary 3.** *Let  $f$  as in Theorem 6. Additionally, if  $f$  is symmetric about the  $x$ -axis, i.e.,  $f(a+b-x) = f(x)$ , we have*

(2.6)

$$\begin{aligned}
\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| &\leq \frac{1}{2^{1/q} (b-a) (p+1)^{1/p}} \\
&\quad \times \left\{ (x-a)^2 [ |f'(a)|^q + |f'(x)|^q ]^{1/q} \right. \\
&\quad + \frac{(a+b-2x)^2}{2} [ |f'(x)|^q + |f'(a+b-x)|^q ]^{1/q} \\
&\quad \left. + (x-a)^2 [ |f'(a+b-x)|^q + |f'(b)|^q ]^{1/q} \right\}
\end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

**Corollary 4.** *In Theorem 6, if we choose*

(1)  $x = a$ , then we get

(2.7)

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{2^{1+\frac{1}{q}} (1+p)^{1/p}} (|f'(a)|^q + |f'(b)|^q)^{1/q}.$$

(2)  $x = \frac{3a+b}{4}$ , then we get

(2.8)

$$\begin{aligned} & \left| \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(b-a)}{2^{4+\frac{1}{q}}(1+p)^{1/p}} \left\{ \left( |f'(a)|^q + \left| f'\left(\frac{3a+b}{4}\right) \right|^q \right)^{1/q} \right. \\ & \quad \left. + 2 \left( \left| f'\left(\frac{3a+b}{4}\right) \right|^q + \left| f'\left(\frac{a+3b}{4}\right) \right|^q \right)^{1/q} + \left( \left| f'\left(\frac{a+3b}{4}\right) \right|^q + |f'(b)|^q \right)^{1/q} \right\}. \end{aligned}$$

(3)  $x = \frac{a+b}{2}$ , then we get

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(b-a)}{2^{2+\frac{1}{q}}(1+p)^{1/p}} \left[ \left( |f'(a)|^q + \left| f'\left(\frac{a+b}{2}\right) \right|^q \right)^{1/q} + \left( \left| f'\left(\frac{a+b}{2}\right) \right|^q + |f'(b)|^q \right)^{1/q} \right]. \end{aligned}$$

**Remark 2.** One may get various inequalities for mappings whose first derivatives absolute value are convex by applying the power-mean inequality. We left the details to the interested reader.

The following result holds for concave mappings.

**Theorem 7.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous mapping on  $I^\circ$ , where  $a, b \in I$  with  $a < b$ , such that  $f' \in L_1[a, b]$ . If  $|f'|^q$ ,  $q > 1$  is concave on  $[a, b]$ , then we have the following inequality:

(2.9)

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{(b-a)(1+p)^{1/p}} \\ & \quad \times \left[ (x-a)^2 \left( \left| f'\left(\frac{a+x}{2}\right) \right| + \left| f'\left(\frac{2b+a-x}{2}\right) \right| \right) + \frac{(a+b-2x)^2}{2} \left| f'\left(\frac{a+b}{2}\right) \right| \right] \end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ .



*Proof.* From Lemma 1, and by (2.5), we have

$$\begin{aligned} & \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{b-a} \left[ \left( \int_a^x |t-a|^p dt \right)^{1/p} \left( \int_a^x |f'(t)|^q dt \right)^{1/q} \right. \\ & \quad + \left( \int_x^{a+b-x} \left| t - \frac{a+b}{2} \right|^p dt \right)^{1/p} \left( \int_x^{a+b-x} |f'(t)|^q dt \right)^{1/q} \\ & \quad \left. + \left( \int_{a+b-x}^b |t-b|^p dt \right)^{1/p} \left( \int_{a+b-x}^b |f'(t)|^q dt \right)^{1/q} \right]. \end{aligned}$$

Now, let us write,

$$\int_a^x |f'(t)|^q dt = (x-a) \int_0^1 |f'(\lambda x + (1-\lambda)a)|^q d\lambda,$$

$$\int_x^{a+b-x} |f'(t)|^q dt = (a+b-2x) \int_0^1 |f'(\lambda(a+b-x) + (1-\lambda)x)|^q d\lambda,$$

and

$$\int_{a+b-x}^b |f'(t)|^q dt = (x-a) \int_0^1 |f'(\lambda b + (1-\lambda)(a+b-x))|^q d\lambda.$$

Since  $|f'|^q$ ,  $q > 1$  is concave on  $[a, b] = [a, x] \cup (x, a+b-x) \cup (a+b-x, b]$ , we can use the Jensen integral inequality to obtain

$$\begin{aligned} & (x-a) \int_0^1 |f'(\lambda x + (1-\lambda)a)|^q d\lambda \\ & = (x-a) \int_0^1 \lambda^0 |f'(\lambda x + (1-\lambda)a)|^q d\lambda \\ & \leq (x-a) \left( \int_0^1 \lambda^0 d\lambda \right) \left| f' \left( \frac{1}{\int_0^1 \lambda^0 d\lambda} \int_0^1 (\lambda x + (1-\lambda)a) d\lambda \right) \right|^q \\ & = (x-a) \left| f' \left( \frac{x+a}{2} \right) \right|^q \end{aligned}$$

and analogously

$$(a+b-2x) \int_0^1 |f'(\lambda(a+b-x) + (1-\lambda)x)|^q d\lambda \leq (a+b-2x) \left| f' \left( \frac{a+b}{2} \right) \right|^q,$$

$$(x-a) \int_0^1 |f'(\lambda b + (1-\lambda)(a+b-x))|^q d\lambda \leq (x-a) \left| f' \left( \frac{2b+a-x}{2} \right) \right|^q.$$

Combining all above inequalities, we get

$$\begin{aligned}
& \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
& \leq \frac{1}{b-a} \left[ \left( \frac{(x-a)^{p+1}}{(p+1)} \right)^{1/p} (x-a)^{1/q} \left| f' \left( \frac{x+a}{2} \right) \right| \right] \\
& \quad + \left( \frac{2}{(p+1)} \left( \frac{a+b}{2} - x \right)^{p+1} \right)^{1/p} (a+b-2x)^{1/q} \left| f' \left( \frac{a+b}{2} \right) \right| \\
& \quad + \left( \frac{(x-a)^{p+1}}{(p+1)} \right)^{1/p} (x-a)^{1/q} \left| f' \left( \frac{2b+a-x}{2} \right) \right| \\
& = \frac{1}{(b-a)(1+p)^{1/p}} \\
& \quad \times \left[ (x-a)^2 \left( \left| f' \left( \frac{a+x}{2} \right) \right| + \left| f' \left( \frac{2b+a-x}{2} \right) \right| \right) + \frac{(a+b-2x)^2}{2} \left| f' \left( \frac{a+b}{2} \right) \right| \right],
\end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $q > 1$ , which is required.  $\square$

Therefore, we may state the following Ostrowski type inequality:

**Corollary 5.** *Let  $f$  as in Theorem 7. Additionally, if  $f$  is symmetric about the  $x$ -axis, i.e.,  $f(a+b-x) = f(x)$ , we have*

(2.10)

$$\begin{aligned}
& \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
& \leq \frac{1}{(b-a)(1+p)^{1/p}} \\
& \quad \times \left[ (x-a)^2 \left( \left| f' \left( \frac{a+x}{2} \right) \right| + \left| f' \left( \frac{2b+a-x}{2} \right) \right| \right) + \frac{(a+b-2x)^2}{2} \left| f' \left( \frac{a+b}{2} \right) \right| \right],
\end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

**Corollary 6.** *In Theorem 7, if we choose*

(1)  $x = a$ , then we get

(2.11)

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{2(1+p)^{1/p}} \left| f' \left( \frac{a+b}{2} \right) \right|.$$

(2)  $x = \frac{3a+b}{4}$ , then we get

(2.12)

$$\begin{aligned} & \left| \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(b-a)}{16(1+p)^{1/p}} \left[ \left| f'\left(\frac{7a+b}{8}\right) \right| + 2 \left| f'\left(\frac{a+b}{2}\right) \right| + \left| f'\left(\frac{a+7b}{8}\right) \right| \right]. \end{aligned}$$

(3)  $x = \frac{a+b}{2}$ , then we get

(2.13)

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{4(1+p)^{1/p}} \left[ \left| f'\left(\frac{3a+b}{4}\right) \right| + \left| f'\left(\frac{a+3b}{4}\right) \right| \right].$$

### 3. A COMPOSITE QUADRATURE FORMULA

Let  $I_n : a = x_0 < x_1 < \dots < x_n = b$  be a division of the interval  $[a, b]$  and  $h_i = x_{i+1} - x_i$ , ( $i = 0, 1, 2, \dots, n-1$ ).

Consider the general quadrature formula

$$(3.1) \quad Q_n(I_n, f) := \frac{1}{2} \sum_{i=0}^{n-1} \left[ f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i.$$

The following result holds.

**Theorem 8.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous mapping on  $I^\circ$ , where  $a, b \in I$  with  $a < b$ , such that  $f' \in L_1[a, b]$ . If  $|f'|$  is convex on  $[a, b]$ . Then, we have*

$$(3.2) \quad \int_a^b f(t) dt = Q_n(I_n, f) + R_n(I_n, f).$$

where,  $Q_n(I_n, f)$  is defined by formula (3.1), and the remainder term  $R_n(I_n, f)$  satisfies the error estimates

$$|R_n(I_n, f)| \leq \frac{1}{96} \sum_{i=0}^{n-1} h_i^2 \left[ |f'(x_i)| + 5 \left| f'\left(\frac{3x_i + x_{i+1}}{4}\right) \right| + 5 \left| f'\left(\frac{x_i + 3x_{i+1}}{4}\right) \right| + |f'(x_{i+1})| \right].$$

*Proof.* Applying inequality (3.1) and (3.2) on the intervals  $[x_i, x_{i+1}]$ , we may state that

$$R_i(I_i, f) = \int_{x_i}^{x_{i+1}} f(t) dt - \frac{1}{2} \left[ f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i.$$

Summing the above inequality over  $i$  from 0 to  $n - 1$ , we get

$$\begin{aligned} R_n(I_n, f) &= \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} f(t) dt - \frac{1}{2} \sum_{i=0}^{n-1} \left[ f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i \\ &= \int_a^b f(t) dt - \frac{1}{2} \sum_{i=0}^{n-1} \left[ f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i, \end{aligned}$$

which follows from (2.2), that

$$\begin{aligned} |R_n(I_n, f)| &= \left| \int_a^b f(t) dt - \frac{1}{2} \sum_{i=0}^{n-1} \left[ f\left(\frac{3x_i + x_{i+1}}{4}\right) + f\left(\frac{x_i + 3x_{i+1}}{4}\right) \right] h_i \right| \\ &\leq \frac{1}{96} \sum_{i=0}^{n-1} h_i^2 \left[ |f'(x_i)| + 5 \left| f'\left(\frac{3x_i + x_{i+1}}{4}\right) \right| + 5 \left| f'\left(\frac{x_i + 3x_{i+1}}{4}\right) \right| + |f'(x_{i+1})| \right]. \end{aligned}$$

which completes the proof.  $\square$

**Remark 3.** One may state more inequalities, using (2.8) and (2.12). We shall omit the details.

#### 4. APPLICATIONS FOR P.D.F.'S

Let  $X$  be a random variable taking values in the finite interval  $[a, b]$ , with the probability density function  $f : [a, b] \rightarrow [0, 1]$  with the cumulative distribution function  $F(x) = Pr(X \leq x) = \int_a^x f(t) dt$ .

**Theorem 9.** With the assumptions of Theorem 5, we have the inequality

$$\begin{aligned} &\left| \frac{1}{2} [F(x) + F(a + b - x)] - \frac{b - E(X)}{b - a} \right| \\ &\leq \frac{(x - a)^2}{6(b - a)} (|F'(a)| + |F'(b)|) \\ &\quad + \frac{8(x - a)^2 + 3(a + b - 2x)^2}{24(b - a)} (|F'(x)| + |F'(a + b - x)|) \end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ , where  $E(X)$  is the expectation of  $X$ .

*Proof.* In the proof of Theorem 5, let  $f = F$ , and taking into account that

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt.$$

We left the details to the interested reader.  $\square$

**Corollary 7.** In Theorem 9, if we choose  $x = \frac{3a+b}{4}$ , then we get

$$\begin{aligned} &\left| \frac{1}{2} \left[ F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) \right] - \frac{b - E(X)}{b - a} \right| \\ &\leq \frac{(b - a)}{96} \left[ |F'(a)| + 5 \left| F'\left(\frac{3a+b}{4}\right) \right| + 5 \left| F'\left(\frac{a+3b}{4}\right) \right| + |F'(b)| \right]. \end{aligned}$$

**Corollary 8.** *In Theorem 9, if  $F$  is symmetric about the  $x$ -axis, i.e.,  $F(a+b-x) = F(x)$ , we have*

$$\begin{aligned} & \left| F(x) - \frac{b - E(X)}{b - a} \right| \\ & \leq \frac{(x-a)^2}{6(b-a)} (|F'(a)| + |F'(b)|) + \frac{8(x-a)^2 + 3(a+b-2x)^2}{24(b-a)} (|F'(x)| + |F'(a+b-x)|) \end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

**Remark 4.** *One may state more inequalities, using Theorem 6 and Theorem 7. We shall omit the details.*

## 5. APPLICATIONS FOR SPECIAL MEANS

Recall the following means which could be considered extensions of arithmetic, logarithmic and generalized logarithmic for positive real numbers.

(1) The arithmetic mean:

$$A = A(a, b) = \frac{a+b}{2}; \quad a, b \in \mathbb{R}^+$$

(2) The logarithmic mean:

$$L(a, b) = \frac{b-a}{\ln|b| - \ln|a|}; \quad |a| \neq |b|, \quad a, b \in \mathbb{R}^+$$

(3) The generalized logarithmic mean:

$$L_n(a, b) = \left[ \frac{b^{n+1} - a^{n+1}}{(b-a)(n+1)} \right]^{\frac{1}{n}}; \quad n \in \mathbb{Z} \setminus \{-1, 0\}, \quad a, b \in \mathbb{R}^+, \quad a \neq b$$

(4) The identric mean:

$$I(a, b) = \begin{cases} \frac{1}{e} \left( \frac{b^b}{a^a} \right)^{\frac{1}{b-a}}, & a \neq b \\ a, & a = b \end{cases} \quad a, b \in \mathbb{R}^+.$$

Now using Results, we give some applications to special means for positive real numbers.

**Proposition 1.** *Let  $a, b \in \mathbb{R}^+$ ,  $a < b$ . Then, we have*

$$\left| A \left( \frac{4}{3a+b}, \frac{4}{a+3b} \right) - L^{-1}(a, b) \right| \leq \frac{(b-a)}{96} \left[ \frac{a^2 + b^2}{a^2 b^2} + 80 \left( \frac{1}{(3a+b)^2} + \frac{1}{(a+3b)^2} \right) \right].$$

*Proof.* The assertion was obtained by the inequality in (2.2) applied to the convex mapping  $f : [a, b] \rightarrow \mathbb{R}$ ,  $f(x) = \frac{1}{x}$ .  $\square$

**Proposition 2.** *Let  $a, b \in \mathbb{R}^+$ ,  $a < b$ , and  $p \in \mathbb{Z}$ ,  $|p| \geq 2$ . Then,*

$$|A^p(a, b) - L_p^p(a, b)| \leq \frac{p(b-a)}{24} [a^{p-1} + 4A^{p-1}(a, b) + b^{p-1}].$$

*Proof.* The assertion was obtained by the inequality in (2.3) applied to the convex mapping  $f : [a, b] \rightarrow \mathbb{R}$ ,  $f(x) = x^p$ .  $\square$

**Proposition 3.** *Let  $a, b \in \mathbb{R}^+$ ,  $a < b$ . Then, we have*

$$|\ln I - A(\ln a, \ln b)| \leq \frac{(b-a)}{2^{1+\frac{1}{q}}(1+p)^{\frac{1}{p}}} \left( \frac{1}{a^q} + \frac{1}{b^q} \right)^{\frac{1}{q}}.$$

*Proof.* The assertion was obtained by the inequality in (2.7) applied to the convex mapping  $f : [a, b] \rightarrow [0, \infty)$ ,  $f(x) = -\ln x$ .  $\square$

#### REFERENCES

- [1] M. Alomari and M. Darus, Some Ostrowski type inequalities for quasi-convex functions with applications to special means, *RGMA Preprint* 13 (2) (2010) article No. 3. [<http://rgmia.org/papers/v13n2/quasi-convex.pdf>]
- [2] M. Alomari, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are  $s$ -convex in the second sense, *Appl. Math. Lett.*, 23 (2010), 1071–1076.
- [3] M.W. Alomari, A companion of Ostrowski's inequality with applications, *Trans. J. Math. Mech.*, to appear.
- [4] N.S. Barnett, S.S. Dragomir and I. Gomma, A companion for the Ostrowski and the generalised trapezoid inequalities, *J. Mathematical and Computer Modelling*, 50 (2009), 179–187.
- [5] A. Guessab and G. Schmeisser, Sharp integral inequalities of the Hermite-Hadamard type, *J. Approx. Th.*, 115 (2002), 260–288.
- [6] S.S. Dragomir and Th.M. Rassias (Ed.), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publishers, Dordrecht, 2002.
- [7] S.S. Dragomir, Some companions of Ostrowski's inequality for absolutely continuous functions and applications, *Bull. Korean Math. Soc.*, 42 (2005), No. 2, pp. 213–230.
- [8] S.S. Dragomir, A companion of Ostrowski's inequality for functions of bounded variation and applications, *RGMA Preprint*, Vol. 5 Supp. (2002) article No. 28. [<http://ajmaa.org/RGMA/papers/v5e/COIFBVApp.pdf>].
- [9] S.S. Dragomir, R.P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Math. Lett.*, 11(5) (1998) 91-95.
- [10] Z. Liu, Some companions of an Ostrowski type inequality and applications, *J. Ineq. Pure & Appl. Math.*, Volume 10 (2009), Issue 2, Article 52, 12 pp.

\*DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, JERASH UNIVERSITY, JERASH, JORDAN.  
E-mail address: [mwomath@gmail.com](mailto:mwomath@gmail.com)

<sup>♣</sup>ATATÜRK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, 25240, CAMPUS, ERZURUM, TURKEY  
E-mail address: [emos@atauni.edu.tr](mailto:emos@atauni.edu.tr)

E-mail address: [hkavurmaci@atauni.edu.tr](mailto:hkavurmaci@atauni.edu.tr)