

**SOME GRÜSS TYPE INEQUALITIES FOR
RIEMANN–STIELTJES INTEGRAL AND APPLICATIONS**

MOHAMMAD WAJEEH ALOMARI

ABSTRACT. In this paper a Grüss' type inequalities for Riemann–Stieltjes integral are proved. Applications to the approximation problem of the Riemann–Stieltjes are also pointed out.

1. INTRODUCTION

In 1935, G. Grüss proved the following famous inequality regarding the integral of the product of two functions and the product of the integrals:

$$(1.1) \quad \left| \frac{1}{b-a} \int_a^b f(x)g(x) dx - \left(\frac{1}{b-a} \int_a^b f(x) dx \right) \cdot \left(\frac{1}{b-a} \int_a^b g(x) dx \right) \right| \leq \frac{1}{4} (\Phi - \phi) (\Gamma - \gamma)$$

provided that f and g are two integrable functions on $[a, b]$ and satisfying the condition $\phi \leq f(x) \leq \Phi$ and $\gamma \leq g(x) \leq \Gamma$, for all $x \in [a, b]$. The constant $\frac{1}{4}$ is best possible in the sense that it cannot be replaced by a smaller one.

In [15], Dragomir and Fedotov, have established the following functional:

$$(1.2) \quad \mathcal{D}(f; u) := \int_a^b f(x) du(x) - \frac{u(b) - u(a)}{b-a} \int_a^b f(t) dt,$$

provided that the Stieltjes integral $\int_a^b f(x) du(x)$ and the Riemann integral $\int_a^b f(t) dt$ exist.

In the same paper [15], the authors have proved the following inequality:

Theorem 1. *Let $f, u : [a, b] \rightarrow \mathbb{R}$ be such that u is of bounded variation on $[a, b]$ and f is Lipschitzian with the constant $K > 0$. Then we have*

$$(1.3) \quad |\mathcal{D}(f; u)| \leq \frac{1}{2} K (b-a) \bigvee_a^b(u),$$

The constant $\frac{1}{2}$ is sharp in the sense that it cannot be replaced by a smaller quantity.

Also, in [7], Dragomir has obtained the following inequality:

Date: November 20, 2011.

2000 Mathematics Subject Classification. 26D15, 26D20, 41A55.

Key words and phrases. Ostrowski inequality, Grüss inequality, Bounded variation, Lipschitzian, Monotonic, Riemann–Stieltjes integral.

Theorem 2. Let $f, u : [a, b] \rightarrow \mathbb{R}$ be such that u is Lipschitzian on $[a, b]$, i.e.,

$$|u(y) - u(x)| \leq L|x - y|, \forall x, y \in [a, b], \quad (L > 0)$$

and f is Riemann integrable on $[a, b]$.

If $m, M \in \mathbb{R}$, are such that $m \leq f(x) \leq M$, for any $x \in [a, b]$, then the inequality

$$(1.4) \quad |\mathcal{D}(f; u)| \leq \frac{1}{2}L(M - m)(b - a)$$

The constant $\frac{1}{2}$ is sharp in the sense that it cannot be replaced by a smaller quantity.

For other recent inequalities for the Riemann–Stieltjes integral, see [1]–[16] and the references therein.

The aim of this paper, is to obtain several new bounds for $\mathcal{D}(f; u)$. More specifically, the integrand f is assumed to be monotonic nondecreasing on both $[a, x]$ and $[x, b]$, and the integrator u is to be of bounded variation, Lipschitzian and monotonic on $[a, b]$.

2. THE CASE OF BOUNDED VARIATION INTEGRATORS

The following result holds:

Theorem 3. Let $x \in [a, b]$. Let $u : [a, b] \rightarrow \mathbb{R}$ be a mapping of bounded variation on $[a, b]$ and $f : [a, b] \rightarrow \mathbb{R}$ is continuous on $[a, b]$. Assume that f is monotonic nondecreasing on both $[a, x]$ and $[x, b]$. Then we have the inequality:

$$(2.1) \quad |\mathcal{D}(f; u)| \leq [f(b) - f(a)] \cdot \bigvee_a^b(u).$$

Proof. It is well-known that for a continuous function $p : [a, b] \rightarrow \mathbb{R}$ and a function $\nu : [a, b] \rightarrow \mathbb{R}$ of bounded variation, one has the inequality

$$\left| \int_a^b p(t) d\nu(t) \right| \leq \sup_{t \in [a, b]} |p(t)| \bigvee_a^b(\nu).$$

Therefore, as u is of bounded variation on $[a, b]$, we have

$$(2.2) \quad \begin{aligned} |\mathcal{D}(f; u)| &= \left| \int_a^b \left[f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right] du(x) \right| \\ &\leq \sup_{x \in [a, b]} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \cdot \bigvee_a^b(u) \\ &= \frac{1}{b-a} \sup_{x \in [a, b]} \left| \int_a^b [f(x) - f(t)] dt \right| \cdot \bigvee_a^b(u) \\ &= \frac{1}{b-a} \sup_{x \in [a, b]} \int_a^b |f(x) - f(t)| dt \cdot \bigvee_a^b(u) \end{aligned}$$

As f is monotonic nondecreasing on $[a, x]$, and monotonic nondecreasing on $[x, b]$, we get

$$\begin{aligned} \int_a^b |f(x) - f(t)| dt &\leq \int_a^x |f(x) - f(t)| dt + \int_x^b |f(x) - f(t)| dt \\ &= (x-a)f(x) - \int_a^x f(t) dt + \int_x^b f(t) dt - (b-x)f(x) \\ &= (2x-a-b)f(x) + \int_x^b f(t) dt - \int_a^x f(t) dt. \end{aligned}$$

On utilizing the monotonicity property of f on both intervals, we have

$$\int_x^b f(t) dt \leq (b-x)f(b), \quad \text{and} \quad \int_a^x f(t) dt \geq (x-a)f(a),$$

which implies that,

$$\int_a^b |f(x) - f(t)| dt \leq (2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a).$$

Taking 'sup' for both side, we get

$$\begin{aligned} \sup_{x \in [a, b]} \int_a^b |f(x) - f(t)| dt &\leq \sup_{x \in [a, b]} \{(2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a)\} \\ (2.3) \qquad \qquad \qquad &= (b-a)[f(b) - f(a)] \end{aligned}$$

combining (2.2) and (2.3), we get

$$\begin{aligned} |\mathcal{D}(f; u)| &\leq \frac{1}{b-a} \sup_{x \in [a, b]} \int_a^b |f(x) - f(t)| dt \cdot \bigvee_a^b(u) \\ &\leq [f(b) - f(a)] \cdot \bigvee_a^b(u), \end{aligned}$$

and the theorem is proved. \square

Corollary 1. *Let f as in Theorem 3. Let $u \in C^{(1)}[a, b]$. Then we have the inequality*

$$(2.4) \qquad \qquad \qquad |\mathcal{D}(f; u)| \leq [f(b) - f(a)] \cdot \|u'\|_{1, [a, b]}$$

where $\|\cdot\|_1$ is the L_1 norm, namely $\|u'\|_{1, [a, b]} := \int_a^b |u'(t)| dt$.

Corollary 2. *Let f as in Theorem 3. Let $u : [a, b] \rightarrow \mathbb{R}$ be a Lipschitzian mapping with the constant $L > 0$. Then we have the inequality*

$$(2.5) \qquad \qquad \qquad |\mathcal{D}(f; u)| \leq L(b-a)[f(b) - f(a)].$$

Corollary 3. *Let f as in Theorem 3. Let $u : [a, b] \rightarrow \mathbb{R}$ be a monotonic mapping. Then we have the inequality*

$$(2.6) \qquad \qquad \qquad |\mathcal{D}(f; u)| \leq [f(b) - f(a)] \cdot |u(b) - u(a)|.$$

3. THE CASE OF LIPSCHITZIAN INTEGRATORS

The following result holds:

Theorem 4. *Let $x \in [a, b]$. Let $u, f : [a, b] \rightarrow \mathbb{R}$ be such that u is L -Lipschitzian on $[a, b]$, and f is monotonic nondecreasing on both $[a, x]$ and $[x, b]$. Then we have the inequality:*

$$(3.1) \quad |\mathcal{D}(f; u)| \leq L \left[\frac{1}{2} (b-a) (f(b) - f(a)) + \int_a^b f(x) dx \right].$$

Proof. It is well-known that for a Riemann integrable function $p : [a, b] \rightarrow \mathbb{R}$ and L -Lipschitzian function $\nu : [a, b] \rightarrow \mathbb{R}$, one has the inequality

$$\left| \int_a^b p(t) d\nu(t) \right| \leq L \int_a^b |p(t)| dt.$$

Therefore, as u is L -Lipschitzian on $[a, b]$, we have

$$(3.2) \quad \begin{aligned} |\mathcal{D}(f; u)| &= \left| \int_a^b \left[f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right] du(x) \right| \leq L \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| dx \\ &= \frac{L}{b-a} \int_a^b \left| \int_a^b [f(x) - f(t)] dt \right| dx \end{aligned}$$

As f is monotonic nondecreasing on $[a, x]$, and monotonic nondecreasing on $[x, b]$, we get

$$\begin{aligned} \left| \int_a^b [f(x) - f(t)] dt \right| &\leq \int_a^b |f(x) - f(t)| dt \\ &= \int_a^x |f(x) - f(t)| dt + \int_x^b |f(x) - f(t)| dt \\ &= (x-a) f(x) - \int_a^x f(t) dt + \int_x^b f(t) dt - (b-x) f(x) \\ &= (2x-a-b) f(x) + \int_x^b f(t) dt - \int_a^x f(t) dt. \end{aligned}$$

On utilizing the monotonicity property of f on both intervals, we have

$$\int_x^b f(t) dt \leq (b-x) f(b), \quad \text{and} \quad \int_a^x f(t) dt \geq (x-a) f(a),$$

which implies that,

$$(3.3) \quad \int_a^b |f(x) - f(t)| dt \leq (2x-a-b) f(x) + (b-x) f(b) - (x-a) f(a).$$

Combining (3.2) and (3.3), we get

$$\begin{aligned}
|\mathcal{D}(f; u)| &\leq \frac{L}{b-a} \int_a^b \left| \int_a^b [f(x) - f(t)] dt \right| dx \\
&\leq \frac{L}{b-a} \int_a^b [(2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a)] dx \\
&= \frac{1}{2}L(b-a)[f(b) - f(a)] + \frac{L}{(b-a)} \int_a^b (2x-a-b)f(x) dx \\
&\leq \frac{1}{2}L(b-a)[f(b) - f(a)] + \frac{L}{(b-a)} \cdot \max_{x \in [a,b]} \{2x-a-b\} \cdot \int_a^b f(x) dx \\
&= L \left[\frac{1}{2}(b-a)(f(b) - f(a)) + \int_a^b f(x) dx \right]
\end{aligned}$$

and the theorem is proved. \square

4. THE CASE OF MONOTONIC INTEGRATORS

Theorem 5. *Let $x \in [a, b]$. Let $u, f : [a, b] \rightarrow \mathbb{R}$ be a continuous mappings on $[a, b]$. Assume that u is monotonic nondecreasing mapping on $[a, b]$ and $f : [a, b] \rightarrow \mathbb{R}$ is monotonic nondecreasing on both $[a, x]$ and $[x, b]$. Then we have the inequality:*

$$(4.1) \quad |\mathcal{D}(f; u)| \leq 2u(b) \cdot \left[f(b) - \frac{1}{b-a} \int_a^b f(x) dx \right].$$

Proof. It is well-known that for a monotonic non-decreasing function $\nu : [a, b] \rightarrow \mathbb{R}$ and continuous function $p : [a, b] \rightarrow \mathbb{R}$, one has the inequality

$$\left| \int_a^b p(t) d\nu(t) \right| \leq \int_a^b |p(t)| d\nu(t).$$

Therefore, as u is monotonic non-decreasing on $[a, b]$, we have

$$\begin{aligned}
|\mathcal{D}(f; u)| &= \left| \int_a^b \left[f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right] du(x) \right| \leq \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| du(x) \\
(4.2) \quad &= \frac{1}{b-a} \int_a^b \left| \int_a^b [f(x) - f(t)] dt \right| du(x)
\end{aligned}$$

As f is monotonic nondecreasing on $[a, x]$, and monotonic nondecreasing on $[x, b]$, we get

$$\begin{aligned}
\left| \int_a^b [f(x) - f(t)] dt \right| &\leq \int_a^b |f(x) - f(t)| dt \\
&\leq \int_a^x |f(x) - f(t)| dt + \int_x^b |f(x) - f(t)| dt \\
&= (x-a)f(x) - \int_a^x f(t) dt + \int_x^b f(t) dt - (b-x)f(x) \\
&= (2x-a-b)f(x) + \int_x^b f(t) dt - \int_a^x f(t) dt.
\end{aligned}$$

On utilizing the monotonicity property of f on both intervals, we have

$$\int_x^b f(t) dt \leq (b-x)f(b), \quad \text{and} \quad \int_a^x f(t) dt \geq (x-a)f(a),$$

which implies that,

$$(4.3) \quad \int_a^b |f(x) - f(t)| dt \leq (2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a).$$

Using (4.2) and (4.3), we get

(4.4)

$$|\mathcal{D}(f; u)| \leq \frac{1}{b-a} \int_a^b [(2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a)] du(x)$$

Now, using Riemann–Stieltjes integral we have

$$\begin{aligned} \int_a^b (2x-a-b)f(x) du(x) &= (b-a)[f(b)u(b) + f(a)u(a)] \\ &\quad - 2 \int_a^b u(x)f(x) dx - \int_a^b (2x-a-b)u(x) df(x) \end{aligned}$$

$$\int_a^b (b-x)f(b) du(x) = f(b) \int_a^b (b-x) du(x) = -(b-a)u(a)f(b) + f(b) \int_a^b u(x) dx$$

and

$$\int_a^b (x-a)f(a) du(x) = f(a) \int_a^b (x-a) du(x) = (b-a)u(b)f(a) - f(a) \int_a^b u(x) dx$$

Therefore, by (4.4), we get

(4.5)

$$\begin{aligned} &\int_a^b [(2x-a-b)f(x) + (b-x)f(b) - (x-a)f(a)] du(x) \\ &= (b-a)[f(b)u(b) + f(a)u(a)] - 2 \int_a^b u(x)f(x) dx - \int_a^b (2x-a-b)u(x) df(x) \\ &\quad - (b-a)u(a)f(b) + f(b) \int_a^b u(x) dx - (b-a)u(b)f(a) + f(a) \int_a^b u(x) dx \\ &= (b-a)(f(b) - f(a))(u(b) - u(a)) + (f(a) + f(b)) \int_a^b u(x) dx \\ &\quad - 2 \int_a^b u(x)f(x) dx - \int_a^b (2x-a-b)u(x) df(x). \end{aligned}$$

Now, by the monotonicity property of u we have $\int_a^b u(x) dx \leq (b-a)u(b)$,

$$\int_a^b u(x)f(x) dx \geq u(a) \int_a^b f(x) dx$$

and

$$\int_a^b (2x-a-b)u(x) df(x) \geq (a-b)u(a) \int_a^b df(x) = (a-b)u(a) \cdot (f(b) - f(a))$$

which gives by (4.5)

$$\begin{aligned}
& \int_a^b [(2x - a - b)f(x) + (b - x)f(b) - (x - a)f(a)] du(x) \\
&= (b - a)(f(b) - f(a))(u(b) - u(a)) + (f(a) + f(b)) \int_a^b u(x) dx \\
&\quad - 2 \int_a^b u(x)f(x) dx - \int_a^b (2x - a - b)u(x) df(x) \\
&\leq (b - a)(f(b) - f(a))(u(b) - u(a)) + (b - a)(f(a) + f(b))u(b) \\
&\quad - 2u(a) \int_a^b f(x) dx - (a - b)u(a) \cdot (f(b) - f(a)) \\
&= (b - a)[(f(b) - f(a))u(b) + (f(a) + f(b))u(b)] - 2u(a) \int_a^b f(x) dx \\
&= 2(b - a)f(b)u(b) - 2u(a) \int_a^b f(x) dx.
\end{aligned}$$

Therefore, by (4.4) we get

$$\begin{aligned}
|\mathcal{D}(f; u)| &\leq \frac{1}{b - a} \int_a^b [(2x - a - b)f(x) + (b - x)f(b) - (x - a)f(a)] du(x) \\
&\leq 2f(b)u(b) - 2\frac{u(a)}{b - a} \int_a^b f(x) dx,
\end{aligned}$$

Now, using the properties of ‘max’ function and the monotonicity of u , we get

$$\begin{aligned}
|\mathcal{D}(f; u)| &\leq 2f(b)u(b) - 2\frac{u(a)}{b - a} \int_a^b f(x) dx \\
&\leq 2 \cdot \max\{u(a), u(b)\} \cdot \left[f(b) - \frac{1}{b - a} \int_a^b f(x) dx \right] \\
&= 2u(b) \cdot \left[f(b) - \frac{1}{b - a} \int_a^b f(x) dx \right],
\end{aligned}$$

which proves the inequality (4.1). \square

5. A NUMERICAL QUADRATURE FORMULA FOR THE RIEMANN-STIELTJES INTEGRAL

In this section, an approximation for the Riemann–Stieltjes integral $\int_a^b f(x) du(x)$, in terms of the Riemann integral $\int_a^b f(t) dt$.

Theorem 6. *Let f, u be as in Theorem 3 and consider*

$$I_h := \{a = x_0 < x_1 < \dots < x_{n-1} < x_n = b\},$$

be a partition of $[a, b]$. Denote $h_i = x_{i+1} - x_i$, $i = 1, 2, \dots, n - 1$. Then we have

$$(5.1) \quad \int_a^b f(x) du(x) = A_n(f, u, I_h) + R_n(f, u, I_h)$$

where,

$$(5.2) \quad A_n(f, u, I_h) = \sum_{i=0}^{n-1} \frac{u(x_{i+1}) - u(x_i)}{h_i} \times \int_{x_i}^{x_{i+1}} f(t) dt$$

and the Remainder $R_n(f, u, I_h)$ satisfies the estimation

$$(5.3) \quad |R_n(f, u, I_h)| \leq [f(b) - f(a)] \cdot \bigvee_a^b(u).$$

Proof. Applying Theorem 3 on the intervals $[x_i, x_{i+1}]$, $i = 1, 2, \dots, n-1$, we get

$$\left| \int_{x_i}^{x_{i+1}} f(x) du(x) - \frac{u(x_{i+1}) - u(x_i)}{h_i} \int_{x_i}^{x_{i+1}} f(t) dt \right| \leq [f(x_{i+1}) - f(x_i)] \bigvee_{x_i}^{x_{i+1}}(u).$$

Summing the above inequality over i from 0 to $n-1$ and using the generalized triangle inequality, we deduce that

$$\begin{aligned} \left| \int_a^b f(x) du(x) - A_n(f, u, I_h) \right| &\leq \sum_{i=0}^{n-1} [f(x_{i+1}) - f(x_i)] \bigvee_{x_i}^{x_{i+1}}(u) \\ &\leq \max_{i=0, n-1} \{f(x_{i+1}) - f(x_i)\} \cdot \sum_{i=0}^{n-1} \bigvee_{x_i}^{x_{i+1}}(u) \\ &= [f(b) - f(a)] \cdot \bigvee_a^b(u), \end{aligned}$$

and the theorem is proved. \square

Theorem 7. Let f, u be as in Theorem 4. Let I_h as above. Then we have

$$(5.4) \quad \int_a^b f(x) du(x) = A_n(f, u, I_h) + R_n(f, u, I_h)$$

where, $A_n(f, u, I_h)$ is defined in (5.2) and the Remainder $R_n(f, u, I_h)$ satisfies the estimation

$$(5.5) \quad |R_n(f, u, I_h)| \leq L \left[\frac{1}{2} \nu(h) (f(b) - f(a)) + \int_a^b f(x) dx \right]$$

where, $\nu(h) = \max_{i=0, n-1} \{h_i\}$.

Proof. Applying Theorem 4 on the intervals $[x_i, x_{i+1}]$, $i = 1, 2, \dots, n-1$, we get

$$\begin{aligned} \left| \int_{x_i}^{x_{i+1}} f(x) du(x) - \frac{u(x_{i+1}) - u(x_i)}{h_i} \int_{x_i}^{x_{i+1}} f(t) dt \right| \\ \leq L \left[\frac{h_i}{2} (f(x_{i+1}) - f(x_i)) + \int_{x_i}^{x_{i+1}} f(x) dx \right]. \end{aligned}$$

Summing the above inequality over i from 0 to $n - 1$ and using the generalized triangle inequality, we deduce that

$$\begin{aligned} & \left| \int_a^b f(x) du(x) - A_n(f, u, I_h) \right| \\ & \leq L \sum_{i=0}^{n-1} \left[\frac{h_i}{2} (f(x_{i+1}) - f(x_i)) + \int_{x_i}^{x_{i+1}} f(x) dx \right] \\ & \leq L \left[\frac{1}{2} \max_{i=0, n-1} \{h_i\} \cdot \sum_{i=0}^{n-1} (f(x_{i+1}) - f(x_i)) + \int_a^b f(x) dx \right] \\ & \leq L \left[\frac{1}{2} \nu(h) (f(b) - f(a)) + \int_a^b f(x) dx \right], \end{aligned}$$

and the theorem is proved. \square

Remark 1. Similarly, one may apply Theorem 5, to approximate $\int_a^b f(x) du(x)$ in terms of $\int_a^b f(t) dt$. We shall omit the details to the interested reader.

REFERENCES

- [1] N.S. Barnett, S.S. Dragomir and I. Gomma, A companion for the Ostrowski and the generalised trapezoid inequalities, *Mathematical and Computer Modelling*, 50 (2009), 179–187.
- [2] N.S. Barnett, W.-S. Cheung, S.S. Dragomir, A. Sofo, Ostrowski and trapezoid type inequalities for the Stieltjes integral with Lipschitzian integrands or integrators, *Comp. Math. Appl.*, 57 (2009), 195–201.
- [3] P. Cerone, W.S. Cheung, S.S. Dragomir, On Ostrowski type inequalities for Stieltjes integrals with absolutely continuous integrands and integrators of bounded variation, *Comp. Math. Appl.*, 54 (2007), 183–191.
- [4] P. Cerone, S.S. Dragomir, New bounds for the three-point rule involving the Riemann-Stieltjes integrals, in: C. Gulati, et al. (Eds.), *Advances in Statistics Combinatorics and Related Areas*, World Science Publishing, 2002, pp. 53–62.
- [5] P. Cerone, S.S. Dragomir, Approximating the Riemann-Stieltjes integral via some moments of the integrand, *Mathematical and Computer Modelling*, 49 (2009), 242–248.
- [6] S.S. Dragomir and Th.M. Rassias (Ed.), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publishers, Dordrecht, 2002.
- [7] S.S. Dragomir, Inequalities of Grüss type for the Stieltjes integral and applications, *Kragujevac J. Math.*, 26 (2004) 89–112.
- [8] S.S. Dragomir, On the Ostrowski inequality for Riemann-Stieltjes integral $\int_a^b f(t) du(t)$ where f is of Hölder type and u is of bounded variation and applications, *J. KSIAM*, 5 (2001), 35–45.
- [9] S.S. Dragomir, On the Ostrowski's inequality for Riemann-Stieltjes integral and applications, *Korean J. Comput. & Appl. Math.*, 7 (2000), 611–627.
- [10] S.S. Dragomir, Some inequalities of midpoint and trapezoid type for the Riemann-Stieltjes integral, *Nonlinear Anal.* 47 (4) (2001) 2333–2340.
- [11] S.S. Dragomir, Approximating the Riemann-Stieltjes integral in terms of generalised trapezoidal rules, *Nonlinear Anal.* TMA 71 (2009) e62–e72.

- [12] S.S. Dragomir, Approximating the Riemann-Stieltjes integral by a trapezoidal quadrature rule with applications, *Mathematical and Computer Modelling* 54 (2011) 243–260.
- [13] S.S. Dragomir, C. Buşe, M.V. Boldea, L. Braescu, A generalisation of the trapezoid rule for the Riemann-Stieltjes integral and applications, *Nonlinear Anal. Forum* 6 (2) (2001) 33–351.
- [14] S.S. Dragomir, I. Fedotov, A Grüss type inequality for mappings of bounded variation and applications to numerical analysis, *Nonlinear Funct. Anal. Appl.*, 6 (3) (2001) 425–433.
- [15] S.S. Dragomir, I. Fedotov, An inequality of Grüss type for Riemann-Stieltjes integral and applications for special means, *Tamkang J. Math.*, 29 (4) (1998) 287–292
- [16] Z. Liu, Refinement of an inequality of Grüss type for Riemann-Stieltjes integral, *Soochow J. Math.*, 30 (4) (2004) 483–489.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, JERASH UNIVERSITY, 26150 JERASH,
JORDAN

E-mail address: mwomath@gmail.com