

**UPPER AND LOWER BOUNDS FOR THE p -ANGULAR
DISTANCE IN NORMED SPACES WITH APPLICATIONS**

S.S. DRAGOMIR^{1,2}

ABSTRACT. For nonzero vectors x and y in the normed linear space $(X, \|\cdot\|)$ we can define the p -angular distance by

$$\alpha_p [x, y] := \left\| \|x\|^{p-1} x - \|y\|^{p-1} y \right\|.$$

In this paper we show among others that

$$\begin{aligned} & \frac{1}{2} \left| \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \|x + y\| - \left(\|x\|^{p-1} + \|y\|^{p-1} \right) \|x - y\| \right| \\ & \leq \alpha_p [x, y] \\ & \leq \frac{1}{2} \left[\left| \|x\|^{p-1} - \|y\|^{p-1} \right| \|x + y\| + \left(\|x\|^{p-1} + \|y\|^{p-1} \right) \|x - y\| \right] \end{aligned}$$

for any $p \in \mathbb{R}$ and for any nonzero $x, y \in X$.

Some reverses of the triangle and the continuity of the norm inequalities are given as well.

Applications for functions f defined by power series in estimating the more general "distance" $\|f(\|x\|)x - f(\|y\|)y\|$ for certain $x, y \in X$ are also provided.

1. INTRODUCTION

Following [3, p. 403] or [12], for nonzero vectors x and y in the normed linear space $(X, \|\cdot\|)$ we define the *angular distance* $\alpha [x, y]$ between x and y by

$$\alpha [x, y] := \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\|.$$

In 1958, Massera and Schäffer [12, Lemma 5.1] showed that

$$(1.1) \quad \alpha [x, y] \leq \frac{2 \|x - y\|}{\max \{\|x\|, \|y\|\}},$$

which is better than the *Dunkl-Williams inequality* [6]

$$(1.2) \quad \alpha [x, y] \leq \frac{4 \|x - y\|}{\|x\| + \|y\|}.$$

We notice that the *Massera-Schäffer inequality* was rediscovered by Gurariĭ in [7] (see also [13, p. 516]).

In [10], Maligranda obtained the double inequality (see also [11]):

$$(1.3) \quad \frac{\|x - y\| - \left| \|x\| - \|y\| \right|}{\min \{\|x\|, \|y\|\}} \leq \alpha [x, y] \leq \frac{\|x - y\| + \left| \|x\| - \|y\| \right|}{\max \{\|x\|, \|y\|\}}.$$

The second inequality in (1.3) is better than Massera-Schäffer's inequality (1.1).

1991 *Mathematics Subject Classification.* Primary 46B20, 46B99; Secondary 51M16.

Key words and phrases. Angular distance, Normed Linear Spaces, Massera-Schäffer's inequality, Dunkl-Williams' inequality, Hile's inequality, Power series.

In the recent paper [10], L. Maligranda has also considered the *p*-angular distance

$$\alpha_p [x, y] := \left\| \|x\|^{p-1} x - \|y\|^{p-1} y \right\|$$

between the vectors x and y in the normed linear space $(X, \|\cdot\|)$ over the real or complex number field \mathbb{K} and showed that

$$(1.4) \quad \alpha_p [x, y] \leq \|x - y\| \times \begin{cases} (2-p) \cdot \frac{\max\{\|x\|^p, \|y\|^p\}}{\max\{\|x\|, \|y\|\}} & \text{if } p \in (-\infty, 0) \text{ and } x, y \neq 0; \\ (2-p) \cdot \frac{1}{[\max\{\|x\|, \|y\|\}]^{1-p}} & \text{if } p \in [0, 1] \text{ and } x, y \neq 0; \\ p \cdot [\max\{\|x\|, \|y\|\}]^{p-1} & \text{if } p \in (1, \infty). \end{cases}$$

The constants $2-p$ and p in (1.1) are best possible in the sense that they cannot be replaced by smaller quantities.

As pointed out in [10], the inequality (1.1) for $p \in [1, \infty)$ is better than the Bourbaki inequality obtained in 1965, [1, p. 257] (see also [13, p. 516]):

$$(1.5) \quad \alpha_p [x, y] \leq 3p \|x - y\| [\|x\| + \|y\|]^{p-1}, \quad x, y \in X.$$

The following results concerning upper bounds for the *p*-angular distance have been obtained by the author in [5]:

$$(1.6) \quad \alpha_p [x, y] \leq \begin{cases} \|x - y\| [\max\{\|x\|, \|y\|\}]^{p-1} + \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \min\{\|x\|, \|y\|\} & \text{if } p \in (1, \infty); \\ \frac{\|x - y\|}{[\min\{\|x\|, \|y\|\}]^{1-p}} + \left| \|x\|^{1-p} - \|y\|^{1-p} \right| \min\left\{ \frac{\|x\|^p}{\|y\|^{1-p}}, \frac{\|y\|^p}{\|x\|^{1-p}} \right\} & \text{if } p \in [0, 1]; \\ \frac{\|x - y\|}{[\min\{\|x\|, \|y\|\}]^{1-p}} + \frac{\left| \|x\|^{1-p} - \|y\|^{1-p} \right|}{\max\{\|x\|^{-p}\|y\|^{1-p}, \|y\|^{-p}\|x\|^{1-p}\}} & \text{if } p \in (-\infty, 0); \end{cases}$$

and

$$(1.7) \quad \alpha_p [x, y] \leq \begin{cases} \|x - y\| [\min\{\|x\|, \|y\|\}]^{p-1} + \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \max\{\|x\|, \|y\|\} & \text{if } p \in (1, \infty); \\ \frac{\|x - y\|}{[\max\{\|x\|, \|y\|\}]^{1-p}} + \left| \|x\|^{1-p} - \|y\|^{1-p} \right| \max\left\{ \frac{\|x\|^p}{\|y\|^{1-p}}, \frac{\|y\|^p}{\|x\|^{1-p}} \right\} & \text{if } p \in [0, 1]; \\ \frac{\|x - y\|}{[\max\{\|x\|, \|y\|\}]^{1-p}} + \frac{\left| \|x\|^{1-p} - \|y\|^{1-p} \right|}{\min\{\|x\|^{-p}\|y\|^{1-p}, \|y\|^{-p}\|x\|^{1-p}\}} & \text{if } p \in (-\infty, 0); \end{cases}$$

for any two nonzero vectors x, y in the normed linear space $(X, \|\cdot\|)$.

The upper bounds for $\alpha_p [x, y]$ provided by (1.4), (1.6) and (1.7) have been compared in [5] to conclude that some of the later ones are better in certain cases. The details are omitted here.

Finally, we recall the results of G. N. Hile from [4]:

$$(1.8) \quad \alpha_p [x, y] \leq \frac{\|x\|^p - \|y\|^p}{\|x\| - \|y\|} \cdot \|x - y\|,$$

for $p \in [1, \infty)$ and $x, y \in X$ with $\|x\| \neq \|y\|$, and

$$(1.9) \quad \alpha_{-p-1} [x, y] \leq \frac{\|x\|^p - \|y\|^p}{\|x\| - \|y\|} \cdot \frac{\|x - y\|}{\|x\|^p \|y\|^p},$$

for $p \in [1, \infty)$ and $x, y \in X \setminus \{0\}$ with $\|x\| \neq \|y\|$.

For other norm inequalities, see [8], [9] and [13].

In this paper we establish some new upper and lower bounds for the p -angular distance. Some reverses of the triangle and the continuity of the norm inequalities are given as well. Applications for functions f defined by power series in estimating the more general "distance" $\|f(\|x\|)x - f(\|y\|)y\|$ for certain $x, y \in X$ are also provided.

2. SOME PRELIMINARY RESULTS

For a pair of scalars $(\gamma, \beta) \in \mathbb{K}^2$ we can introduce the following (γ, β) -angular distance $d_{(\gamma, \beta)} : X^2 \rightarrow [0, \infty)$ given by

$$(2.1) \quad d_{(\gamma, \beta)} [x, y] := \|\gamma x - \beta y\|.$$

When $\gamma = \|x\|^{p-1}$ and $\beta = \|y\|^{p-1}$, then we have

$$d_{(\|x\|^{p-1}, \|y\|^{p-1})} [x, y] = \alpha_p [x, y]$$

for $p \in \mathbb{R}$.

In [5] we have shown that, for any $(\gamma, \beta) \in \mathbb{K}^2$ and $x, y \in X$ we have the following upper bounds for the (γ, β) -angular distance:

$$(2.2) \quad d_{(\gamma, \beta)} [x, y] \leq \|x - y\| \max\{|\gamma|, |\beta|\} + |\gamma - \beta| \min\{\|x\|, \|y\|\}$$

and

$$(2.3) \quad d_{(\gamma, \beta)} [x, y] \leq \|x - y\| \min\{|\gamma|, |\beta|\} + |\gamma - \beta| \max\{\|x\|, \|y\|\}$$

respectively.

By adding these two upper bounds we have the symmetrical bound

$$(2.4) \quad d_{(\gamma, \beta)} [x, y] \leq \|x - y\| \cdot \frac{|\gamma| + |\beta|}{2} + |\gamma - \beta| \cdot \frac{\|x\| + \|y\|}{2}.$$

We also proved in [5] that the following lower bounds may be provided as well:

$$(2.5) \quad \|x - y\| \min\{|\gamma|, |\beta|\} - |\gamma - \beta| \min\{\|x\|, \|y\|\} \leq d_{(\gamma, \beta)} [x, y]$$

and

$$(2.6) \quad \|x - y\| \max\{|\gamma|, |\beta|\} - |\gamma - \beta| \max\{\|x\|, \|y\|\} \leq d_{(\gamma, \beta)} [x, y],$$

and, by addition

$$\|x - y\| \cdot \frac{|\gamma| + |\beta|}{2} - |\gamma - \beta| \cdot \frac{\|x\| + \|y\|}{2} \leq d_{(\gamma, \beta)} [x, y].$$

We provide now some different upper and lower bounds for the (γ, β) -angular distance:

Lemma 1. For any $(\gamma, \beta) \in \mathbb{K}^2$ and $x, y \in X$ we have the inequalities

$$(2.7) \quad \begin{aligned} & \frac{1}{2} \|\gamma - \beta\| \|x + y\| - |\gamma + \beta| \|x - y\| \\ & \leq d_{(\gamma, \beta)} [x, y] \\ & \leq \frac{1}{2} [\|\gamma - \beta\| \|x + y\| + |\gamma + \beta| \|x - y\|]. \end{aligned}$$

Proof. We observe that for any $(\gamma, \beta) \in \mathbb{K}^2$ and $x, y \in X$ we have the key equality

$$\gamma x - \beta y = (\gamma - \beta) \frac{x + y}{2} + \frac{\gamma + \beta}{2} (x - y).$$

Utilising the triangle inequality we have

$$\begin{aligned} \|\gamma x - \beta y\| &= \left\| (\gamma - \beta) \frac{x + y}{2} + \frac{\gamma + \beta}{2} (x - y) \right\| \\ &\leq \left\| (\gamma - \beta) \frac{x + y}{2} \right\| + \left\| \frac{\gamma + \beta}{2} (x - y) \right\| \\ &= |\gamma - \beta| \left\| \frac{x + y}{2} \right\| + \left| \frac{\gamma + \beta}{2} \right| \|x - y\| \end{aligned}$$

and the second inequality in (2.7) is proved.

Utilising the continuity inequality of the norm, i.e., $\|u - v\| \geq \| \|u\| - \|v\| \|$ we also have

$$\begin{aligned} \|\gamma x - \beta y\| &= \left\| (\gamma - \beta) \frac{x + y}{2} - \frac{\gamma + \beta}{2} (y - x) \right\| \\ &\geq \left| \left\| (\gamma - \beta) \frac{x + y}{2} \right\| - \left\| \frac{\gamma + \beta}{2} (y - x) \right\| \right| \\ &= \left| |\gamma - \beta| \left\| \frac{x + y}{2} \right\| - \left| \frac{\gamma + \beta}{2} \right| \|x - y\| \right|, \end{aligned}$$

and the proof is complete. \square

Corollary 1. For any $(\gamma, \beta) \in \mathbb{K}^2$ and $x, y \in X$ we have the inequalities

$$\left| d_{(\gamma, \beta)} [x, y] - \frac{1}{2} |\gamma - \beta| \|x + y\| \right| \leq \frac{1}{2} |\gamma + \beta| \|x - y\|$$

and

$$\left| d_{(\gamma, \beta)} [x, y] - \frac{1}{2} |\gamma + \beta| \|x - y\| \right| \leq \frac{1}{2} |\gamma - \beta| \|x + y\|.$$

Remark 1. We observe that the upper bound for $d_{(\gamma, \beta)} [x, y]$ provided by (2.7) is better than the one provided by (2.4).

3. NEW BOUNDS FOR THE ANGULAR DISTANCE

We can state the following result providing upper and lower bounds for the angular distance:

Theorem 1. For any nonzero vectors x and y in the normed linear space $(X, \|\cdot\|)$ we have

$$\begin{aligned}
(3.1) \quad 0 &\leq \frac{1}{2\|y\|\|x\|} \max \{ \|x+y\| [\|x-y\| - \|\|x\| - \|y\|\|], \\
&\quad \|\|x\| - \|y\|\| [\|y\| + \|x\| - \|x+y\|] \} \\
&\leq \frac{1}{2\|y\|\|x\|} [(\|y\| + \|x\|) \|x-y\| - \|\|x\| - \|y\|\| \|x+y\|] \\
&\leq \alpha[x, y] \\
&\leq \frac{1}{2\|y\|\|x\|} [\|\|x\| - \|y\|\| \|x+y\| + (\|x\| + \|y\|) \|x-y\|] \\
&\leq \frac{1}{2\|y\|\|x\|} \min \{ (\|x\| + \|y\|) [\|\|x\| - \|y\|\| + \|x-y\|], \\
&\quad \|x-y\| [\|x+y\| + \|x\| + \|y\|] \}.
\end{aligned}$$

Proof. If we take $\gamma = \frac{1}{\|x\|}$ and $\beta = \frac{1}{\|y\|}$ (in 2.7) then we get

$$\begin{aligned}
&\frac{1}{2} \left| \left| \frac{1}{\|x\|} - \frac{1}{\|y\|} \right| \|x+y\| - \left| \frac{1}{\|x\|} + \frac{1}{\|y\|} \right| \|x-y\| \right| \\
&\leq \alpha[x, y] \\
&\leq \frac{1}{2} \left[\left| \frac{1}{\|x\|} - \frac{1}{\|y\|} \right| \|x+y\| + \left| \frac{1}{\|x\|} + \frac{1}{\|y\|} \right| \|x-y\| \right],
\end{aligned}$$

which is equivalent with

$$\begin{aligned}
&\frac{1}{2\|y\|\|x\|} [\|\|x\| - \|y\|\| \|x+y\| - (\|y\| + \|x\|) \|x-y\|] \\
&\leq \alpha[x, y] \\
&\leq \frac{1}{2\|x\|\|y\|} [\|\|x\| - \|y\|\| \|x+y\| + (\|x\| + \|y\|) \|x-y\|].
\end{aligned}$$

We notice that, by the triangle inequality and by the continuity of norm inequality we have

$$\begin{aligned}
&\|\|x\| - \|y\|\| \|x+y\| - (\|y\| + \|x\|) \|x-y\| \\
&= (\|y\| + \|x\|) \|x-y\| - \|\|x\| - \|y\|\| \|x+y\| \geq 0.
\end{aligned}$$

We also have

$$\begin{aligned}
&(\|y\| + \|x\|) \|x-y\| - \|\|x\| - \|y\|\| \|x+y\| \\
&\geq \|x+y\| [\|x-y\| - \|\|x\| - \|y\|\|] \geq 0
\end{aligned}$$

and

$$\begin{aligned}
&(\|y\| + \|x\|) \|x-y\| - \|\|x\| - \|y\|\| \|x+y\| \\
&\geq \|\|x\| - \|y\|\| [\|y\| + \|x\| - \|x+y\|] \geq 0
\end{aligned}$$

which implies that

$$\begin{aligned} & (\|y\| + \|x\|) \|x - y\| - \|x\| - \|y\| \|x + y\| \\ & \geq \max \{ \|x + y\| [\|x - y\| - \|x\| - \|y\|], \\ & \|x\| - \|y\| [\|y\| + \|x\| - \|x + y\|] \} \\ & \geq 0. \end{aligned}$$

These prove the first three inequalities in (3.1).

We also have that

$$\begin{aligned} & \|x\| - \|y\| \|x + y\| + (\|x\| + \|y\|) \|x - y\| \\ & \leq (\|x\| + \|y\|) [\|x\| - \|y\| + \|x - y\|] \end{aligned}$$

and

$$\begin{aligned} & \|x\| - \|y\| \|x + y\| + (\|x\| + \|y\|) \|x - y\| \\ & \leq \|x - y\| [\|x + y\| + \|x\| + \|y\|] \end{aligned}$$

which implies that

$$\begin{aligned} & \|x\| - \|y\| \|x + y\| + (\|x\| + \|y\|) \|x - y\| \\ & \leq \min \{ (\|x\| + \|y\|) [\|x\| - \|y\| + \|x - y\|], \\ & \|x - y\| [\|x + y\| + \|x\| + \|y\|] \} \end{aligned}$$

and the last part of (3.1) is proved. \square

Problem 1. *It is an open question for the author whether or not the upper bound for the angular distance provided by the inequality*

$$\alpha[x, y] \leq \frac{1}{2 \frac{\|y\|}{\|x\|}} [\|x\| - \|y\| \|x + y\| + (\|x\| + \|y\|) \|x - y\|]$$

is better than the one in (1.3). The same question applies for the lower bound.

4. BOUNDS FOR THE 2-ANGULAR DISTANCE

The case $p = 2$ in the p -angular distance is of interest. It generates the 2-angular distance which has the expression

$$(4.1) \quad \alpha_2[x, y] := \| \|x\| x - \|y\| y \|$$

and can be defined for any x, y in a normed linear space.

We observe that from Maligranda's result (1.4) we get the upper bound

$$(4.2) \quad \alpha_2[x, y] \leq 2 \max \{ \|x\|, \|y\| \} \|x - y\|$$

while from Hile's inequality we have

$$(4.3) \quad \alpha_2[x, y] \leq (\|x\| + \|y\|) \|x - y\|,$$

for any vectors x and y in the normed linear space $(X, \|\cdot\|)$.

Since

$$\frac{\|x\| + \|y\|}{2} \leq \max \{ \|x\|, \|y\| \}$$

then (4.3) is a better inequality than (4.2).

Moreover, if we employ (2.2) and (2.3) for $\gamma = \|x\|$ and $\beta = \|y\|$, then we get

$$\alpha_2[x, y] \leq \|x - y\| \max \{ \|x\|, \|y\| \} + \| \|x\| - \|y\| \| \min \{ \|x\|, \|y\| \}$$

and

$$\alpha_2 [x, y] \leq \|x - y\| \min \{\|x\|, \|y\|\} + \left| \|x\| - \|y\| \right| \max \{\|x\|, \|y\|\}$$

implying that

$$(4.4) \quad \alpha_2 [x, y] \leq \min \{ \|x - y\| \max \{\|x\|, \|y\|\} + \left| \|x\| - \|y\| \right| \min \{\|x\|, \|y\|\} \\ \|x - y\| \min \{\|x\|, \|y\|\} + \left| \|x\| - \|y\| \right| \max \{\|x\|, \|y\|\} \}$$

for any vectors x and y in the normed linear space $(X, \|\cdot\|)$.

Since

$$\begin{aligned} & \|x - y\| \max \{\|x\|, \|y\|\} + \left| \|x\| - \|y\| \right| \min \{\|x\|, \|y\|\} \\ & \leq \|x - y\| [\max \{\|x\|, \|y\|\} + \min \{\|x\|, \|y\|\}] \\ & = \|x - y\| (\|x\| + \|y\|) \end{aligned}$$

we can conclude that (4.4) is better than Hile's inequality (4.3).

Now, utilizing (2.5) and (2.6) we also have the lower bounds

$$(4.5) \quad \min \{\|x\|, \|y\|\} [\|x - y\| - \left| \|x\| - \|y\| \right|] \leq \alpha_2 [x, y]$$

and

$$(4.6) \quad \max \{\|x\|, \|y\|\} [\|x - y\| - \left| \|x\| - \|y\| \right|] \leq \alpha_2 [x, y].$$

Obviously (4.6) is better than (4.5) and produces the following reverse of continuity of norm inequality

$$(4.7) \quad 0 \leq \|x - y\| - \left| \|x\| - \|y\| \right| \leq \frac{1}{\max \{\|x\|, \|y\|\}} \alpha_2 [x, y]$$

that holds for any nonzero vectors x and y in the normed linear space $(X, \|\cdot\|)$.

Theorem 2. For any vectors x and y in the normed linear space $(X, \|\cdot\|)$ we have

$$(4.8) \quad \begin{aligned} 0 & \leq \max \{ \|x + y\| [\|x - y\| - \left| \|x\| - \|y\| \right|], \\ & \quad \| \|x\| - \|y\| \| [\|y\| + \|x\| - \|x + y\|] \} \\ & \leq (\|y\| + \|x\|) \|x - y\| - \left| \|x\| - \|y\| \right| \|x + y\| \\ & \leq \alpha_2 [x, y] \\ & \leq \| \|x\| - \|y\| \| \|x + y\| + (\|x\| + \|y\|) \|x - y\| \\ & \leq \min \{ (\|x\| + \|y\|) [\| \|x\| - \|y\| \| + \|x - y\|], \\ & \quad \|x - y\| [\|x + y\| + \|x\| + \|y\|] \}. \end{aligned}$$

Proof. If we take $\gamma = \|x\|$ and $\beta = \|y\|$ in the inequality (2.7) then we get

$$\begin{aligned} & \frac{1}{2} [\| \|x\| - \|y\| \| \|x + y\| - (\|x\| + \|y\|) \|x - y\|] \\ & \leq \alpha_2 [x, y] \\ & \leq \frac{1}{2} [\| \|x\| - \|y\| \| \|x + y\| + (\|x\| + \|y\|) \|x - y\|], \end{aligned}$$

which is equivalent with (see the proof of the above Theorem 1)

$$\begin{aligned} & \frac{1}{2} [(\|x\| + \|y\|) \|x - y\| - \| \|x\| - \|y\| \| \|x + y\|] \\ & \leq \alpha_2 [x, y] \\ & \leq \frac{1}{2} [\| \|x\| - \|y\| \| \|x + y\| + (\|x\| + \|y\|) \|x - y\|]. \end{aligned}$$

The rest follows as in the proof of the above Theorem 1 and the details are omitted. \square

Remark 2. From the inequality (4.8) we get the following reverse for the triangle inequality

$$(4.9) \quad 0 \leq \|y\| + \|x\| - \|x + y\| \leq \frac{1}{\left| \|x\| - \|y\| \right|} \alpha_2 [x, y]$$

that holds for any vectors x and y in the normed linear space $(X, \|\cdot\|)$ with $\|x\| \neq \|y\|$.

From the inequality (4.8) we also have the following reverse of continuity of norm inequality

$$(4.10) \quad 0 \leq \|x - y\| - \left| \|x\| - \|y\| \right| \leq \frac{1}{\|x + y\|} \alpha_2 [x, y]$$

that holds for any vectors x and y in the normed linear space $(X, \|\cdot\|)$ with $x \neq -y$.

Since, in general, the quantity $\|x + y\|$ is not always greater than $\max \{\|x\|, \|y\|\}$ we can conclude that not always the inequality (4.10) is better than the previous inequality from (4.7).

5. BOUNDS FOR THE p -ANGULAR DISTANCE

If we write the Lemma 1 for $\gamma = \|x\|^{p-1}$ and $\beta = \|y\|^{p-1}$ we can state the following result for the p -angular distance:

Theorem 3. For any nonzero vectors x and y in the normed linear space $(X, \|\cdot\|)$ we have

$$(5.1) \quad \begin{aligned} & \frac{1}{2} \left| \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \|x + y\| - \left(\|x\|^{p-1} + \|y\|^{p-1} \right) \|x - y\| \right| \\ & \leq \alpha_p [x, y] \\ & \leq \frac{1}{2} \left[\left| \|x\|^{p-1} - \|y\|^{p-1} \right| \|x + y\| + \left(\|x\|^{p-1} + \|y\|^{p-1} \right) \|x - y\| \right] \end{aligned}$$

for any $p \in \mathbb{R}$.

Now, for $s \in [-\infty, \infty]$ and $a, b > 0, a \neq b$, by following [2, p. 385], we can introduce the s -generalized logarithmic means by

$$L^{[s]}(a, b) := \begin{cases} \left(\frac{b^{s+1} - a^{s+1}}{(s+1)(b-a)} \right)^{1/s} & \text{if } s \neq -1, 0, \pm\infty; \\ \frac{b-a}{\ln b - \ln a} & \text{if } s = -1; \\ \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{1/(b-a)} & \text{if } s = 0; \\ \max \{a, b\} & \text{if } s = \infty; \\ \min \{a, b\} & \text{if } s = -\infty. \end{cases}$$

The mapping $\mathbb{R} \ni s \rightarrow L^{[s]}(a, b)$ is strictly increasing and (see [2, p. 386])

$$(5.2) \quad \min \{a, b\} < L^{[s]}(a, b) < \max \{a, b\}$$

for any $s \in \mathbb{R}$ and $a, b > 0$, with $a \neq b$.

Utilising the properties of the s -generalized logarithmic means we proved the following lemma in [5]:

Lemma 2. *For any two nonzero vectors $x, y \in X$ we have*

$$(5.3) \quad (p-1) [\min \{\|x\|, \|y\|\}]^{p-2} \|x\| - \|y\| \\ \leq \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \leq (p-1) \|x\| - \|y\| [\max \{\|x\|, \|y\|\}]^{p-2}$$

if $p \in (2, \infty)$,

$$(5.4) \quad (p-1) \frac{1}{[\max \{\|x\|, \|y\|\}]^{2-p}} \|x\| - \|y\| \\ \leq \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \leq (p-1) \|x\| - \|y\| \frac{1}{[\min \{\|x\|, \|y\|\}]^{2-p}}$$

if $p \in [1, 2]$, and

$$(5.5) \quad (1-p) \frac{\|x\|^{1-p} \|y\|^{1-p}}{[\max \{\|x\|, \|y\|\}]^{2-p}} \|x\| - \|y\| \\ \leq \left| \|x\|^{1-p} - \|y\|^{1-p} \right| \leq (1-p) \|x\| - \|y\| \frac{\|x\|^{1-p} \|y\|^{1-p}}{[\min \{\|x\|, \|y\|\}]^{2-p}}$$

if $p \in (-\infty, 1)$, respectively.

By utilizing this lemma and Theorem 3 we can state the following sequence of upper bounds for the p -angular distance:

$$(5.6) \quad \alpha_p [x, y] \\ \leq \frac{1}{2} \left[\left| \|x\|^{p-1} - \|y\|^{p-1} \right| \|x + y\| + (\|x\|^{p-1} + \|y\|^{p-1}) \|x - y\| \right] \\ \leq (p-1) \|x\| - \|y\| [\max \{\|x\|, \|y\|\}]^{p-2} \left\| \frac{x+y}{2} \right\| \\ + \frac{\|x\|^{p-1} + \|y\|^{p-1}}{2} \|x - y\| \\ \leq [\max \{\|x\|, \|y\|\}]^{p-1} [(p-1) \|x\| - \|y\| + \|x - y\|] \\ \leq p [\max \{\|x\|, \|y\|\}]^{p-1} \|x - y\|$$

for $p \geq 1$, which is a better inequality than Maligranda's result (1.4).

Similar results can be stated for $p < 1$, however the details are not presented here.

6. APPLICATIONS FOR POWER SERIES

For power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ with complex coefficients we can naturally construct another power series which have as coefficients the absolute values of the coefficients of the original series, namely, $f_a(z) := \sum_{n=0}^{\infty} |a_n| z^n$. It is obvious that this new power series have the same radius of convergence as the original series, and that if all coefficients $a_n \geq 0$, then $f_a = f$.

As some natural examples that are useful for applications, we can point out that, if

$$(6.1) \quad \begin{aligned} f(z) &= \sum_{n=1}^{\infty} \frac{(-1)^n}{n} z^n = \ln \frac{1}{1+z}, \quad z \in D(0,1); \\ g(z) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} z^{2n} = \cos z, \quad z \in \mathbb{C}; \\ h(z) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1} = \sin z, \quad z \in \mathbb{C}; \\ l(z) &= \sum_{n=0}^{\infty} (-1)^n z^n = \frac{1}{1+z}, \quad z \in D(0,1); \end{aligned}$$

then the corresponding functions constructed by the use of the absolute values of the coefficients are

$$(6.2) \quad \begin{aligned} f_a(z) &= \sum_{n=1}^{\infty} \frac{1}{n} z^n = \ln \frac{1}{1-z}, \quad z \in D(0,1); \\ g_a(z) &= \sum_{n=0}^{\infty} \frac{1}{(2n)!} z^{2n} = \cosh z, \quad z \in \mathbb{C}; \\ h_A(z) &= \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} z^{2n+1} = \sinh z, \quad z \in \mathbb{C}; \\ l_A(z) &= \sum_{n=0}^{\infty} z^n = \frac{1}{1-z}, \quad z \in D(0,1). \end{aligned}$$

Other important examples of functions as power series representations with non-negative coefficients are:

$$(6.3) \quad \begin{aligned} \exp(z) &= \sum_{n=0}^{\infty} \frac{1}{n!} z^n, \quad z \in \mathbb{C}, \\ \frac{1}{2} \ln \left(\frac{1+z}{1-z} \right) &= \sum_{n=1}^{\infty} \frac{1}{2n-1} z^{2n-1}, \quad z \in D(0,1); \\ \sin^{-1}(z) &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\frac{1}{2})}{\sqrt{\pi}(2n+1)n!} z^{2n+1}, \quad z \in D(0,1); \\ \tanh^{-1}(z) &= \sum_{n=1}^{\infty} \frac{1}{2n-1} z^{2n-1}, \quad z \in D(0,1) \\ {}_2F_1(\alpha, \beta, \gamma, z) &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)\Gamma(n+\beta)\Gamma(\gamma)}{n!\Gamma(\alpha)\Gamma(\beta)\Gamma(n+\gamma)} z^n, \quad \alpha, \beta, \gamma > 0, \\ &z \in D(0,1); \end{aligned}$$

where Γ is *Gamma function*.

Theorem 4. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a function defined by power series with complex coefficients and convergent on the open disk $D(0, R) \subset \mathbb{C}$, $R > 0$. If

$(X; \|\cdot\|)$ is a normed linear space and $x, y \in X$ with $\|x\|, \|y\| < R$, then

$$\begin{aligned}
(6.4) \quad \|f(\|x\|)x - f(\|y\|)y\| &\leq \left\| \frac{x+y}{2} \right\| \|\|x\| - \|y\|\| \max\{f'_a(\|x\|), f'_a(\|y\|)\} \\
&\quad + \|x-y\| \frac{f_a(\|x\|) + f_a(\|y\|)}{2} \\
&\leq \|x-y\| \left\{ \left\| \frac{x+y}{2} \right\| \max\{f'_a(\|x\|), f'_a(\|y\|)\} \right. \\
&\quad \left. + \frac{f_a(\|x\|) + f_a(\|y\|)}{2} \right\} \\
&\leq \|x-y\| [\max\{\|x\|, \|y\|\} \max\{f'_a(\|x\|), f'_a(\|y\|)\} \\
&\quad + \max\{f_a(\|x\|), f_a(\|y\|)\}].
\end{aligned}$$

Proof. If we write the inequality (5.6) for $p = n + 1$, n a natural number, then we get

$$\begin{aligned}
(6.5) \quad &\|\|x\|^n x - \|y\|^n y\| \\
&\leq n [\max\{\|x\|, \|y\|\}]^{n-1} \left\| \frac{x+y}{2} \right\| \|\|x\| - \|y\|\| \\
&\quad + \frac{\|x\|^n + \|y\|^n}{2} \|x-y\|
\end{aligned}$$

for any $x, y \in X$.

Let $m \geq 1$. Then we have, by the generalized triangle inequality and by (6.5), that

$$\begin{aligned}
(6.6) \quad &\left\| \left(\sum_{n=0}^m a_n \|x\|^n \right) x - \left(\sum_{n=0}^m a_n \|y\|^n \right) y \right\| \\
&\leq \sum_{n=0}^m |a_n| \|\|x\|^n x - \|y\|^n y\| \\
&\leq \left\| \frac{x+y}{2} \right\| \|\|x\| - \|y\|\| \sum_{n=0}^m n |a_n| [\max\{\|x\|, \|y\|\}]^{n-1} \\
&\quad + \|x-y\| \sum_{n=0}^m |a_n| \frac{\|x\|^n + \|y\|^n}{2}.
\end{aligned}$$

Since $\|x\|, \|y\| < R$, the series

$$\sum_{n=0}^{\infty} a_n \|x\|^n, \sum_{n=0}^{\infty} a_n \|y\|^n, \sum_{n=0}^{\infty} |a_n| \|x\|^n, \sum_{n=0}^{\infty} |a_n| \|y\|^n$$

and

$$\sum_{n=0}^{\infty} n |a_n| [\max\{\|x\|, \|y\|\}]^{n-1}$$

are convergent and

$$\sum_{n=0}^{\infty} a_n \|x\|^n = f(\|x\|), \sum_{n=0}^{\infty} a_n \|y\|^n = f(\|y\|),$$

$$\sum_{n=0}^{\infty} |a_n| \|x\|^n = f_a(\|x\|), \quad \sum_{n=0}^{\infty} |a_n| \|y\|^n = f_a(\|y\|)$$

while

$$\begin{aligned} \sum_{n=0}^{\infty} n |a_n| [\max\{\|x\|, \|y\|\}]^{n-1} &= f'_a(\max\{\|x\|, \|y\|\}) \\ &= \max\{f'_a(\|x\|), f'_a(\|y\|)\}. \end{aligned}$$

Taking the limit over $m \rightarrow \infty$ in (6.6) we obtain the first part of (6.4).

The second part is obvious. \square

Remark 3. If we take $f(z) := \exp(z) = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$ then we have from (6.4) the following inequality

$$\begin{aligned} (6.7) \quad & \|\exp(\|x\|)x - \exp(\|y\|)y\| \\ & \leq \left\| \frac{x+y}{2} \right\| \| \|x\| - \|y\| \| \max\{\exp\|x\|, \exp\|y\|\} \\ & + \|x-y\| \frac{\exp\|x\| + \exp\|y\|}{2} \\ & \leq \|x-y\| \left\{ \left\| \frac{x+y}{2} \right\| \max\{\exp\|x\|, \exp\|y\|\} \right. \\ & \quad \left. + \frac{\exp\|x\| + \exp\|y\|}{2} \right\} \\ & \leq \|x-y\| \max\{\exp\|x\|, \exp\|y\|\} (\max\{\|x\|, \|y\|\} + 1) \end{aligned}$$

for any $x, y \in X$.

If we apply the inequality (6.4) for the functions $f(z) := \frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$ and $f(z) := \frac{1}{1+z} = \sum_{n=0}^{\infty} (-1)^n z^n$ then we have

$$\begin{aligned} (6.8) \quad & \left\| \frac{x}{1 \pm \|x\|} - \frac{y}{1 \pm \|y\|} \right\| \\ & \leq \left\| \frac{x+y}{2} \right\| \| \|x\| - \|y\| \| \max\{(1 - \|x\|)^{-2}, (1 - \|y\|)^{-2}\} \\ & + \|x-y\| \frac{(1 - \|x\|)^{-1} + (1 - \|y\|)^{-1}}{2} \\ & \leq \|x-y\| \left\{ \left\| \frac{x+y}{2} \right\| \max\{(1 - \|x\|)^{-2}, (1 - \|y\|)^{-2}\} \right. \\ & \quad \left. + \frac{(1 - \|x\|)^{-1} + (1 - \|y\|)^{-1}}{2} \right\} \\ & \leq \|x-y\| \left[\max\{\|x\|, \|y\|\} \max\{(1 - \|x\|)^{-2}, (1 - \|y\|)^{-2}\} \right. \\ & \quad \left. + \max\{(1 - \|x\|)^{-1}, (1 - \|y\|)^{-1}\} \right] \end{aligned}$$

for any $x, y \in X$ with $\|x\|, \|y\| < 1$.

The interested reader may apply the above results for other power series as pointed out in (6.1)-(6.3). However the details are not provided here.

REFERENCES

- [1] N. Bourbaki, *Integration*, Herman, Paris, 1965.
- [2] P. S. Bullen, *Handbook of Means and Their Inequalities*, Kluwer Academic Publishers, Dordrecht/Boston/London, 2003.
- [3] J. A. Clarkson, Uniform convex spaces, *Trans. Amer. Math. Soc.* **40**(1936), 396-414.
- [4] G. N. Hile, Entire solutions of linear elliptic equations with Laplacian principal part, *Pacific J. Math.* **62**(1976), 124-140.
- [5] S. S. Dragomir, Inequalities for the p -angular distance in normed linear spaces. *Math. Inequal. Appl.* **12** (2009), no. 2, 391–401.
- [6] C. F. Dunkl and K. S. Williams, A simple norm inequality, *Amer. Math. Month.* **71**(1964), 53-54.
- [7] V. I. Gurariĭ, Strengthening the Dunkl-Williams inequality on the norms of elements of Banach spaces (Ukrainian), *Dopovidi Akad. Nauk Ukrain. RSR* **1966**(1966), 35-38.
- [8] Eder Kikianty and S. S. Dragomir, Hermite-Hadamard's inequality and the p -HH-norm on the Cartesian product of two copies of a normed space. *Math. Inequal. Appl.* **13** (2010), no. 1, 1–32.
- [9] Eder Kikianty, G. Sinnamon, The p -HH norms on Cartesian powers and sequence spaces. *J. Math. Anal. Appl.* **359** (2009), no. 2, 765–779.
- [10] L. Maligranda, Simple norm inequalities, *Amer. Math. Month.* **113**(2006), 256-260.
- [11] L. Maligranda, Some remarks on the triangle inequality for norms, *Banach J. Math. Anal.* **2** (2008), no. 2, 31–41.
- [12] J. L. Massera and J. J. Schäffer, Linear differential equations and functional analysis. I., *Ann. of Math.* **67**(1958), 517-573.
- [13] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Dordrecht, 1993. Kluwer

¹MATHEMATICS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.org/dragomir>

²SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA.