

Received 19/02/13

**SOME HERMITE-HADAMARD TYPE INEQUALITIES FOR
FUNCTIONS WHOSE PARTIAL DERIVATIVES IN ABSOLUTE
VALUE ARE PREINTEX ON THE CO-ORDINATES**

M. A. LATIF

ABSTRACT. In this paper we point out some inequalities of Hermite-Hadamard type for double integrals of functions whose partial derivatives in absolute value are preinvex on the co-ordinates on rectangle from the plane. Our established results generalize those results proved in [33] for functions whose partial derivatives in absolute value are convex on the co-ordinates on the rectangle from the plane.

1. INTRODUCTION

The following definition is well known in literature:

A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if the inequality

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y),$$

holds for all $x, y \in I$ and $\lambda \in [0, 1]$.

Many important inequalities have been established for the class of convex functions but the most famous is the Hermite-Hadamard's inequality. This double inequality is stated as:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}, \quad (1.1)$$

where $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$ a convex function, $a, b \in I$ with $a < b$. The inequalities in (1.1) are in reversed order if f a concave function.

The inequalities (1.1) have become an important cornerstone in mathematical analysis and optimization and many uses of these inequalities have been discovered in a variety of settings. The Hermite-Hadamard inequality (1.1) has been extended, refined and generalized in a number of ways, see for instance [6, 7, 9, 20, 24, 29, 32, 34] and the references therein.

In recent years, lot of efforts have been made by many mathematicians to generalize the classical convexity. These studies include among others the work of Hanson in [?], Ben-Israel and Mond [6], Pini [26], M.A.Noor [21, 22], Yang and Li [35] and Weir [34]. Mond [6], Weir [34] and Noor [21, 22], have studied the basic properties of the preinvex functions and their role in optimization, variational inequalities and equilibrium problems. Hanson in [?], introduced invex functions as a significant generalization of convex functions. Ben-Israel and Mond [6], gave the

Date: January 25, 2013.

2000 Mathematics Subject Classification. Primary: 26A15, 26A51; Scodary 52A30.

Key words and phrases. convex function, Hermite-Hadamard type inequality, preinvex function on the co-ordinaes, Hölder's integral integral inequality, power-mean integral inequality.

This paper is in final form and no version of it will be submitted for publication elsewhere.

concept of preinvex function which is special case of invexity. Pini [26], introduced the concept of prequasiinvex functions as a generalization of invex functions.

Let us recall some known results concerning invexity and preinvexity

Let K be a closed set in \mathbb{R}^n and let $f : K \rightarrow \mathbb{R}$ and $\eta : K \times K \rightarrow \mathbb{R}$ be continuous functions. Let $x \in K$, then the set K is said to be invex at x with respect to $\eta(\cdot, \cdot)$, if

$$x + t\eta(y, x) \in K, \forall x, y \in K, t \in [0, 1].$$

K is said to be an invex set with respect to η if K is invex at each $x \in K$. The invex set K is also called a η -connected set.

Definition 1. [34] *The function f on the invex set K is said to be preinvex with respect to η , if*

$$f(u + t\eta(v, u)) \leq (1 - t)f(u) + tf(v), \forall u, v \in K, t \in [0, 1].$$

The function f is said to be preconcave if and only if $-f$ is preinvex.

It is to be noted that every convex function is preinvex with respect to the map $\eta(x, y) = x - y$ but the converse is not true see for instance [34].

In the recent paper, Noor [20] has obtained the following Hermite-Hadamard inequalities for the preinvex functions:

Theorem 1. [20] *Let $f : [a, a + \eta(b, a)] \rightarrow (0, \infty)$ be a preinvex function on the interval of the real numbers K° (the interior of K) and $a, b \in K^\circ$ with $a < a + \eta(b, a)$. Then the following inequality holds:*

$$f\left(\frac{2a + \eta(b, a)}{2}\right) \leq \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} f(x) dx \leq \frac{f(a) + f(b)}{2}. \quad (1.2)$$

Barani, Ghazanfari and Dragomir in [2], presented the following estimates of the right-side of a Hermite-Hadamard type inequality in which some preinvex functions are involved.

Theorem 2. [2] *Let $K \subseteq \mathbb{R}$ be an open invex subset with respect to $\eta : K \times K \rightarrow \mathbb{R}$. Suppose that $f : K \rightarrow \mathbb{R}$ is a differentiable function. Assume $p \in \mathbb{R}$ with $p > 1$. If $|f'|^{p-1}$ is preinvex on K then, for every $a, b \in K$ with $\eta(b, a) \neq 0$, then the following inequality holds:*

$$\left| \frac{f(a) + f(a + \eta(b, a))}{2} - \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} f(x) dx \right| \leq \frac{\eta(b, a)}{2(1+p)^{\frac{1}{p}}} \left[\frac{|f'(a)|^{\frac{p}{p-1}} + |f'(b)|^{\frac{p}{p-1}}}{2} \right]^{\frac{p-1}{p}}. \quad (1.3)$$

Theorem 3. [2] *Let $K \subseteq \mathbb{R}$ be an open invex subset with respect to $\eta : K \times K \rightarrow \mathbb{R}$. Suppose that $f : K \rightarrow \mathbb{R}$ is a differentiable function. If $|f'|$ is preinvex on K then,*

for every $a, b \in K$ with $\eta(b, a) \neq 0$, then the following inequality holds:

$$\left| \frac{f(a) + f(a + \eta(b, a))}{2} - \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} f(x) dx \right| \leq \frac{\eta(b, a)}{8} \left(|f'(a)| + |f'(b)| \right). \quad (1.4)$$

For more recent results on Hermite-Hadamard type inequalities for preinvex, log-preinvex functions, we refer the readers to the latest papers of M. Z. Sarikaya et. al, [31].

Let us consider now a bidimensional interval $\Delta =: [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$ and $c < d$. A mapping $f : \Delta \rightarrow \mathbb{R}$ is said to be convex on Δ if the inequality

$$f(\lambda x + (1 - \lambda)z, \lambda y + (1 - \lambda)w) \leq \lambda f(x, y) + (1 - \lambda)f(z, w),$$

holds for all $(x, y), (z, w) \in \Delta$ and $\lambda \in [0, 1]$.

A modification for convex functions on Δ , known as co-ordinated convex functions, was introduced by S. S. Dragomir [10] as follows:

A function $f : \Delta \rightarrow \mathbb{R}$ is said to be convex on the co-ordinates on Δ if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}$, $f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}$, $f_x(v) = f(x, v)$ are convex where defined for all $x \in [a, b]$, $y \in [c, d]$.

A formal definition for co-ordinated convex functions may be stated as follows:

Definition 2. [15] *A function $f : \Delta \rightarrow \mathbb{R}$ is said to be convex on the co-ordinates on Δ if the inequality*

$$\begin{aligned} & f(tx + (1 - t)y, su + (1 - s)w) \\ & \leq tsf(x, u) + t(1 - s)f(x, w) + s(1 - t)f(y, u) + (1 - t)(1 - s)f(y, w), \end{aligned}$$

holds for all $t, s \in [0, 1]$ and $(x, u), (y, w) \in \Delta$.

Clearly, every convex mapping $f : \Delta \rightarrow \mathbb{R}$ is convex on the co-ordinates but converse may not be true [10].

The following Hermite-Hadamard type for co-ordinated convex functions on the rectangle from the plane \mathbb{R}^2 were established in ??:

Theorem 4. [10] *Suppose that $f : \Delta \rightarrow \mathbb{R}$ is co-ordinated convex on Δ , then*

$$\begin{aligned} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) & \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right] \\ & \leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\ & \leq \frac{1}{4} \left[\frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) + f(b, y)] dy \right] \\ & \leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}. \quad (1.5) \end{aligned}$$

The above inequalities are sharp.

For several recent results on Hermite-Hadamard type inequalities for functions that satisfy different kinds of convexity on the co-ordinates on the rectangle from the plane \mathbb{R}^2 we refer the reader to [1]-[4], [10]-[11], [15]-[17], [25]-[28] and [33].

By using the following lemma:

Lemma 1. [21, Lemma 1] *Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$, $c < d$. If $\frac{\partial^2 f}{\partial t \partial s} \in L(\Delta)$, then the following equality holds:*

$$\begin{aligned} & \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\ & - \frac{1}{2} \left[\frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) dy + f(b, y)] dy \right] \\ & = \frac{(b-a)(d-c)}{4} \int_0^1 \int_0^1 (1-2t)(1-2s) \frac{\partial^2 f(ta + (1-t)b, sc + (1-s)d)}{\partial t \partial s} dt ds. \end{aligned} \quad (1.6)$$

Sarikaya, et. al [33], proved the following Hermite-Hadamard type inequalities for differentiable co-ordinated convex functions:

Theorem 5. [21, Theorem 2, Page 4] *Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$, $c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$ is convex on the co-ordinates on Δ , then one has the inequalities:*

$$\begin{aligned} & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\ & \leq \frac{(b-a)(d-c)}{16} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|}{4} \right), \end{aligned} \quad (1.7)$$

where

$$A = \frac{1}{2} \left[\frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) dy + f(b, y)] dy \right].$$

Theorem 6. [33, Theorem 3, Page 6-7] *Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$, $c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$, $q > 1$, is convex on the co-ordinates on Δ , then one has the inequalities:*

$$\begin{aligned} & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\ & \leq \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|}{4} \right)^{\frac{1}{q}}, \end{aligned} \quad (1.8)$$

where

$$A = \frac{1}{2} \left[\frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) dy + f(b, y)] dy \right]$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 7. [33, Theorem 4, Page 8-9] *Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b, c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q, q \geq 1$, is convex on the co-ordinates on Δ , then one has the inequalities:*

$$\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \leq \frac{(b-a)(d-c)}{16} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|}{4} \right)^{\frac{1}{q}}, \quad (1.9)$$

where

$$A = \frac{1}{2} \left[\frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) + f(b, y)] dy \right].$$

In a recent paper, M. Matloka [19] introduced a new class of functions which are (h_1, h_2) -preinvex on the co-ordinates and established some Hermite-Hadamard and Fejér type inequalities for this class of functions.

Motivated by the results established in [19], the main aim of the present paper is to define preinvex functions on the co-ordinates and to establish some Hermite-Hadamard type inequalities for functions whose second order partial derivatives in absolute value are preinvex on the co-ordinates. Our established results generalize those result proved above in Theorem 5-Theorem 7.

2. MAIN RESULTS

In this section we first give notion of preinvex functions on the co-ordinates which generalize the classical convexity on the co-ordinates and then we prove some inequalities of Hermite-Hadamard type for such functions.

Definition 3. [19] *Let K_1 and K_2 be non-empty subsets of \mathbb{R}^n and let $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}^n$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}^n$. We say $K_1 \times K_2$ is invex with respect to η_1 and η_2 at $(u, v) \in K_1 \times K_2$ if for each $(x, y) \in K_1 \times K_2$ and $t, s \in [0, 1]$, we have*

$$(u + t\eta_1(x, u), v + s\eta_2(y, v)) \in X_1 \times X_2.$$

$K_1 \times K_2$ is said to be invex set with respect to η_1 and η_2 if $K_1 \times K_2$ is invex at each $(u, v) \in K_1 \times K_2$.

Definition 4. *Let $K_1 \times K_2$ is invex set with respect to $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}^n$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}^n$. A function $f : K_1 \times K_2 \rightarrow \mathbb{R}$ is said to be preinvex if for every $(x, y), (u, v) \in K_1 \times K_2$ and $t \in [0, 1]$, we have*

$$f(u + t\eta_1(x, u), v + t\eta_2(y, v)) \leq (1-t)f(x, y) + tf(u, v).$$

Definition 5. *Let $K_1 \times K_2$ be an invex set with respect to $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}^n$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}^n$. A function $f : K_1 \times K_2 \rightarrow \mathbb{R}$ is said to preinvex on the co-ordinates if the partial mappings $f_y : K_1 \rightarrow \mathbb{R}, f_y(u) = f(u, y)$ and $f_x : K_2 \rightarrow \mathbb{R}, f_x(v) = f(x, v)$ are preinvex with respect to η_1 and η_2 respectively for all $y \in K_2$ and $x \in K_1$.*

Remark 1. *If $\eta_1(x, u) = x - u$ and $\eta_2(y, v) = y - v$ then f will be a convex function on the coordinates.*

Remark 2. From the definition 5 it follows that if f is preinvex on the co-ordinates on $K_1 \times K_2$ then

$$\begin{aligned} f(u + t\eta_1(x, u), v + s\eta_2(y, v)) \\ \leq (1-t)(1-s)f(u, v) + (1-t)sf(u, y) \\ + (1-s)tf(x, v) + tsf(x, y). \end{aligned}$$

Remark 3. Every convex function on the co-ordinates is preinvex on the co-ordinates but the converse is not true. For example the function $f(u, v) = -|u||v|$ is not convex on the co-ordinates but it is a preinvex function with respect to the mappings

$$\eta_1(u, z) = \begin{cases} u - z, & u \geq 0, z \geq 0 \text{ and } u \leq 0, z \leq 0 \\ z - u, & \text{otherwise} \end{cases}$$

and

$$\eta_2(v, w) = \begin{cases} v - w, & v \geq 0, w \geq 0 \text{ and } v \leq 0, w \leq 0 \\ w - v, & \text{otherwise} \end{cases}.$$

The following Lemma is essential to establish our results:

Lemma 2. Let $K_1 \times K_2$ be an open invex subset of \mathbb{R}^2 with respect to the mappings $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}$. Suppose $f : K_1 \times K_2 \rightarrow \mathbb{R}$ be a twice partial differentiable mapping such that $\frac{\partial^2 f}{\partial t \partial s} \in L([a + t\eta_1(b, a)] \times [c + s\eta_2(d, c)])$ with $\eta_1(b, a) \neq 0$, $\eta_2(d, c) \neq 0$, where $a, b \in K_1$ and $c, d \in K_2$. Then the following equality holds:

$$\begin{aligned} & \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \\ & + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \\ & = \frac{\eta_1(b, a)\eta_2(d, c)}{4} \int_0^1 \int_0^1 (1-2t)(1-2s) \\ & \quad \times \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} dt ds, \quad (2.1) \end{aligned}$$

where

$$\begin{aligned} A' & = \frac{1}{2(\eta_1(b, a))} \int_a^{a+\eta_1(b, a)} [f(x, c) + f(x, c + \eta_2(d, c))] dx \\ & + \frac{1}{2(\eta_2(d, c))} \int_c^d [f(a, y) dy + f(a + \eta_1(b, a), y)] dy. \end{aligned}$$

Proof. By integration by parts with respect to t , we have

$$\begin{aligned}
 & \frac{\eta_1(b, a)\eta_2(d, c)}{4} \int_0^1 (1-2s) \left[\frac{-\frac{\partial f(a+t\eta_1(b, a), c+s\eta_2(d, c))}{\partial s} - \frac{\partial f(a, c+s\eta_2(d, c))}{\partial s}}{\eta_1(b, a)} \right]_0^1 \\
 & \quad + \frac{2}{\eta_1(b, a)} \int_0^1 \frac{\partial f(a+t\eta_1(b, a), c+s\eta_2(d, c))}{\partial s} dt \Big] ds \\
 & = -\frac{\eta_2(d, c)}{4} \int_0^1 (1-2s) \frac{\partial f(a+t\eta_1(b, a), c+s\eta_2(d, c))}{\partial s} ds \\
 & \quad - \frac{\eta_2(d, c)}{4} \int_0^1 (1-2s) \frac{\partial f(a, c+s\eta_2(d, c))}{\partial s} ds \\
 & \quad + \frac{\eta_2(d, c)}{2} \int_0^1 \int_0^1 (1-2s) \frac{\partial f(a+t\eta_1(b, a), c+s\eta_2(d, c))}{\partial s} ds dt. \quad (2.2)
 \end{aligned}$$

Integrating each integral on right hand side of (2.2) by parts with respect to s and using the substitution $x = a + t\eta_1(b, a)$ and $y = c + s\eta_2(d, c)$, we get the desired identity. This completes the proof of the lemma. \square

Theorem 8. Let $K_1 \times K_2$ be an open invex subset of \mathbb{R}^2 with respect to the mappings $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}$. Suppose $f : K_1 \times K_2 \rightarrow \mathbb{R}$ be a twice partial differentiable mapping such that $\frac{\partial^2 f}{\partial t \partial s} \in L([a + t\eta_1(b, a)] \times [c + s\eta_2(d, c)])$ with $\eta_1(b, a) \neq 0$, $\eta_2(d, c) \neq 0$, where $a, b \in K_1$ and $c, d \in K_2$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$ is preinvex on the co-ordinates on $K_1 \times K_2$, then the following inequality holds:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
 & \quad \left. + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
 & \leq \frac{\eta_1(b, a)\eta_2(d, c)}{16} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right| + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|}{4} \right), \quad (2.3)
 \end{aligned}$$

where A' is as defined in Lemma 2.

Proof. From Lemma 2 we have:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
 & \quad \left. + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
 & \leq \frac{\eta_1(b, a)\eta_2(d, c)}{4} \int_0^1 \int_0^1 |1-2t||1-2s| \left| \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} \right| dt ds \quad (2.4)
 \end{aligned}$$

By preinvexity of $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$ on the co-ordinates on $K_1 \times K_2$, we have

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
& \quad \left. + \frac{1}{\eta_1(b, a) \eta_2(d, c)} \int_a^{a + t\eta_1(b, a)} \int_c^{c + s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
& \leq \frac{\eta_1(b, a) \eta_2(d, c)}{4} \left[\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right| \int_0^1 \int_0^1 |1 - 2t| |1 - 2s| (1 - t) (1 - s) ds dt \right. \\
& \quad + \left| \frac{\partial f(a, d)}{\partial t \partial s} \right| \int_0^1 \int_0^1 |1 - 2t| |1 - 2s| (1 - t) s ds dt \\
& \quad + \left| \frac{\partial f(b, c)}{\partial t \partial s} \right| \int_0^1 \int_0^1 |1 - 2t| |1 - 2s| t (1 - s) ds dt \\
& \quad \left. + \left| \frac{\partial f(b, d)}{\partial t \partial s} \right| \int_0^1 \int_0^1 |1 - 2t| |1 - 2s| t s ds dt \right]. \quad (2.5)
\end{aligned}$$

Since

$$\begin{aligned}
\int_0^1 |1 - 2t| (1 - t) dt &= \int_0^{\frac{1}{2}} (1 - 2t) (1 - t) dt - \int_{\frac{1}{2}}^1 (1 - 2t) (1 - t) dt \\
&= \frac{1}{4}
\end{aligned}$$

and

$$\int_0^1 |1 - 2t| t dt = \int_0^{\frac{1}{2}} (1 - 2t) t dt - \int_{\frac{1}{2}}^1 (1 - 2t) t dt = \frac{1}{4}.$$

Making use the above in (2.5), we get the inequality (2.3). This completes the proof of the theorem. \square

Theorem 9. *Let $K_1 \times K_2$ be an open invex subset of \mathbb{R}^2 with respect to the mappings $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}$. Suppose $f : K_1 \times K_2 \rightarrow \mathbb{R}$ be a twice partial differentiable mapping such that $\frac{\partial^2 f}{\partial t \partial s} \in L([a + t\eta_1(b, a)] \times [c + s\eta_2(d, c)])$ with $\eta_1(b, a) \neq 0$, $\eta_2(d, c) \neq 0$, where $a, b \in K_1$ and $c, d \in K_2$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ is preinvex on the co-ordinates on $K_1 \times K_2$, $q \in (1, \infty)$, then the following inequality holds:*

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
& \quad \left. + \frac{1}{\eta_1(b, a) \eta_2(d, c)} \int_a^{a + t\eta_1(b, a)} \int_c^{c + s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
& \leq \frac{\eta_1(b, a) \eta_2(d, c)}{4(p+1)^{\frac{2}{p}}} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q}{4} \right)^{\frac{1}{q}}, \quad (2.6)
\end{aligned}$$

where A' is as defined in Lemma 2 and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 2 and Hölder's integral inequality, we have:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
 & \quad \left. + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
 & \leq \frac{\eta_1(b, a)\eta_2(d, c)}{4} \left(\int_0^1 \int_0^1 |1-2t|^p |1-2s|^p ds dt \right)^{\frac{1}{p}} \\
 & \quad \times \left(\int_0^1 \int_0^1 \left| \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} \right|^q dt ds \right)^{\frac{1}{q}} \quad (2.7)
 \end{aligned}$$

Since $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ is preinvex on the co-ordinates on $K_1 \times K_2$, we have

$$\begin{aligned}
 & \int_0^1 \int_0^1 \left| \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} \right|^q dt ds \\
 & \leq \left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-t)(1-s) dt ds + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-t) s dt ds \\
 & \quad + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-s) t dt ds + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 t s dt ds \\
 & = \frac{1}{4} \left[\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q \right]. \quad (2.8)
 \end{aligned}$$

Using (2.8) and

$$\int_0^1 \int_0^1 |1-2t|^p |1-2s|^p = \frac{1}{(p+1)^2}$$

in (2.7) gives us the desired inequality (2.6). This completes the proof of the theorem. \square

Theorem 10. Let $K_1 \times K_2$ be an open invex subset of \mathbb{R}^2 with respect to the mappings $\eta_1 : K_1 \times K_1 \rightarrow \mathbb{R}$ and $\eta_2 : K_2 \times K_2 \rightarrow \mathbb{R}$. Suppose $f : K_1 \times K_2 \rightarrow \mathbb{R}$ be a twice partial differentiable mapping such that $\frac{\partial^2 f}{\partial t \partial s} \in L([a + t\eta_1(b, a)] \times [c + s\eta_2(d, c)])$ with $\eta_1(b, a) \neq 0$, $\eta_2(d, c) \neq 0$, where $a, b \in K_1$ and $c, d \in K_2$. If $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ is preinvex on the co-ordinates on $K_1 \times K_2$, $q \in [1, \infty)$, then the following inequality holds:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\
 & \quad \left. + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\
 & \leq \frac{\eta_1(b, a)\eta_2(d, c)}{16} \left(\frac{\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q}{4} \right)^{\frac{1}{q}}, \quad (2.9)
 \end{aligned}$$

where A' is as defined in Lemma 2.

Proof. For $q = 1$, the proof is similar to that of Theorem 8. Suppose now that $q > 1$ then from Lemma 2 and the power-mean integral inequality, we have:

$$\begin{aligned} & \left| \frac{f(a, c) + f(a, c + \eta_2(d, c)) + f(a + \eta_1(b, a), c) + f(a + \eta_1(b, a), c + \eta_2(d, c))}{4} \right. \\ & \quad \left. + \frac{1}{\eta_1(b, a)\eta_2(d, c)} \int_a^{a+t\eta_1(b, a)} \int_c^{c+s\eta_2(d, c)} f(x, y) dy dx - A' \right| \\ & \leq \frac{\eta_1(b, a)\eta_2(d, c)}{4} \left(\int_0^1 \int_0^1 |1-2t||1-2s| ds dt \right)^{1-\frac{1}{q}} \\ & \quad \times \left(\int_0^1 \int_0^1 |1-2t||1-2s| \left| \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} \right|^q dt ds \right)^{\frac{1}{q}} \quad (2.10) \end{aligned}$$

Since $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ is preinvex on the co-ordinates on $K_1 \times K_2$, we have

$$\begin{aligned} & \int_0^1 \int_0^1 |1-2t||1-2s| \left| \frac{\partial^2 f(a + t\eta_1(b, a), c + s\eta_2(d, c))}{\partial t \partial s} \right|^q dt ds \\ & \leq \left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-t)(1-s)|1-2t||1-2s| dt ds \\ & \quad + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-t)s|1-2t||1-2s| dt ds \\ & \quad + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 (1-s)t|1-2t||1-2s| dt ds \\ & \quad + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q \int_0^1 \int_0^1 ts|1-2t||1-2s| dt ds \\ & = \frac{1}{16} \left[\left| \frac{\partial^2 f(a, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(a, d)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, c)}{\partial t \partial s} \right|^q + \left| \frac{\partial^2 f(b, d)}{\partial t \partial s} \right|^q \right]. \quad (2.11) \end{aligned}$$

Using (2.11) and

$$\int_0^1 \int_0^1 |1-2t||1-2s| = \frac{1}{4}$$

in (2.10) gives us the desired results. \square

Remark 4. Since $\frac{1}{4} < \frac{1}{(1+p)^{\frac{2}{p}}} < 1$, if $p > 1$; the estimation given in Theorem 9 is better than the one given in Theorem 10.

Remark 5. If $\eta_1(b, a) = b - a$ and $\eta_2(d, c) = d - c$, then we get those results proved in Theorem 5-Theorem 7 from [21]. This also reveals that our results are more general than those proved in [21].

REFERENCES

- [1] M. Alomari and M. Darus, The Hadamard's inequality for s -convex function of 2-variables on the co-ordinates, Int. Journal of Math. Analysis, 2 (13) (2008), 629–638.
- [2] M. Alomari and M. Darus, The Hadamard's inequality for s -convex function, Int. Journal of Math. Analysis, 2 (13) (2008), 639–646.
- [3] M. Alomari and M. Darus, On co-ordinated s -convex functions, International Mathematical Forum, 3, 2008, no. 40, 1977 - 1989.

- [4] M. Alomari and M. Darus, Hadamard-type Inequalities for s -convex functions, *International Mathematical Forum*, 3, 2008, no. 40, 1965-1975
- [5] T. Antczak, Mean value in invexity analysis, *Nonl. Anal.*, 60 (2005), 1473-1484.
- [6] A. Barani, A.G. Ghazanfari, S.S. Dragomir, Hermite-Hadamard inequality through prequasi- s -invex functions, *RGMA Research Report Collection*, 14(2011), Article 48, 7 pp.
- [7] A. Barani, A.G. Ghazanfari, S.S. Dragomir, Hermite-Hadamard inequality for functions whose derivatives absolute values are preinvex, *RGMA Research Report Collection*, 14(2011), Article 64, 11 pp.
- [8] A. Ben-Israel and B. Mond, What is invexity?, *J. Austral. Math. Soc., Ser. B*, 28(1986), No. 1, 1-9.
- [9] S. S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to Trapezoidal formula, *Appl. Math. Lett.* 11(5) (1998) 91-95.
- [10] S.S. Dragomir, On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, *Taiwanese Journal of Mathematics*, 4 (2001), 775-788.
- [11] D. Y. Hwang, K. L. Tseng and G. S. Yang, Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane, *Taiwanese Journal of Mathematics*, 11 (2007), 63-73.
- [12] M. A. Hanson, On sufficiency of the Kuhn-Tucker conditions, *J. Math. Anal. Appl.* 80 (1981) 545-550.
- [13] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, *J. Math Pures Appl.*, 58 (1893), 171-215.
- [14] I. Iscan, Ostrowski type inequalities for functions whose derivatives are preinvex, *arXiv:1204.2010v1.st. Math.*, 31 (1998), 354-364.
- [15] M. A. Latif and M. Alomari, Hadamard-type inequalities for product two convex functions on the co-ordinates, *Int. Math. Forum*, 4(47), 2009, 2327-2338.
- [16] M. A. Latif and M. Alomari, On the Hadamard-type inequalities for h -convex functions on the co-ordinates, *Int. J. of Math. Analysis*, 3(33), 2009, 1645-1656.
- [17] M. A. Latif and S. S. Dragomir, On some new inequalities for differentiable co-ordinated convex functions, *Journal of Inequalities and Applications* 2012, 2012:28 doi:10.1186/1029-242X-2012-28.
- [18] S. R. Mohan and S. K. Neogy, On invex sets and preinvex functions, *J. Math. Anal. Appl.* 189 (1995), 901-908.
- [19] M. Matloka, On some Hadamard-type inequalities for (h_1, h_2) -preinvex functions on the co-ordinates. (Submitted)
- [20] M. Aslam Noor, Hermite-Hadamard integral inequalities for log-preinvex functions, Preprint.
- [21] M. A. Noor, Variational-like inequalities, *Optimization*, 30 (1994), 323-330.
- [22] M. A. Noor, Invex equilibrium problems, *J. Math. Anal. Appl.*, 302 (2005), 463-475.
- [23] M. A. Noor, Some new classes of nonconvex functions, *Nonl. Funct. Anal. Appl.*, 11(2006), 165-171
- [24] M. A. Noor, On Hadamard integral inequalities involving two log-preinvex functions, *J. Inequal. Pure Appl. Math.*, 8(2007), No. 3, 1
- [25] M.E. Özdemir, E. Set and M.Z. Sarikaya, Some new Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions, *Hacet. J. Math Stat*, 40 (2) (2011), 219-229.
- [26] M. E. Özdemir, Havva Kavurmacı, Ahmet Ocak Akdemir and Merve Avcı, Inequalities for convex and s -convex functions on $\Delta = [a, b] \times [c, d]$, *Journal of Inequalities and Applications* 2012:20, doi:10.1186/1029-242X-2012-20.
- [27] M. E. Özdemir, M. A. Latif and A. O. Akdemir, On some Hadamard-type inequalities for product of two s -convex functions on the co-ordinates, *Journal of Inequalities and Applications*, 2012:21, doi:10.1186/1029-242X-2012-21.
- [28] M. E. Özdemir, Ahmet Ocak Akdemir and Melit Tunc, On the Hadamard-type inequalities for co-ordinated convex functions, *arXiv:1203.4327v1*.
- [29] C. M. E. Pearce and J. E. Pečarić, Inequalities for differentiable mappings with applications to special means and quadrature formula, *Appl. Math. Lett.* 13(2) (2000) 51-55.
- [30] M.Z. Sarikaya, E. Set, M.E. Özdemir and S. S. Dragomir, New some Hadamard's type inequalities for co-ordinated convex functions, *arXiv:1005.0700v1 [math.CA]*.
- [31] R. Pini, Invexity and generalized Convexity, *Optimization* 22 (1991) 513-525.

- [32] M. Z. Sarikaya, H. Bozkurt and N. Alp, On Hermite-Hadamard type integral inequalities for preinvex and log-preinvex functions, arXiv:1203.4759v1.
- [33] M. Z. Sarikaya and E. Set, New some Hadamard's type inequalities for co-ordinated convex functions, Tamsui Oxford Journal of Information and Mathematical Sciences 28(2) (2012) 137-152.
- [34] A. Saglam and H. Yidirim and M. Z. Sarikaya, Some new inequalities of Hermite-Hadamard's type, Kyungpook Math. J. 50(2010), 399-410.
- [35] T. Weir, and B. Mond, Preinvex functions in multiple bjective optimization, Journal of Mathematical Analysis and Applications, 136 (1998) 29-38.
- [36] X. M. Yang and D. Li, On properties of preinvex functions, J. Math. Anal. Appl. 256 (2001), 229-241.

COLLEGE OF SCIENCE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIL, HAIL-2440, SAUDI ARABIA

E-mail address: m_amer_latif@hotmail.com