

Generalised fractional Hermite-Hadamard Inequalities involving m -convexity and (s, m) -convexity

George A. Anastassiou
Department of Mathematical Sciences
University of Memphis
Memphis, TN 38152, U.S.A.
ganastss@memphis.edu

Abstract

Here we present generalised fractional Hermite-Hadamard type inequalities involving m -convexity and (s, m) -convexity. These inequalities are with respect to generalised Riemann-Liouville fractional integrals. Our work is motivated by and expands [7] to the greatest generality and all possible directions.

2010 AMS Subject Classification : 26A33, 26D10, 26D15.

Keywords and Phrases: fractional Hermite-Hadamard inequality, generalised fractional Riemann-Liouville integral, m -convexity, (s, m) -convexity.

1 Background

We use a lot here the following generalised fractional integrals.

Definition 1 (see also [3, p. 99]) *The left and right fractional integrals, respectively, of a function f with respect to given function g are defined as follows:*

Let $a, b \in \mathbb{R}$, $a < b$, $\alpha > 0$. Here $g \in AC([a, b])$ (absolutely continuous functions) and is strictly increasing, $f \in L_\infty([a, b])$. We set

$$(I_{a+;g}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (g(x) - g(t))^{\alpha-1} g'(t) f(t) dt, \quad x \geq a, \quad (1)$$

clearly $(I_{a+;g}^\alpha f)(a) = 0$,
and

$$(I_{b-;g}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (g(t) - g(x))^{\alpha-1} g'(t) f(t) dt, \quad x \leq b, \quad (2)$$

clearly $(I_{b-;g}^\alpha f)(b) = 0$.

When g is the identity function id , we get that $I_{a+;id}^\alpha = I_{a+}^\alpha$ and $I_{b-;id}^\alpha = I_{b-}^\alpha$ the ordinary left and right Riemann-Liouville fractional integrals, where

$$(I_{a+}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x \geq a, \quad (3)$$

$(I_{a+}^\alpha f)(a) = 0$, and

$$(I_{b-}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x \leq b, \quad (4)$$

$(I_{b-}^\alpha f)(b) = 0$.

Remark 2 (see also [1]) We observe that

$$(I_{a+;g}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (g(x) - g(t))^{\alpha-1} (f \circ g^{-1})(g(t)) g'(t) dt =$$

(by change of variable for Lebesgue integrals)

$$\frac{1}{\Gamma(\alpha)} \int_{g(a)}^{g(x)} (g(x) - z)^{\alpha-1} (f \circ g^{-1})(z) dz = \left(I_{g(a)+}^\alpha (f \circ g^{-1}) \right) (g(x)), \quad x \geq a, \quad (5)$$

equivalently $g(x) \geq g(a)$.

That is in the terms and assumptions of Definition 1 we get

$$(I_{a+;g}^\alpha f)(x) = \left(I_{g(a)+}^\alpha (f \circ g^{-1}) \right) (g(x)), \quad \text{for } x \geq a. \quad (6)$$

Similarly we observe that

$$\begin{aligned} (I_{b-;g}^\alpha f)(x) &= \frac{1}{\Gamma(\alpha)} \int_x^b (g(t) - g(x))^{\alpha-1} (f \circ g^{-1})(g(t)) g'(t) dt \\ &= \frac{1}{\Gamma(\alpha)} \int_{g(x)}^{g(b)} (z - g(x))^{\alpha-1} (f \circ g^{-1})(z) dz = \left(I_{g(b)-}^\alpha (f \circ g^{-1}) \right) (g(x)), \end{aligned} \quad (7)$$

for $x \leq b$.

That is

$$(I_{b-;g}^\alpha f)(x) = \left(I_{g(b)-}^\alpha (f \circ g^{-1}) \right) (g(x)), \quad \text{for } x \leq b. \quad (8)$$

So by (6) and (8) we have reduced the general fractional integrals to the ordinary left and right Riemann-Liouville fractional integrals.

When $g(x) = e^x$, $x \in [a, b]$ we have the application

Definition 3 The left and right fractional exponential integrals are defined as follows: Let $a, b \in \mathbb{R}$, $a < b$, $\alpha > 0$, $f \in L_\infty([a, b])$. We set

$$(I_{a+;e^x}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (e^x - e^t)^{\alpha-1} e^t f(t) dt, \quad x \geq a, \quad (9)$$

and

$$(I_{b-;e^x}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (e^t - e^x)^{\alpha-1} e^t f(t) dt, \quad x \leq b. \quad (10)$$

Note 4 We see that

$$(I_{a+;e^x}^\alpha f)(x) = (I_{e^a+}^\alpha (f \circ \ln))(e^x), \quad x \geq a, \quad (11)$$

and

$$(I_{b-;e^x}^\alpha f)(x) = (I_{e^b-}^\alpha (f \circ \ln))(e^x), \quad x \leq b. \quad (12)$$

Another example follows:

Definition 5 Let $a, b \in \mathbb{R}$, $a < b$, $\alpha > 0$, $f \in L_\infty([a, b])$, $A > 1$. We introduce the fractional integrals:

$$(I_{a+;A^x}^\alpha f)(x) = \frac{\ln A}{\Gamma(\alpha)} \int_a^x (A^x - A^t)^{\alpha-1} A^t f(t) dt, \quad x \geq a, \quad (13)$$

and

$$(I_{b-;A^x}^\alpha f)(x) = \frac{\ln A}{\Gamma(\alpha)} \int_x^b (A^t - A^x)^{\alpha-1} A^t f(t) dt, \quad x \leq b. \quad (14)$$

We are motivated by

Theorem 6 (1881, Hermite-Hadamard inequality, [4]) Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a convex function on the interval I of real numbers, and $a, b \in I$, with $a < b$. Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a) + f(b)}{2}. \quad (15)$$

Additionally to the classical convex functions, Toader [6], Hudzik and Maignanda [2] and Pinheiro [5] generalized the concepts of classical convex functions to the concepts of m -convex function and (s, m) -convex function.

Definition 7 The function $f : [0, b^*] \rightarrow \mathbb{R}$ is said to be m -convex, where $m \in [0, 1]$ and $b^* > 0$ if for every $x, y \in [0, b^*]$ and $t \in [0, 1]$, we have

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y). \quad (16)$$

Definition 8 The function $f : [0, b^*] \rightarrow \mathbb{R}$ is said to be (s, m) -convex, where $(s, m) \in [0, 1]^2$ and $b^* > 0$, if for every $x, y \in [0, b^*]$ and $t \in [0, 1]$, we have

$$f(tx + m(1-t)y) \leq t^s f(x) + m(1-t^s) f(y). \quad (17)$$

We need the following list of Lemmas and Theorems from [7].

Lemma 9 Let $\alpha > 0$, $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping on (a, b) with $a < b$. If $f'' \in L_1([a, b])$, then

$$\begin{aligned} & \frac{\Gamma(\alpha + 1)}{2(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] - f\left(\frac{a+b}{2}\right) = \\ & \frac{(b-a)^2}{2} \int_0^1 m(t) f''(ta + (1-t)b) dt, \end{aligned} \quad (18)$$

where

$$m(t) = \begin{cases} t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [0, \frac{1}{2}), \\ 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [\frac{1}{2}, 1]. \end{cases} \quad (19)$$

Lemma 10 Let $\alpha > 0$, $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping on (a, b) with $a < b$. If $f'' \in L_1([a, b])$, $r > 0$, then

$$\begin{aligned} & \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha + 1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] = \\ & (b-a)^2 \int_0^1 k(t) f''(ta + (1-t)b) dt, \end{aligned} \quad (20)$$

where

$$k(t) = \begin{cases} \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{r(\alpha+1)} - \frac{t}{r+1}, & t \in [0, \frac{1}{2}), \\ \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{r(\alpha+1)} - \frac{1-t}{r+1}, & t \in [\frac{1}{2}, 1]. \end{cases} \quad (21)$$

Lemma 11 Let $\alpha > 0$, $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping on (a, b) with $a < mb \leq b$. If $f'' \in L_1([a, b])$, $r > 0$, then

$$\begin{aligned} & \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha + 1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] = \\ & (mb-a)^2 \int_0^1 k(t) f''(ta + m(1-t)b) dt, \end{aligned} \quad (22)$$

where $k(t)$ is defined in (21).

The following fractional m -convex Hermite-Hadamard type inequalities also come from [7].

Theorem 12 Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q \geq 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then

$$H^m(f) := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha + 1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] \right|$$

$$\leq (b-a)^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right) \cdot \left(\frac{|f''(a)|^q + m|f''(\frac{b}{m})|^q}{2} \right)^{\frac{1}{q}} =: R_1^m(f). \quad (23)$$

Theorem 13 Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q > 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then

$$H^m(f) := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] \right| \leq \frac{(b-a)^2}{r(\alpha+1)} \left(1 - \frac{2}{p(\alpha+1)+1} \right)^{\frac{1}{p}} \left(\frac{|f''(a)|^q + m|f''(\frac{b}{m})|^q}{2} \right)^{\frac{1}{q}} =: R_2^m(f), \quad (24)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 14 Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q > 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then

$$H^m(f) := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] \right| \leq \frac{(b-a)^2}{r(\alpha+1)} \left(\frac{q(\alpha+1)-1}{q(\alpha+1)+1} \right)^{\frac{1}{q}} \left(\frac{|f''(a)|^q + m|f''(\frac{b}{m})|^q}{2} \right)^{\frac{1}{q}} =: R_3^m(f). \quad (25)$$

Theorem 15 Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q > 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then

$$H^m(f) := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] \right| \leq \left(\frac{2}{p+1} \right)^{\frac{1}{p}} \frac{(b-a)^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{p+1} - \left(\frac{r+1}{r(\alpha+1)} \right)^{p+1} \right]^{\frac{1}{p}} \cdot \left(\frac{|f''(a)|^q + m|f''(\frac{b}{m})|^q}{2} \right)^{\frac{1}{q}} =: R_4^m(f), \quad (26)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 16 Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q > 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then

$$\begin{aligned} H^m(f) &:= \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{r(b-a)^\alpha} [I_{a+}^\alpha f(b) + I_{b-}^\alpha f(a)] \right| \\ &\leq \left(\frac{2}{q+1} \right)^{\frac{1}{q}} \frac{(b-a)^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{q+1} - \left(\frac{r+1}{r(\alpha+1)} \right)^{q+1} \right]^{\frac{1}{q}} \\ &\quad \left(\frac{|f''(a)|^q + m |f''(\frac{b}{m})|^q}{2} \right)^{\frac{1}{q}} =: R_5^m(f). \end{aligned} \quad (27)$$

The following fractional (s, m) -convex Hermite-Hadamard type inequalities also come from [7].

Theorem 17 Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q \geq 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$\begin{aligned} H_s^m(f) &:= \\ &\left| \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] \right| \\ &\leq (mb-a)^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right)^{1-\frac{1}{q}}. \end{aligned} \quad (28)$$

$$\left[|f''(a)|^q I + m |f''(b)|^q \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} - I \right) \right]^{\frac{1}{q}} =: R_{1s}^m(f),$$

where

$$\begin{aligned} I &= \frac{1}{r(s+1)(s+\alpha+2)} - \frac{1}{r(\alpha+1)} B(s+1, \alpha+2) \\ &\quad + \frac{1}{(r+1)(s+1)(s+2)} \left(1 - \left(\frac{1}{2} \right)^{s+1} \right). \end{aligned}$$

Theorem 18 Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$\begin{aligned} H_s^m(f) &:= \\ &\left| \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] \right| \end{aligned}$$

$$\leq \frac{(mb-a)^2}{r(\alpha+1)} \left(1 - \frac{2}{p(\alpha+1)+1}\right)^{\frac{1}{p}} \left(\frac{1}{s+1} |f''(a)|^q + \frac{ms}{s+1} |f''(b)|^q\right)^{\frac{1}{q}} \quad (29)$$

$$=: R_{2s}^m(f),$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 19 Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$H_s^m(f) :=$$

$$\left| \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] \right|$$

$$\leq \frac{(mb-a)^2}{r(\alpha+1)} \left[|f''(a)|^q \left(\frac{1}{s+1} - \frac{1}{q(s+1)+s+1} - B(s+1, q(\alpha+1)+1) \right) \right. \quad (30)$$

$$\left. + m |f''(b)|^q \left(\frac{s}{s+1} - \frac{2}{q(\alpha+1)+1} + \frac{1}{q(\alpha+1)+s+1} \right) \right. \\ \left. + B(s+1, q(\alpha+1)+1) \right] =: R_{3s}^m(f).$$

Theorem 20 Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$H_s^m(f) :=$$

$$\left| \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] \right|$$

$$\leq \frac{(mb-a)^2}{r+1} \left(\frac{2}{p+1}\right)^{\frac{1}{p}} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)}\right)^{p+1} - \left(\frac{r+1}{r(\alpha+1)}\right)^{p+1} \right]^{\frac{1}{p}}.$$

$$\left(\frac{1}{s+1} |f''(a)|^q + \frac{ms}{s+1} |f''(b)|^q\right)^{\frac{1}{q}} =: R_{4s}^m(f), \quad (31)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 21 Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$H_s^m(f) :=$$

$$\left| \frac{f(a) + f(mb)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+mb}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mb-a)^\alpha} [I_{a+}^\alpha f(mb) + I_{mb-}^\alpha f(a)] \right|$$

$$\leq \frac{(mb-a)^2}{r+1} \left[|f''(a)|^q H + m |f''(b)|^q \left(\frac{2}{q+1} \left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{q+1} - \frac{2}{q+1} \left(\frac{r+1}{r(\alpha+1)} \right)^{q+1} - H \right) \right] =: R_{5s}^m(f), \quad (32)$$

where

$$H = \int_0^{\frac{1}{2}} \left(\frac{r+1}{r(\alpha+1)} + t \right)^q t^s dt + \int_{\frac{1}{2}}^1 \left(\frac{r+1}{r(\alpha+1)} + 1 - t \right)^q t^s dt. \quad (33)$$

The aim of this article is to extend the results of [7] to generalized fractional integrals (1) and (2), in particular to fractional exponential integrals (9), (10) and to fractional trigonometric integrals (60), (61). That is to produce very general fractional m -convex and (s, m) -convex Hermite-Hadamard type inequalities.

2 Main Results

Combining Theorems 12-16 we get the following m -convex Hermite-Hadamard type inequality.

Theorem 22 *Let $f : [0, b^*] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $b^* > 0$, $\alpha > 0$. If $|f''|^q$ is measurable and m -convex on $[a, \frac{b}{m}]$ for some fixed $q > 1$, $0 \leq a < b$ and $m \in (0, 1]$ with $\frac{b}{m} \leq b^*$, $r > 0$, then*

$$H^m(f) \leq \min \{R_1^m(f), R_2^m(f), R_3^m(f), R_4^m(f), R_5^m(f)\}. \quad (34)$$

Combining Theorems 17-21 we obtain the following (s, m) -convex Hermite-Hadamard type inequality.

Theorem 23 *Let $f : [0, b] \rightarrow \mathbb{R}$ be a twice differentiable mapping with $0 \leq a < mb \leq b$, $\alpha > 0$. If $|f''|^q$ is measurable and (s, m) -convex on $[a, b]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then*

$$H_s^m(f) \leq \min \{R_{1s}^m(f), R_{2s}^m(f), R_{3s}^m(f), R_{4s}^m(f), R_{5s}^m(f)\}. \quad (35)$$

Next we generalize Lemmas 9-11.

Lemma 24 *Let $\alpha > 0$, $a < b$, $f \in C([a, b])$, $g \in C^1([a, b])$, g strictly increasing on $[a, b]$, $(f \circ g^{-1})$ is twice differentiable function on $(g(a), g(b))$ with $(f \circ g^{-1})'' \in L_1([g(a), g(b)])$. Then*

$$\frac{\Gamma(\alpha+1)}{2(g(b)-g(a))^\alpha} [I_{a+;g}^\alpha f(b) + I_{b-;g}^\alpha f(a)] - (f \circ g^{-1}) \left(\frac{g(a)+g(b)}{2} \right) =$$

$$\frac{(g(b) - g(a))^2}{2} \int_0^1 m(t) (f \circ g^{-1})'' (tg(a) + (1-t)g(b)) dt, \quad (36)$$

where $m(t)$ as in (19).

Lemma 25 *Let all as in Lemma 24, $r > 0$. Then*

$$\begin{aligned} & \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} (f \circ g^{-1}) \left(\frac{g(a) + g(b)}{2} \right) \\ & - \frac{\Gamma(\alpha+1)}{r(g(b) - g(a))^\alpha} [I_{a+;g}^\alpha f(b) + I_{b-;g}^\alpha f(a)] \\ & = (g(b) - g(a))^2 \int_0^1 k(t) (f \circ g^{-1})'' (tg(a) + (1-t)g(b)) dt, \end{aligned} \quad (37)$$

where $k(t)$ as in (21).

Lemma 26 *Let all as Lemma 25, with $g(a) < mg(b) \leq g(b)$. Then*

$$\begin{aligned} & \frac{f(a) + (f \circ g^{-1})(mg(b))}{r(r+1)} + \frac{2}{r+1} (f \circ g^{-1}) \left(\frac{g(a) + mg(b)}{2} \right) \\ & - \frac{\Gamma(\alpha+1)}{r(mg(b) - g(a))^\alpha} \left[I_{g(a)+}^\alpha (f \circ g^{-1})(mg(b)) + I_{mg(b)-}^\alpha (f \circ g^{-1})(g(a)) \right] \\ & = (mg(b) - g(a))^2 \int_0^1 k(t) (f \circ g^{-1})'' (tg(a) + m(1-t)g(b)) dt, \end{aligned} \quad (38)$$

where $k(t)$ as in (21).

We apply Lemmas 24-26 to $g(x) = e^x$.

Lemma 27 *Let $\alpha > 0$, $a < b$, $f \in C([a, b])$, $(f \circ \ln)$ is twice differentiable function on (e^a, e^b) with $(f \circ \ln)'' \in L_1([e^a, e^b])$. Then*

$$\begin{aligned} & \frac{\Gamma(\alpha+1)}{2(e^b - e^a)^\alpha} [I_{a+;e^x}^\alpha f(b) + I_{b-;e^x}^\alpha f(a)] - (f \circ \ln) \left(\frac{e^a + e^b}{2} \right) = \\ & \frac{(e^b - e^a)^2}{2} \int_0^1 m(t) (f \circ \ln)'' (te^a + (1-t)e^b) dt, \end{aligned} \quad (39)$$

where $m(t)$ as in (19).

Lemma 28 *Let all as in Lemma 27, $r > 0$. Then*

$$\begin{aligned} & \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} (f \circ \ln) \left(\frac{e^a + e^b}{2} \right) - \frac{\Gamma(\alpha+1)}{r(e^b - e^a)^\alpha} [I_{a+;e^x}^\alpha f(b) + I_{b-;e^x}^\alpha f(a)] \\ & = (e^b - e^a)^2 \int_0^1 k(t) (f \circ \ln)'' (te^a + (1-t)e^b) dt, \end{aligned} \quad (40)$$

where $k(t)$ as in (21).

Lemma 29 *Let all as in Lemma 28, with $e^a < me^b \leq e^b$. Then*

$$\begin{aligned} & \frac{f(a) + (f \circ \ln)(me^b)}{r(r+1)} + \frac{2}{r+1} (f \circ \ln) \left(\frac{e^a + me^b}{2} \right) \\ & - \frac{\Gamma(\alpha+1)}{r(me^b - e^a)^\alpha} [I_{e^a+}^\alpha (f \circ \ln)(me^b) + I_{me^b-}^\alpha (f \circ \ln)(e^a)] \\ & = (me^b - e^a)^2 \int_0^1 k(t) (f \circ \ln)''(te^a + (1-t)e^b) dt, \end{aligned} \quad (41)$$

where $k(t)$ as in (21).

We need

Notation 30 *We denote by*

$$\begin{aligned} H^m(f, g) & := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} (f \circ g^{-1}) \left(\frac{g(a) + g(b)}{2} \right) - \right. \\ & \quad \left. \frac{\Gamma(\alpha+1)}{r(g(b) - g(a))^\alpha} [I_{a+;g}^\alpha f(b) + I_{b-;g}^\alpha f(a)] \right|, \end{aligned} \quad (42)$$

$$\begin{aligned} R_1^m(f, g) & := (g(b) - g(a))^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right) \cdot \\ & \quad \left(\frac{|(f \circ g^{-1})''(g(a))|^q + m |(f \circ g^{-1})''(\frac{g(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \end{aligned} \quad (43)$$

$$\begin{aligned} R_2^m(f, g) & := \frac{(g(b) - g(a))^2}{r(\alpha+1)} \left(1 - \frac{2}{p(\alpha+1)+1} \right)^{\frac{1}{p}} \cdot \\ & \quad \left(\frac{|(f \circ g^{-1})''(g(a))|^q + m |(f \circ g^{-1})''(\frac{g(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \end{aligned} \quad (44)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,

$$\begin{aligned} R_3^m(f, g) & := \frac{(g(b) - g(a))^2}{r(\alpha+1)} \left(\frac{q(\alpha+1)-1}{q(\alpha+1)+1} \right)^{\frac{1}{q}} \cdot \\ & \quad \left(\frac{|(f \circ g^{-1})''(g(a))|^q + m |(f \circ g^{-1})''(\frac{g(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \end{aligned} \quad (45)$$

$$R_4^m(f, g) := \left(\frac{2}{p+1} \right)^{\frac{1}{p}} \frac{(g(b) - g(a))^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{p+1} - \right.$$

$$-\left(\frac{r+1}{r(\alpha+1)}\right)^{p+1}\right]^{\frac{1}{p}}\left(\frac{|(f\circ g^{-1})''(g(a))|^q+m|(f\circ g^{-1})''\left(\frac{g(b)}{m}\right)|^q}{2}\right)^{\frac{1}{q}}, \quad (46)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,
and

$$R_5^m(f, g) := \left(\frac{2}{q+1}\right)^{\frac{1}{q}} \frac{(g(b) - g(a))^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)}\right)^{q+1} - \left(\frac{r+1}{r(\alpha+1)}\right)^{q+1}\right]^{\frac{1}{q}} \left(\frac{|(f\circ g^{-1})''(g(a))|^q+m|(f\circ g^{-1})''\left(\frac{g(b)}{m}\right)|^q}{2}\right)^{\frac{1}{q}}. \quad (47)$$

We present the following fractional generalised m -convex Hermite-Hadamard type inequality.

Theorem 31 *Let all as in Notation 30. Here $\alpha > 0$, $b^* > 0$, $f \in C([0, b^*])$, $g \in C^1([0, b^*])$, g is strictly increasing on $[0, b^*]$ with $g(0) = 0$. Assume that $f \circ g^{-1} : [0, g(b^*)] \rightarrow \mathbb{R}$ is twice differentiable mapping. If $|(f \circ g^{-1})''|^q$ is measurable and m -convex on $\left[g(a), \frac{g(b)}{m}\right]$ for some fixed $q > 1$, $0 \leq a < b \leq b^*$ and $m \in (0, 1]$ with $\frac{g(b)}{m} \leq g(b^*)$, $r > 0$, then*

$$H^m(f, g) \leq \min\{R_1^m(f, g), R_2^m(f, g), R_3^m(f, g), R_4^m(f, g), R_5^m(f, g)\}. \quad (48)$$

Proof. By Theorem 22. ■

We need

Notation 32 *We denote by*

$$H_s^m(f, g) := \left|\frac{f(a) + (f \circ g^{-1})(mg(b))}{r(r+1)} + \frac{2}{r+1}(f \circ g^{-1})\left(\frac{g(a) + mg(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(mg(b) - g(a))^\alpha} \left[I_{g(a)+}^\alpha (f \circ g^{-1})(mg(b)) + I_{mg(b)-}^\alpha (f \circ g^{-1})(g(a)) \right]\right|, \quad (49)$$

$$R_{1s}^m(f, g) := (mg(b) - g(a))^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)}\right)^{1-\frac{1}{q}} \left[|(f \circ g^{-1})''(g(a))|^q I + m |(f \circ g^{-1})''(g(b))|^q \right]. \quad (50)$$

$$\left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} - I\right)^{\frac{1}{q}},$$

where

$$I := \frac{1}{r(s+1)(s+\alpha+2)} - \frac{1}{r(\alpha+1)}B(s+1, \alpha+2) \\ + \frac{1}{(r+1)(s+1)(s+2)} \left(1 - \left(\frac{1}{2}\right)^{s+1}\right), \quad (51)$$

$$R_{2s}^m(f, g) := \frac{(mg(b) - g(a))^2}{r(\alpha+1)} \left(1 - \frac{2}{p(\alpha+1)+1}\right)^{\frac{1}{p}} \\ \left(\frac{1}{s+1} \left|(f \circ g^{-1})''(g(a))\right|^q + \frac{ms}{s+1} \left|(f \circ g^{-1})''(g(b))\right|^q\right)^{\frac{1}{q}}, \quad (52)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,

$$R_{3s}^m(f, g) := \frac{(mg(b) - g(a))^2}{r(\alpha+1)} \left[\left|(f \circ g^{-1})''(g(a))\right|^q \left(\frac{1}{s+1} - \frac{1}{q(\alpha+1)+s+1}\right) \right. \\ \left. - B(s+1, q(\alpha+1)+1) + m \left|(f \circ g^{-1})''(g(b))\right|^q \left(\frac{s}{s+1} - \frac{2}{q(\alpha+1)+1}\right) \right. \\ \left. + \frac{1}{q(\alpha+1)+s+1} + B(s+1, q(\alpha+1)+1) \right], \quad (53)$$

$$R_{4s}^m(f, g) := \frac{(mg(b) - g(a))^2}{r+1} \left(\frac{2}{p+1}\right)^{\frac{1}{p}} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)}\right)^{p+1} - \right. \\ \left. \left(\frac{r+1}{r(\alpha+1)}\right)^{p+1} \right]^{\frac{1}{p}} \left(\frac{1}{s+1} \left|(f \circ g^{-1})''(g(a))\right|^q + \frac{ms}{s+1} \left|(f \circ g^{-1})''(g(b))\right|^q\right)^{\frac{1}{q}}, \quad (54)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,
and

$$R_{5s}^m(f, g) := \frac{(mg(b) - g(a))^2}{r+1} \left[\left|(f \circ g^{-1})''(g(a))\right|^q H + m \left|(f \circ g^{-1})''(g(b))\right|^q \right. \\ \left. \left(\frac{2}{q+1} \left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)}\right)^{q+1} - \frac{2}{q+1} \left(\frac{r+1}{r(\alpha+1)}\right)^{q+1} - H\right) \right], \quad (55)$$

where

$$H = \int_0^{\frac{1}{2}} \left(\frac{r+1}{r(\alpha+1)} + t\right)^q t^s dt + \int_{\frac{1}{2}}^1 \left(\frac{r+1}{r(\alpha+1)} + 1 - t\right)^q t^s dt. \quad (56)$$

Next we present a fractional generalised (s, m) -convex Hermite-Hadamard type inequality.

Theorem 33 Here all as in Notation 32. Let $\alpha > 0$, $b > 0$, $f \in C([0, b])$, $g \in C^1([0, b])$, g is strictly increasing on $[0, b]$ with $g(0) = 0$. Assume that $f \circ g^{-1} : [0, g(b)] \rightarrow \mathbb{R}$ is twice differentiable mapping, with $0 \leq g(a) < mg(b) \leq g(b)$, $a \in [0, b]$. If $\left| (f \circ g^{-1})'' \right|^q$ is measurable and (s, m) -convex on $[g(a), g(b)]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$H_s^m(f, g) \leq \min \{R_{1s}^m(f, g), R_{2s}^m(f, g), R_{3s}^m(f, g), R_{4s}^m(f, g), R_{5s}^m(f, g)\}. \quad (57)$$

Proof. By Theorem 23. ■

The case $q = 1$ is met separately.

Proposition 34 Here $H^m(f, g)$ as in (42) of Notation 30. The rest of the assumptions as in Theorem 31 with $q = 1$. Then

$$H^m(f, g) \leq (g(b) - g(a))^2 \left(\frac{\alpha}{r(\alpha + 1)(\alpha + 2)} + \frac{1}{4(r + 1)} \right) \cdot \left(\frac{\left| (f \circ g^{-1})''(g(a)) \right| + m \left| (f \circ g^{-1})''\left(\frac{g(b)}{m}\right) \right|}{2} \right). \quad (58)$$

Proof. By Theorem 12. ■

Proposition 35 Here $H_s^m(f, g)$ as in (49) of Notation 32. The rest of the assumptions as in Theorem 33 with $q = 1$. Then

$$H_s^m(f, g) \leq (mg(b) - g(a))^2 \left[\left| (f \circ g^{-1})''(g(a)) \right| I + m \left| (f \circ g^{-1})''(g(b)) \right| \cdot \left(\frac{\alpha}{r(\alpha + 1)(\alpha + 2)} + \frac{1}{4(r + 1)} - I \right) \right], \quad (59)$$

where I as in (51).

Proof. By Theorem 17. ■

We need

Definition 36 Let $a, b \in [0, \frac{\pi}{2}]$, $a < b$, $\alpha > 0$, $f \in L_\infty([a, b])$. We consider the left and right fractional trigonometric integrals of f with respect to sine function denoted by \sin :

$$(I_{a+; \sin}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (\sin x - \sin t)^{\alpha-1} \cos t f(t) dt, \quad x \geq a, \quad (60)$$

and

$$(I_{b-; \sin}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (\sin t - \sin x)^{\alpha-1} \cos t f(t) dt, \quad x \leq b. \quad (61)$$

We need

Notation 37 We denote by

$$H_*^m(f, \sin) := \left| \frac{f(a) + f(b)}{r(r+1)} + \frac{2}{r+1} (f \circ \sin^{-1}) \left(\frac{\sin(a) + \sin(b)}{2} \right) - \frac{\Gamma(\alpha+1)}{r(\sin(b) - \sin(a))^\alpha} [I_{a+; \sin}^\alpha f(b) + I_{b-; \sin}^\alpha f(a)] \right|, \quad (62)$$

$$R_{1*}^m(f, \sin) := (\sin(b) - \sin(a))^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right) \cdot \left(\frac{|(f \circ \sin^{-1})''(\sin(a))|^q + m |(f \circ \sin^{-1})''(\frac{\sin(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \quad (63)$$

$$R_{2*}^m(f, \sin) := \frac{(\sin(b) - \sin(a))^2}{r(\alpha+1)} \left(1 - \frac{2}{p(\alpha+1)+1} \right)^{\frac{1}{p}} \cdot \left(\frac{|(f \circ \sin^{-1})''(\sin(a))|^q + m |(f \circ \sin^{-1})''(\frac{\sin(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \quad (64)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,

$$R_{3*}^m(f, \sin) := \frac{(\sin(b) - \sin(a))^2}{r(\alpha+1)} \left(\frac{q(\alpha+1)-1}{q(\alpha+1)+1} \right)^{\frac{1}{q}} \cdot \left(\frac{|(f \circ \sin^{-1})''(\sin(a))|^q + m |(f \circ \sin^{-1})''(\frac{\sin(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \quad (65)$$

$$R_{4*}^m(f, \sin) := \left(\frac{2}{p+1} \right)^{\frac{1}{p}} \frac{(\sin(b) - \sin(a))^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{p+1} - \left(\frac{r+1}{r(\alpha+1)} \right)^{p+1} \right]^{\frac{1}{p}} \cdot \left(\frac{|(f \circ \sin^{-1})''(\sin(a))|^q + m |(f \circ \sin^{-1})''(\frac{\sin(b)}{m})|^q}{2} \right)^{\frac{1}{q}}, \quad (66)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,
and

$$R_{5*}^m(f, \sin) := \left(\frac{2}{q+1} \right)^{\frac{1}{q}} \frac{(\sin(b) - \sin(a))^2}{r+1} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha+1)} \right)^{q+1} - \left(\frac{r+1}{r(\alpha+1)} \right)^{q+1} \right]^{\frac{1}{q}}$$

$$\left(\frac{r+1}{r(\alpha+1)} \right)^{q+1} \Big]^\frac{1}{q} \left(\frac{|(f \circ \sin^{-1})''(\sin(a))|^q + m |(f \circ \sin^{-1})''(\frac{\sin(b)}{m})|^q}{2} \right)^\frac{1}{q}. \quad (67)$$

We present the following fractional generalised m -convex Hermite-Hadamard type inequality for sin function. So here $g(x) = \sin(x)$, $x \in [0, \frac{\pi}{2}]$.

Theorem 38 *Let all as in Notation 37. Here $\alpha > 0$, $f \in C([0, \frac{\pi}{2}])$. Assume that $f \circ \sin^{-1} : [0, 1] \rightarrow \mathbb{R}$ is twice differentiable mapping. If $|(f \circ \sin^{-1})''|^q$ is measurable and m -convex on $[\sin(a), \frac{\sin(b)}{m}]$ for some fixed $q > 1$, $0 \leq a < b \leq \frac{\pi}{2}$ and $m \in (0, 1]$ with $\sin(b) \leq m$, $r > 0$, then*

$$H_*^m(f, \sin) \leq \min \{ R_{1*}^m(f, \sin), R_{2*}^m(f, \sin), R_{3*}^m(f, \sin), R_{4*}^m(f, \sin), R_{5*}^m(f, \sin) \}. \quad (68)$$

Proof. By Theorem 31. ■

We need

Notation 39 *We denote by*

$$H_{s*}^m(f, \sin) := \left| \frac{f(a) + (f \circ \sin^{-1})(m \sin(b))}{r(r+1)} + \frac{2}{r+1} (f \circ \sin^{-1}) \left(\frac{\sin(a) + m \sin(b)}{2} \right) - \frac{\Gamma(\alpha+1)}{r(m \sin(b) - \sin(a))^\alpha} \right. \\ \left. \left[I_{\sin(a)+}^\alpha (f \circ \sin^{-1})(m \sin(b)) + I_{m \sin(b)-}^\alpha (f \circ \sin^{-1})(\sin(a)) \right] \right|, \quad (69)$$

$$R_{1s*}^m(f, \sin) := (m \sin(b) - \sin(a))^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right)^{1-\frac{1}{q}} \\ \left[\left| (f \circ \sin^{-1})''(\sin(a)) \right|^q I + m \left| (f \circ \sin^{-1})''(\sin(b)) \right|^q \right. \\ \left. \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} - I \right) \right]^\frac{1}{q}, \quad (70)$$

where

$$I := \frac{1}{r(s+1)(s+\alpha+2)} - \frac{1}{r(\alpha+1)} B(s+1, \alpha+2) \\ + \frac{1}{(r+1)(s+1)(s+2)} \left(1 - \left(\frac{1}{2} \right)^{s+1} \right), \quad (71)$$

$$R_{2s^*}^m(f, \sin) := \frac{(m \sin(b) - \sin(a))^2}{r(\alpha + 1)} \left(1 - \frac{2}{p(\alpha + 1) + 1}\right)^{\frac{1}{p}} \cdot \left(\frac{1}{s+1} \left| (f \circ \sin^{-1})''(\sin(a)) \right|^q + \frac{ms}{s+1} \left| (f \circ \sin^{-1})''(\sin(b)) \right|^q\right)^{\frac{1}{q}}, \quad (72)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,

$$R_{3s^*}^m(f, \sin) := \frac{(m \sin(b) - \sin(a))^2}{r(\alpha + 1)} \left[\left| (f \circ \sin^{-1})''(\sin(a)) \right|^q \left(\frac{1}{s+1} - \frac{1}{q(\alpha + 1) + s + 1} - B(s+1, q(\alpha + 1) + 1) \right) + m \left| (f \circ \sin^{-1})''(\sin(b)) \right|^q \left(\frac{s}{s+1} - \frac{2}{q(\alpha + 1) + 1} + \frac{1}{q(\alpha + 1) + s + 1} + B(s+1, q(\alpha + 1) + 1) \right) \right], \quad (73)$$

$$R_{4s^*}^m(f, \sin) := \frac{(m \sin(b) - \sin(a))^2}{r+1} \left(\frac{2}{p+1}\right)^{\frac{1}{p}} \left[\left(\frac{1}{2} + \frac{r+1}{r(\alpha + 1)}\right)^{p+1} - \left(\frac{r+1}{r(\alpha + 1)}\right)^{p+1} \right]^{\frac{1}{p}} \left(\frac{1}{s+1} \left| (f \circ \sin^{-1})''(\sin(a)) \right|^q + \frac{ms}{s+1} \left| (f \circ \sin^{-1})''(\sin(b)) \right|^q \right)^{\frac{1}{q}}, \quad (74)$$

where $\frac{1}{p} + \frac{1}{q} = 1$,
and

$$R_{5s^*}^m(f, \sin) := \frac{(m \sin(b) - \sin(a))^2}{r+1} \left[\left| (f \circ \sin^{-1})''(\sin(a)) \right|^q H + m \left| (f \circ \sin^{-1})''(\sin(b)) \right|^q \left(\frac{2}{q+1} \left(\frac{1}{2} + \frac{r+1}{r(\alpha + 1)}\right)^{q+1} - \frac{2}{q+1} \left(\frac{r+1}{r(\alpha + 1)}\right)^{q+1} - H \right) \right], \quad (75)$$

where

$$H = \int_0^{\frac{1}{2}} \left(\frac{r+1}{r(\alpha + 1)} + t\right)^q t^s dt + \int_{\frac{1}{2}}^1 \left(\frac{r+1}{r(\alpha + 1)} + 1 - t\right)^q t^s dt. \quad (76)$$

Next we present a fractional generalised (s, m) -convex Hermite-Hadamard type inequality involving $g(x) = \sin x$, $x \in [0, \frac{\pi}{2}]$.

Theorem 40 Here all as in Notation 39. Let $\alpha > 0$, $a, b \in [0, \frac{\pi}{2}]$, $a < b$, $f \in C([0, b])$. Assume that $f \circ \sin^{-1} : [0, \sin(b)] \rightarrow \mathbb{R}$ is twice differentiable mapping, with $0 \leq \sin(a) < m \sin(b) \leq \sin(b)$. If $\left| (f \circ \sin^{-1})'' \right|^q$ is measurable and (s, m) -convex on $[\sin(a), \sin(b)]$ for some fixed $q > 1$ and $(s, m) \in (0, 1]^2$, $r > 0$, then

$$H_{s^*}^m(f, \sin) \leq \min \{R_{1s^*}^m(f, \sin), R_{2s^*}^m(f, \sin), R_{3s^*}^m(f, \sin), R_{4s^*}^m(f, \sin), R_{5s^*}^m(f, \sin)\}. \quad (77)$$

Proof. By Theorem 33. ■

Finally we treat the case of $q = 1$ when $g(x) = \sin x$, $x \in [0, \frac{\pi}{2}]$.

Proposition 41 Here $H_{s^*}^m(f, \sin)$ as in (62) of Notation 37. The rest of the assumptions as in Theorem 38 with $q = 1$. Then

$$H_{s^*}^m(f, \sin) \leq (\sin(b) - \sin(a))^2 \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} \right) \cdot \left(\frac{\left| (f \circ \sin^{-1})''(\sin(a)) \right| + m \left| (f \circ \sin^{-1})''\left(\frac{\sin(b)}{m}\right) \right|}{2} \right). \quad (78)$$

Proof. By Proposition 34. ■

Proposition 42 Here $H_{s^*}^m(f, \sin)$ as in (69) of Notation 39. The rest of the assumptions as in Theorem 40 with $q = 1$. Then

$$H_{s^*}^m(f, \sin) \leq (m \sin(b) - \sin(a))^2 \left[\left| (f \circ \sin^{-1})''(\sin(a)) \right| I + m \left| (f \circ \sin^{-1})''(\sin(b)) \right| \left(\frac{\alpha}{r(\alpha+1)(\alpha+2)} + \frac{1}{4(r+1)} - I \right) \right], \quad (79)$$

where I as in (51).

Proof. By Proposition 35. ■

References

- [1] G.A. Anastassiou, *The reduction method in fractional Calculus and fractional Ostrowski type inequalities*, submitted 2013.
- [2] H. Hudzik, L. Maligranda, *Some remarks on s -convex functions*, Aequationes Math. 48 (1994), 100-111.

- [3] A.A. Kilbas, H.M. Srivastava and J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, vol. 204 of North-Holland Mathematics Studies, Elsevier, New York, NY, USA, 2006.
- [4] D.S. Mitrinović, I.B. Lacković, *Hermite and convexity*, Aequationes Math., 28 (1985), 229-232.
- [5] M.R. Pinheiro, *Exploring the concept of s -convexity*, Aequat. Math., 74 (2007), 201-209.
- [6] G.H. Toader, *Some generalisations of the convexity*, Proc. Colloq. Approx. Optim., (1984), 329-338.
- [7] Yuruo Zhang, Jin Rong Wang, *On some New Hermite-Hadamard inequalities involving Riemann-Liouville Fractional Integrals*, Journal of Inequalities and Applications, 2013, 2013: 220, doi: 10.1186 / 1029-242X-2013-220.