

**TRACE INEQUALITIES OF CASSELS AND GRÜSS TYPE FOR
OPERATORS IN HILBERT SPACES**

S. S. DRAGOMIR^{1,2}

ABSTRACT. Some trace inequalities of Cassels type for operators in Hilbert spaces are provided. Applications in connection to Grüss inequality and for convex functions of selfadjoint operators are also given.

1. INTRODUCTION

Let $\bar{\mathbf{a}} = (a_1, \dots, a_n)$ and $\bar{\mathbf{b}} = (b_1, \dots, b_n)$ be two positive n -tuples with

$$(1.1) \quad 0 < m_1 \leq a_i \leq M_1 < \infty \text{ and } 0 < m_2 \leq b_i \leq M_2 < \infty;$$

for each $i \in \{1, \dots, n\}$, and some constants m_1, m_2, M_1, M_2 .

The following reverses of the Cauchy-Bunyakovsky-Schwarz inequality for positive sequences of real numbers are well known:

a) *Pólya-Szegő's inequality* [51]:

$$\frac{\sum_{k=1}^n a_k^2 \sum_{k=1}^n b_k^2}{\left(\sum_{k=1}^n a_k b_k\right)^2} \leq \frac{1}{4} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right)^2.$$

b) *Shisha-Mond's inequality* [55]:

$$\frac{\sum_{k=1}^n a_k^2}{\sum_{k=1}^n a_k b_k} - \frac{\sum_{k=1}^n a_k b_k}{\sum_{k=1}^n b_k^2} \leq \left[\left(\frac{M_1}{m_2} \right)^{\frac{1}{2}} - \left(\frac{m_1}{M_2} \right)^{\frac{1}{2}} \right]^2.$$

c) *Ozeki's inequality* [48]:

$$\sum_{k=1}^n a_k^2 \sum_{k=1}^n b_k^2 - \left(\sum_{k=1}^n a_k b_k \right)^2 \leq \frac{n^2}{4} (M_1 M_2 - m_1 m_2)^2.$$

d) *Diaz-Metcalf's inequality* [17]:

$$\sum_{k=1}^n b_k^2 + \frac{m_2 M_2}{m_1 M_1} \sum_{k=1}^n a_k^2 \leq \left(\frac{M_2}{m_1} + \frac{m_2}{M_1} \right) \sum_{k=1}^n a_k b_k.$$

If $\bar{\mathbf{w}} = (w_1, \dots, w_n)$ is a positive sequence, then the following weighted inequalities also hold:

e) *Cassels' inequality* [58]. If the positive real sequences $\bar{\mathbf{a}} = (a_1, \dots, a_n)$ and $\bar{\mathbf{b}} = (b_1, \dots, b_n)$ satisfy the condition

$$(1.2) \quad 0 < m \leq \frac{a_k}{b_k} \leq M < \infty \text{ for each } k \in \{1, \dots, n\},$$

1991 *Mathematics Subject Classification.* 47A63; 47A99.

Key words and phrases. Trace class operators, Hilbert-Schmidt operators, Schwarz inequality, Grüss inequality.

then

$$\frac{\left(\sum_{k=1}^n w_k a_k^2\right) \left(\sum_{k=1}^n w_k b_k^2\right)}{\left(\sum_{k=1}^n w_k a_k b_k\right)^2} \leq \frac{(M+m)^2}{4mM}.$$

f) *Greub-Reinboldt's inequality* [38]. We have

$$\left(\sum_{k=1}^n w_k a_k^2\right) \left(\sum_{k=1}^n w_k b_k^2\right) \leq \frac{(M_1 M_2 + m_1 m_2)^2}{4m_1 m_2 M_1 M_2} \left(\sum_{k=1}^n w_k a_k b_k\right)^2,$$

provided $\bar{\mathbf{a}} = (a_1, \dots, a_n)$ and $\bar{\mathbf{b}} = (b_1, \dots, b_n)$ satisfy the condition (1.1).

g) *Generalized Diaz-Metcalf's inequality* [17], see also [46, p. 123]. If $u, v \in [0, 1]$ and $v \leq u$, $u + v = 1$ and (1.2) holds, then one has the inequality

$$u \sum_{k=1}^n w_k b_k^2 + v M m \sum_{k=1}^n w_k a_k^2 \leq (vm + uM) \sum_{k=1}^n w_k a_k b_k.$$

h) *Klamkin-McLenaghan's inequality* [40]. If $\bar{\mathbf{a}}, \bar{\mathbf{b}}$ satisfy (1.2), then

$$(1.3) \quad \begin{aligned} & \left(\sum_{i=1}^n w_i a_i^2\right) \left(\sum_{i=1}^n w_i b_i^2\right) - \left(\sum_{i=1}^n w_i a_i b_i\right)^2 \\ & \leq \left(M^{\frac{1}{2}} - m^{\frac{1}{2}}\right)^2 \sum_{i=1}^n w_i a_i b_i \sum_{i=1}^n w_i a_i^2. \end{aligned}$$

For other recent results providing discrete reverse inequalities, see the monograph online [19].

The following reverse of Schwarz's inequality in inner product spaces holds [20].

Theorem 1 (Dragomir, 2003, [20]). *Let $A, a \in \mathbb{C}$ and $x, y \in H$, a complex inner product space with the inner product $\langle \cdot, \cdot \rangle$. If*

$$(1.4) \quad \operatorname{Re} \langle Ay - x, x - ay \rangle \geq 0,$$

or equivalently,

$$(1.5) \quad \left\| x - \frac{a + A}{2} \cdot y \right\| \leq \frac{1}{2} |A - a| \|y\|,$$

holds, then we have the inequality

$$(1.6) \quad 0 \leq \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 \leq \frac{1}{4} |A - a|^2 \|y\|^4.$$

The constant $\frac{1}{4}$ is sharp in (1.6).

In 1935, G. Grüss [39] proved the following integral inequality which gives an approximation of the integral mean of the product in terms of the product of the integrals means as follows:

$$(1.7) \quad \left| \frac{1}{b-a} \int_a^b f(x) g(x) dx - \frac{1}{b-a} \int_a^b f(x) dx \cdot \frac{1}{b-a} \int_a^b g(x) dx \right| \leq \frac{1}{4} (\Phi - \phi) (\Gamma - \gamma),$$

where $f, g : [a, b] \rightarrow \mathbb{R}$ are integrable on $[a, b]$ and satisfy the condition

$$(1.8) \quad \phi \leq f(x) \leq \Phi, \quad \gamma \leq g(x) \leq \Gamma$$

for each $x \in [a, b]$, where $\phi, \Phi, \gamma, \Gamma$ are given real constants.

Moreover, the constant $\frac{1}{4}$ is sharp in the sense that it cannot be replaced by a smaller one.

In [22], in order to generalize the Grüss integral inequality in abstract structures the author has proved the following inequality in inner product spaces.

Theorem 2 (Dragomir, 1999, [22]). *Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space over \mathbb{K} ($\mathbb{K} = \mathbb{R}, \mathbb{C}$) and $e \in H$, $\|e\| = 1$. If $\varphi, \gamma, \Phi, \Gamma$ are real or complex numbers and x, y are vectors in H such that the conditions*

$$(1.9) \quad \operatorname{Re} \langle \Phi e - x, x - \varphi e \rangle \geq 0 \text{ and } \operatorname{Re} \langle \Gamma e - y, y - \gamma e \rangle \geq 0$$

hold, then we have the inequality

$$(1.10) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle| \leq \frac{1}{4} |\Phi - \varphi| |\Gamma - \gamma|.$$

The constant $\frac{1}{4}$ is best possible in the sense that it can not be replaced by a smaller constant.

For other results of this type, see the recent monograph [25] and the references therein.

For other Grüss type results for integral and sums see the papers [1]-[3], [8]-[10], [11]-[13], [21]-[28], [35], [49], [62] and the references therein.

In order to state some reverses of Schwarz and Grüss type inequalities for trace operators on complex Hilbert spaces we need some preparations as follows.

2. SOME FACTS ON TRACE OF OPERATORS

Let $(H, \langle \cdot, \cdot \rangle)$ be a complex Hilbert space and $\{e_i\}_{i \in I}$ an *orthonormal basis* of H . We say that $A \in \mathcal{B}(H)$ is a *Hilbert-Schmidt operator* if

$$(2.1) \quad \sum_{i \in I} \|Ae_i\|^2 < \infty.$$

It is well know that, if $\{e_i\}_{i \in I}$ and $\{f_j\}_{j \in J}$ are orthonormal bases for H and $A \in \mathcal{B}(H)$ then

$$(2.2) \quad \sum_{i \in I} \|Ae_i\|^2 = \sum_{j \in J} \|Af_j\|^2 = \sum_{j \in J} \|A^* f_j\|^2$$

showing that the definition (2.1) is independent of the orthonormal basis and A is a Hilbert-Schmidt operator iff A^* is a Hilbert-Schmidt operator.

Let $\mathcal{B}_2(H)$ the set of Hilbert-Schmidt operators in $\mathcal{B}(H)$. For $A \in \mathcal{B}_2(H)$ we define

$$(2.3) \quad \|A\|_2 := \left(\sum_{i \in I} \|Ae_i\|^2 \right)^{1/2}$$

for $\{e_i\}_{i \in I}$ an orthonormal basis of H . This definition does not depend on the choice of the orthonormal basis.

Using the triangle inequality in $l^2(I)$, one checks that $\mathcal{B}_2(H)$ is a *vector space* and that $\|\cdot\|_2$ is a norm on $\mathcal{B}_2(H)$, which is usually called in the literature as the *Hilbert-Schmidt norm*.

Denote *the modulus* of an operator $A \in \mathcal{B}(H)$ by $|A| := (A^*A)^{1/2}$.

Because $\| |A|x \| = \|Ax\|$ for all $x \in H$, A is Hilbert-Schmidt iff $|A|$ is Hilbert-Schmidt and $\|A\|_2 = \||A|\|_2$. From (2.2) we have that if $A \in \mathcal{B}_2(H)$, then $A^* \in \mathcal{B}_2(H)$ and $\|A\|_2 = \|A^*\|_2$.

The following theorem collects some of the most important properties of Hilbert-Schmidt operators:

Theorem 3. *We have*

(i) $(\mathcal{B}_2(H), \|\cdot\|_2)$ is a Hilbert space with inner product

$$(2.4) \quad \langle A, B \rangle_2 := \sum_{i \in I} \langle Ae_i, Be_i \rangle = \sum_{i \in I} \langle B^* Ae_i, e_i \rangle$$

and the definition does not depend on the choice of the orthonormal basis $\{e_i\}_{i \in I}$;

(ii) We have the inequalities

$$(2.5) \quad \|A\| \leq \|A\|_2$$

for any $A \in \mathcal{B}_2(H)$ and

$$(2.6) \quad \|AT\|_2, \|TA\|_2 \leq \|T\| \|A\|_2$$

for any $A \in \mathcal{B}_2(H)$ and $T \in \mathcal{B}(H)$;

(iii) $\mathcal{B}_2(H)$ is an operator ideal in $\mathcal{B}(H)$, i.e.

$$\mathcal{B}(H) \mathcal{B}_2(H) \mathcal{B}(H) \subseteq \mathcal{B}_2(H);$$

(iv) $\mathcal{B}_{fin}(H)$, the space of operators of finite rank, is a dense subspace of $\mathcal{B}_2(H)$;

(v) $\mathcal{B}_2(H) \subseteq \mathcal{K}(H)$, where $\mathcal{K}(H)$ denotes the algebra of compact operators on H .

If $\{e_i\}_{i \in I}$ an orthonormal basis of H , we say that $A \in \mathcal{B}(H)$ is trace class if

$$(2.7) \quad \|A\|_1 := \sum_{i \in I} \langle |A| e_i, e_i \rangle < \infty.$$

The definition of $\|A\|_1$ does not depend on the choice of the orthonormal basis $\{e_i\}_{i \in I}$. We denote by $\mathcal{B}_1(H)$ the set of trace class operators in $\mathcal{B}(H)$.

The following proposition holds:

Proposition 1. *If $A \in \mathcal{B}(H)$, then the following are equivalent:*

(i) $A \in \mathcal{B}_1(H)$;

(ii) $|A|^{1/2} \in \mathcal{B}_2(H)$;

(iii) A (or $|A|$) is the product of two elements of $\mathcal{B}_2(H)$.

The following properties are also well known:

Theorem 4. *With the above notations:*

(i) We have

$$(2.8) \quad \|A\|_1 = \|A^*\|_1 \quad \text{and} \quad \|A\|_2 \leq \|A\|_1$$

for any $A \in \mathcal{B}_1(H)$;

(ii) $\mathcal{B}_1(H)$ is an operator ideal in $\mathcal{B}(H)$, i.e.

$$\mathcal{B}(H) \mathcal{B}_1(H) \mathcal{B}(H) \subseteq \mathcal{B}_1(H);$$

(iii) We have

$$\mathcal{B}_2(H) \mathcal{B}_2(H) = \mathcal{B}_1(H);$$

(iv) We have

$$\|A\|_1 = \sup \{ \langle A, B \rangle_2 \mid B \in \mathcal{B}_2(H), \|B\| \leq 1 \};$$

(v) $(\mathcal{B}_1(H), \|\cdot\|_1)$ is a Banach space.

(iv) We have the following isometric isomorphisms

$$\mathcal{B}_1(H) \cong K(H)^* \text{ and } \mathcal{B}_1(H)^* \cong \mathcal{B}(H),$$

where $K(H)^*$ is the dual space of $K(H)$ and $\mathcal{B}_1(H)^*$ is the dual space of $\mathcal{B}_1(H)$.

We define the *trace* of a trace class operator $A \in \mathcal{B}_1(H)$ to be

$$(2.9) \quad \text{tr}(A) := \sum_{i \in I} \langle Ae_i, e_i \rangle,$$

where $\{e_i\}_{i \in I}$ an orthonormal basis of H . Note that this coincides with the usual definition of the trace if H is finite-dimensional. We observe that the series (2.9) converges absolutely and it is independent from the choice of basis.

The following result collects some properties of the trace:

Theorem 5. *We have*

(i) If $A \in \mathcal{B}_1(H)$ then $A^* \in \mathcal{B}_1(H)$ and

$$(2.10) \quad \text{tr}(A^*) = \overline{\text{tr}(A)};$$

(ii) If $A \in \mathcal{B}_1(H)$ and $T \in \mathcal{B}(H)$, then $AT, TA \in \mathcal{B}_1(H)$ and

$$(2.11) \quad \text{tr}(AT) = \text{tr}(TA) \text{ and } |\text{tr}(AT)| \leq \|A\|_1 \|T\|;$$

(iii) $\text{tr}(\cdot)$ is a bounded linear functional on $\mathcal{B}_1(H)$ with $\|\text{tr}\| = 1$;

(iv) If $A, B \in \mathcal{B}_2(H)$ then $AB, BA \in \mathcal{B}_1(H)$ and $\text{tr}(AB) = \text{tr}(BA)$;

(v) $\mathcal{B}_{fin}(H)$ is a dense subspace of $\mathcal{B}_1(H)$.

Utilising the trace notation we obviously have that

$$\langle A, B \rangle_2 = \text{tr}(B^*A) = \text{tr}(AB^*) \text{ and } \|A\|_2^2 = \text{tr}(A^*A) = \text{tr}(|A|^2)$$

for any $A, B \in \mathcal{B}_2(H)$.

The following Hölder's type inequality has been obtained by Ruskai in [52]

$$(2.12) \quad |\text{tr}(AB)| \leq \text{tr}(|AB|) \leq \left[\text{tr}(|A|^{1/\alpha}) \right]^\alpha \left[\text{tr}(|B|^{1/(1-\alpha)}) \right]^{1-\alpha}$$

where $\alpha \in (0, 1)$ and $A, B \in \mathcal{B}(H)$ with $|A|^{1/\alpha}, |B|^{1/(1-\alpha)} \in \mathcal{B}_1(H)$.

In particular, for $\alpha = \frac{1}{2}$ we get the Schwarz inequality

$$(2.13) \quad |\text{tr}(AB)| \leq \text{tr}(|AB|) \leq \left[\text{tr}(|A|^2) \right]^{1/2} \left[\text{tr}(|B|^2) \right]^{1/2}$$

with $A, B \in \mathcal{B}_2(H)$.

For the theory of trace functionals and their applications the reader is referred to [56].

For some classical trace inequalities see [14], [16], [47] and [61], which are continuations of the work of Bellman [5]. For related works the reader can refer to [4], [6], [14], [36], [41], [42], [44], [53] and [57].

We denote by

$$\mathcal{B}_1^+(H) := \{P : P \in \mathcal{B}_1(H), P \text{ and is selfadjoint and } P \geq 0\}.$$

We obtained recently the following result [33]:

Theorem 6. For any $A, C \in \mathcal{B}(H)$ and $P \in \mathcal{B}_1^+(H) \setminus \{0\}$ we have the inequality

$$(2.14) \quad \begin{aligned} & \left| \frac{\operatorname{tr}(PAC)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right| \\ & \leq \inf_{\lambda \in \mathbb{C}} \|A - \lambda \cdot 1_H\| \frac{1}{\operatorname{tr}(P)} \operatorname{tr} \left(\left| \left(C - \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} 1_H \right) P \right| \right) \\ & \leq \inf_{\lambda \in \mathbb{C}} \|A - \lambda \cdot 1_H\| \left[\frac{\operatorname{tr}(P|C|^2)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right|^2 \right]^{1/2}, \end{aligned}$$

where $\|\cdot\|$ is the operator norm.

We also have [33]:

Corollary 1. Let $\alpha, \beta \in \mathbb{C}$ and $A \in \mathcal{B}(H)$ such that

$$\left\| A - \frac{\alpha + \beta}{2} \cdot 1_H \right\| \leq \frac{1}{2} |\beta - \alpha|.$$

For any $C \in \mathcal{B}(H)$ and $P \in \mathcal{B}_1^+(H) \setminus \{0\}$ we have the inequality

$$(2.15) \quad \begin{aligned} & \left| \frac{\operatorname{tr}(PAC)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right| \\ & \leq \frac{1}{2} |\beta - \alpha| \frac{1}{\operatorname{tr}(P)} \operatorname{tr} \left(\left| \left(C - \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} 1_H \right) P \right| \right) \\ & \leq \frac{1}{2} |\beta - \alpha| \left[\frac{\operatorname{tr}(P|C|^2)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right|^2 \right]^{1/2}. \end{aligned}$$

In particular, if $C \in \mathcal{B}(H)$ is such that

$$\left\| C - \frac{\alpha + \beta}{2} \cdot 1_H \right\| \leq \frac{1}{2} |\beta - \alpha|,$$

then

$$(2.16) \quad \begin{aligned} 0 & \leq \frac{\operatorname{tr}(P|C|^2)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right|^2 \\ & \leq \frac{1}{2} |\beta - \alpha| \frac{1}{\operatorname{tr}(P)} \operatorname{tr} \left(\left| \left(C - \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} 1_H \right) P \right| \right) \\ & \leq \frac{1}{2} |\beta - \alpha| \left[\frac{\operatorname{tr}(P|C|^2)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right|^2 \right]^{1/2} \leq \frac{1}{4} |\beta - \alpha|^2. \end{aligned}$$

Also

$$\begin{aligned}
(2.17) \quad & \left| \frac{\operatorname{tr}(PC^2)}{\operatorname{tr}(P)} - \left(\frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right)^2 \right| \\
& \leq \frac{1}{2} |\beta - \alpha| \frac{1}{\operatorname{tr}(P)} \operatorname{tr} \left(\left| \left(C - \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} 1_H \right) P \right| \right) \\
& \leq \frac{1}{2} |\beta - \alpha| \left[\frac{\operatorname{tr}(P|C|^2)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PC)}{\operatorname{tr}(P)} \right|^2 \right]^{1/2} \leq \frac{1}{4} |\beta - \alpha|^2.
\end{aligned}$$

For other related results see [33].

3. CASSELS TYPE TRACE INEQUALITIES

For two given operators $T, U \in B(H)$ and two given scalars $\alpha, \beta \in \mathbb{C}$ consider the transform

$$\mathcal{C}_{\alpha,\beta}(T, U) = (T^* - \bar{\alpha}U^*)(\beta U - T).$$

This transform generalizes the transform

$$\mathcal{C}_{\alpha,\beta}(T) := (T^* - \bar{\alpha}1_H)(\beta 1_H - T) = \mathcal{C}_{\alpha,\beta}(T, 1_H),$$

where 1_H is the identity operator, which has been introduced in [31] in order to provide some generalizations of the well known Kantorovich inequality for operators in Hilbert spaces.

We recall that a bounded linear operator T on the complex Hilbert space $(H, \langle \cdot, \cdot \rangle)$ is called *accretive* if $\operatorname{Re} \langle Ty, y \rangle \geq 0$ for any $y \in H$.

Utilizing the following identity

$$\begin{aligned}
(3.1) \quad \operatorname{Re} \langle \mathcal{C}_{\alpha,\beta}(T, U)x, x \rangle &= \operatorname{Re} \langle \mathcal{C}_{\beta,\alpha}(T, U)x, x \rangle \\
&= \frac{1}{4} |\beta - \alpha|^2 \|Ux\|^2 - \left\| Tx - \frac{\alpha + \beta}{2} \cdot Ux \right\|^2 \\
&= \frac{1}{4} |\beta - \alpha|^2 \langle |U|^2 x, x \rangle - \left\langle \left| T - \frac{\alpha + \beta}{2} \cdot U \right|^2 x, x \right\rangle
\end{aligned}$$

that holds for any scalars α, β and any vector $x \in H$, we can give a simple characterization result that is useful in the following:

Lemma 1. *For $\alpha, \beta \in \mathbb{C}$ and $T, U \in B(H)$ the following statements are equivalent:*

- (i) *The transform $\mathcal{C}_{\alpha,\beta}(T, U)$ (or, equivalently, $\mathcal{C}_{\beta,\alpha}(T, U)$) is accretive;*
- (ii) *We have the norm inequality*

$$(3.2) \quad \left\| Tx - \frac{\alpha + \beta}{2} \cdot Ux \right\| \leq \frac{1}{2} |\beta - \alpha| \|Ux\|$$

for any $x \in H$;

- (iii) *We have the following inequality in the operator order*

$$\left| T - \frac{\alpha + \beta}{2} \cdot U \right|^2 \leq \frac{1}{4} |\beta - \alpha|^2 |U|^2.$$

As a consequence of the above lemma we can state:

Corollary 2. *Let $\alpha, \beta \in \mathbb{C}$ and $T, U \in B(H)$. If $\mathcal{C}_{\alpha, \beta}(T, U)$ is accretive, then*

$$(3.3) \quad \left\| T - \frac{\alpha + \beta}{2} \cdot U \right\| \leq \frac{1}{2} |\beta - \alpha| \|U\|.$$

Remark 1. *In order to give examples of linear operators $T, U \in B(H)$ and numbers $\alpha, \beta \in \mathbb{C}$ such that the transform $\mathcal{C}_{\alpha, \beta}(T, U)$ is accretive, it suffices to select two bounded linear operator S and V and the complex numbers z, w ($w \neq 0$) with the property that $\|Sx - zVx\| \leq |w| \|Vx\|$ for any $x \in H$, and, by choosing $T = S$, $U = V$, $\alpha = \frac{1}{2}(z + w)$ and $\beta = \frac{1}{2}(z - w)$ we observe that T and U satisfy (3.2), i.e., $\mathcal{C}_{\alpha, \beta}(T, U)$ is accretive.*

The following result also holds:

Lemma 2. *Let, either $P \in \mathcal{B}_+(H)$, $A, B \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B \in \mathcal{B}(H)$ and $\gamma, \Gamma \in \mathbb{C}$. Then*

$$(3.4) \quad \operatorname{Re}(\operatorname{tr}[P(A^* - \bar{\gamma}B^*)(\Gamma B - A)]) \geq 0$$

if and only if

$$(3.5) \quad \operatorname{tr} \left(P \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \right) \leq \frac{1}{4} |\Gamma - \gamma|^2 \operatorname{tr} (P |B|^2).$$

To simplify the writing, we say that (A, B) satisfies the P - (γ, Γ) -trace property.

Proof. We have the equalities

$$(3.6) \quad \begin{aligned} & \frac{1}{4} |\Gamma - \gamma|^2 P |B|^2 - P \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \\ &= P \left[\frac{1}{4} |\Gamma - \gamma|^2 |B|^2 - \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \right] \\ &= P \left[\frac{1}{4} |\Gamma - \gamma|^2 |B|^2 - \left(A - \frac{\gamma + \Gamma}{2} B \right)^* \left(A - \frac{\gamma + \Gamma}{2} B \right) \right] \\ &= P \left[\frac{1}{4} |\Gamma - \gamma|^2 |B|^2 \right. \\ & \quad \left. - |A|^2 + \frac{\overline{\gamma + \Gamma}}{2} B^* A + \frac{\gamma + \Gamma}{2} A^* B - \left| \frac{\gamma + \Gamma}{2} \right|^2 |B|^2 \right] \\ &= P \left[-|A|^2 + \frac{\overline{\gamma + \Gamma}}{2} B^* A + \frac{\gamma + \Gamma}{2} A^* B \right. \\ & \quad \left. + \left(\frac{1}{4} |\Gamma - \gamma|^2 - \left| \frac{\gamma + \Gamma}{2} \right|^2 \right) |B|^2 \right] \\ &= P \left[-|A|^2 + \frac{\overline{\gamma + \Gamma}}{2} B^* A + \frac{\gamma + \Gamma}{2} A^* B - \operatorname{Re}(\Gamma \bar{\gamma}) |B|^2 \right] \end{aligned}$$

for any bounded operators A, B, P and the complex numbers $\gamma, \Gamma \in \mathbb{C}$.

Taking the trace in (3.6) we get

$$\begin{aligned}
(3.7) \quad & \frac{1}{4} |\Gamma - \gamma|^2 \operatorname{tr} (P |B|^2) - \operatorname{tr} \left(P \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \right) \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) \\
&+ \frac{\gamma + \Gamma}{2} \operatorname{tr} (PB^* A) + \frac{\gamma + \Gamma}{2} \operatorname{tr} (PA^* B) \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) \\
&+ \frac{\gamma + \Gamma}{2} \operatorname{tr} (PB^* A) + \frac{\gamma + \Gamma}{2} \overline{\operatorname{tr} (PB^* A)} \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) + \frac{\gamma + \Gamma}{2} \operatorname{tr} (PB^* A) + \overline{\frac{\gamma + \Gamma}{2} \operatorname{tr} (PB^* A)} \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) + 2 \operatorname{Re} \left[\frac{\gamma + \Gamma}{2} \operatorname{tr} (PB^* A) \right] \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) + \operatorname{Re} [\bar{\gamma} \operatorname{tr} (PB^* A)] + \operatorname{Re} [\Gamma \operatorname{tr} (PB^* A)] \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) + \operatorname{Re} [\bar{\gamma} \operatorname{tr} (PB^* A)] + \operatorname{Re} [\overline{\Gamma \operatorname{tr} (PB^* A)}] \\
&= -\operatorname{tr} (P |A|^2) - \operatorname{Re} (\Gamma \bar{\gamma}) \operatorname{tr} (P |B|^2) + \operatorname{Re} [\bar{\gamma} \operatorname{tr} (PB^* A)] + \operatorname{Re} [\Gamma \operatorname{tr} (PB^* A)].
\end{aligned}$$

Since

$$\begin{aligned}
& \operatorname{Re} (\operatorname{tr} [P (A^* - \bar{\gamma} B^*) (\Gamma B - A)]) \\
&= \operatorname{Re} [\operatorname{tr} (\Gamma P A^* B + \bar{\gamma} P B^* A - \bar{\gamma} \Gamma P B^* B - P A^* A)] \\
&= \operatorname{Re} [\Gamma \operatorname{tr} (P A^* B) + \bar{\gamma} \operatorname{tr} (P B^* A)] \\
&- \bar{\gamma} \Gamma \operatorname{tr} (P |B|^2) - \operatorname{tr} (P |A|^2) \\
&= \operatorname{Re} [\Gamma \overline{\operatorname{tr} (P B^* A)} + \bar{\gamma} \operatorname{tr} (P B^* A)] \\
&- \operatorname{tr} (P |B|^2) \operatorname{Re} (\bar{\gamma} \Gamma) - \operatorname{tr} (P |A|^2),
\end{aligned}$$

then we get

$$\begin{aligned}
(3.8) \quad & \frac{1}{4} |\Gamma - \gamma|^2 \operatorname{tr} (P |B|^2) - \operatorname{tr} \left(P \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \right) \\
&= \operatorname{Re} (\operatorname{tr} [P (A^* - \bar{\gamma} B^*) (\Gamma B - A)]),
\end{aligned}$$

which proves the desired equivalence. \square

Corollary 3. *Let, either $P \in \mathcal{B}_+(H)$, $A, B \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B \in \mathcal{B}(H)$ and $\gamma, \Gamma \in \mathbb{C}$. If the transform $\mathcal{C}_{\gamma, \Gamma}(A, B)$ is accretive, then (A, B) satisfies the P - (γ, Γ) -trace property.*

We have the following result:

Theorem 7. *Let, either $P \in \mathcal{B}_+(H)$, $A, B \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B \in \mathcal{B}(H)$ and $\gamma, \Gamma \in \mathbb{C}$ with $\operatorname{Re} (\Gamma \bar{\gamma}) = \operatorname{Re} (\Gamma) \operatorname{Re} (\gamma) + \operatorname{Im} (\Gamma) \operatorname{Im} (\gamma) > 0$.*

(i) If (A, B) satisfies the P - (γ, Γ) -trace property, then we have

$$(3.9) \quad \begin{aligned} & \operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |B|^2 \right) \\ & \leq \frac{1}{4} \cdot \frac{[\operatorname{Re}(\gamma + \Gamma) \operatorname{Re} \operatorname{tr}(PB^*A) + \operatorname{Im}(\gamma + \Gamma) \operatorname{Im} \operatorname{tr}(PB^*A)]^2}{\operatorname{Re}(\Gamma) \operatorname{Re}(\gamma) + \operatorname{Im}(\Gamma) \operatorname{Im}(\gamma)} \\ & \leq \frac{1}{4} \cdot \frac{|\gamma + \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} |\operatorname{tr}(PB^*A)|^2. \end{aligned}$$

(ii) If the transform $\mathcal{C}_{\gamma, \Gamma}(A, B)$ is accretive, then the inequality (3.9) also holds.

Proof. (i) If (A, B) satisfies the P - (γ, Γ) -trace property, then, on utilizing the calculations above, we have

$$\begin{aligned} 0 & \leq \frac{1}{4} |\Gamma - \gamma|^2 \operatorname{tr} \left(P |B|^2 \right) - \operatorname{tr} \left(P \left| A - \frac{\gamma + \Gamma}{2} B \right|^2 \right) \\ & = -\operatorname{tr} \left(P |A|^2 \right) - \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right) \\ & \quad + \operatorname{Re}[\bar{\gamma} \operatorname{tr}(PB^*A)] + \operatorname{Re}[\Gamma \overline{\operatorname{tr}(PB^*A)}] \\ & = -\operatorname{tr} \left(P |A|^2 \right) - \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right) \\ & \quad + \operatorname{Re}[\bar{\gamma} \operatorname{tr}(PB^*A)] + \operatorname{Re}[\Gamma \overline{\operatorname{tr}(PB^*A)}] \\ & = -\operatorname{tr} \left(P |A|^2 \right) - \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right) \\ & \quad + \operatorname{Re}[\bar{\gamma} \operatorname{tr}(PB^*A)] + \operatorname{Re}[\bar{\Gamma} \operatorname{tr}(PB^*A)] \\ & = -\operatorname{tr} \left(P |A|^2 \right) - \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right) + \operatorname{Re}[(\bar{\gamma} + \bar{\Gamma}) \operatorname{tr}(PB^*A)], \end{aligned}$$

which implies that

$$(3.10) \quad \begin{aligned} & \operatorname{tr} \left(P |A|^2 \right) + \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right) \\ & \leq \operatorname{Re}[(\bar{\gamma} + \bar{\Gamma}) \operatorname{tr}(PB^*A)] \\ & = \operatorname{Re}(\gamma + \Gamma) \operatorname{Re} \operatorname{tr}(PB^*A) + \operatorname{Im}(\gamma + \Gamma) \operatorname{Im} \operatorname{tr}(PB^*A). \end{aligned}$$

Making use of the elementary inequality

$$2\sqrt{pq} \leq p + q, \quad p, q \geq 0,$$

we also have

$$(3.11) \quad 2\sqrt{\operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |B|^2 \right)} \leq \operatorname{tr} \left(P |A|^2 \right) + \operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{tr} \left(P |B|^2 \right).$$

Utilising (3.10) and (3.11) we get

$$(3.12) \quad \begin{aligned} & \sqrt{\operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |B|^2 \right)} \\ & \leq \frac{\operatorname{Re}(\gamma + \Gamma) \operatorname{Re} \operatorname{tr}(PB^*A) + \operatorname{Im}(\gamma + \Gamma) \operatorname{Im} \operatorname{tr}(PB^*A)}{2\sqrt{\operatorname{Re}(\Gamma\bar{\gamma})}} \end{aligned}$$

that is equivalent with the first inequality in (3.9).

The second inequality in (3.9) is obvious by Schwarz inequality

$$(ab + cd)^2 \leq (a^2 + c^2)(b^2 + d^2), \quad a, b, c, d \in \mathbb{R}.$$

The (ii) is obvious from (i). \square

Remark 2. We observe that the inequality between the first and last term in (3.9) is equivalent to

$$(3.13) \quad 0 \leq \operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |B|^2 \right) - |\operatorname{tr} (PB^*A)|^2 \leq \frac{1}{4} \cdot \frac{|\gamma - \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} |\operatorname{tr} (PB^*A)|^2.$$

Corollary 4. Let, either $P \in \mathcal{B}_+(H)$, $A \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A \in \mathcal{B}(H)$ and $\gamma, \Gamma \in \mathbb{C}$ with $\operatorname{Re}(\Gamma\bar{\gamma}) = \operatorname{Re}(\Gamma)\operatorname{Re}(\gamma) + \operatorname{Im}(\Gamma)\operatorname{Im}(\gamma) > 0$.

(i) If A satisfies the P - (γ, Γ) -trace property, namely

$$(3.14) \quad \operatorname{Re}(\operatorname{tr} [P(A^* - \bar{\gamma}1_H)(\Gamma1_H - A)]) \geq 0$$

or, equivalently

$$(3.15) \quad \operatorname{tr} \left(P \left| A - \frac{\gamma + \Gamma}{2} 1_H \right|^2 \right) \leq \frac{1}{4} |\Gamma - \gamma|^2 \operatorname{tr}(P),$$

then we have

$$(3.16) \quad \begin{aligned} & \frac{\operatorname{tr} \left(P |A|^2 \right)}{\operatorname{tr}(P)} \\ & \leq \frac{1}{4} \cdot \frac{\left[\operatorname{Re}(\gamma + \Gamma) \frac{\operatorname{Re} \operatorname{tr}(PA)}{\operatorname{tr}(P)} + \operatorname{Im}(\gamma + \Gamma) \frac{\operatorname{Im} \operatorname{tr}(PA)}{\operatorname{tr}(P)} \right]^2}{\operatorname{Re}(\Gamma)\operatorname{Re}(\gamma) + \operatorname{Im}(\Gamma)\operatorname{Im}(\gamma)} \\ & \leq \frac{1}{4} \cdot \frac{|\gamma + \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} \left| \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right|^2. \end{aligned}$$

(ii) If the transform $\mathcal{C}_{\gamma, \Gamma}(A)$ is accretive, then the inequality (3.9) also holds.

(iii) We have

$$(3.17) \quad 0 \leq \frac{\operatorname{tr} \left(P |A|^2 \right)}{\operatorname{tr}(P)} - \left| \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right|^2 \leq \frac{1}{4} \cdot \frac{|\gamma - \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} \left| \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right|^2.$$

Remark 3. The case of selfadjoint operators is as follows.

Let A, B be selfadjoint operators and either $P \in \mathcal{B}_+(H)$, $A, B \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B \in \mathcal{B}(H)$ and $m, M \in \mathbb{R}$ with $mM > 0$.

(i) If (A, B) satisfies the P - (m, M) -trace property, then we have

$$(3.18) \quad \operatorname{tr} (PA^2) \operatorname{tr} (PB^2) \leq \frac{(m+M)^2}{4mM} [\operatorname{tr} (PBA)]^2$$

or, equivalently

$$(3.19) \quad 0 \leq \operatorname{tr} (PA^2) \operatorname{tr} (PB^2) - [\operatorname{tr} (PBA)]^2 \leq \frac{(m-M)^2}{4mM} [\operatorname{tr} (PBA)]^2.$$

(ii) If the transform $\mathcal{C}_{m, M}(A, B)$ is accretive, then the inequality (3.18) also holds.

(iii) If $(A - mB)(MB - A) \geq 0$, then (3.18) is valid.

We observe that the inequality (3.18) is the operator trace inequality version of Cassels' inequality from Introduction.

4. TRACE INEQUALITIES OF GRÜSS TYPE

Let P be a selfadjoint operator with $P \geq 0$. The functional $\langle \cdot, \cdot \rangle_{2,P}$ defined by

$$\langle A, B \rangle_{2,P} := \operatorname{tr}(PB^*A) = \operatorname{tr}(APB^*) = \operatorname{tr}(B^*AP)$$

is a *nonnegative Hermitian form* on $\mathcal{B}_2(H)$, i.e. $\langle \cdot, \cdot \rangle_{2,P}$ satisfies the properties:

- (h) $\langle A, A \rangle_{2,P} \geq 0$ for any $A \in \mathcal{B}_2(H)$;
- (hh) $\langle \cdot, \cdot \rangle_{2,P}$ is linear in the first variable;
- (hhh) $\langle B, A \rangle_{2,P} = \overline{\langle A, B \rangle_{2,P}}$ for any $A, B \in \mathcal{B}_2(H)$.

Using the properties of the trace we also have the following representations

$$\|A\|_{2,P}^2 := \operatorname{tr}(P|A|^2) = \operatorname{tr}(APA^*) = \operatorname{tr}(|A|^2P)$$

and

$$\langle A, B \rangle_{2,P} = \operatorname{tr}(APB^*) = \operatorname{tr}(B^*AP)$$

for any $A, B \in \mathcal{B}_2(H)$.

The same definitions can be considered if $P \in \mathcal{B}_1^+(H)$ and $A, B \in \mathcal{B}(H)$.

We have the following Grüss type inequality:

Theorem 8. *Let, either $P \in \mathcal{B}_+(H)$, $A, B, C \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B, C \in \mathcal{B}(H)$ with $P|A|^2, P|B|^2, P|C|^2 \neq 0$ and $\lambda, \Gamma, \delta, \Delta \in \mathbb{C}$ with $\operatorname{Re}(\Gamma\bar{\gamma})$, $\operatorname{Re}(\Delta\bar{\delta}) > 0$. If (A, C) has the trace P - (λ, Γ) -property and (B, C) has the trace P - (δ, Δ) -property, then*

$$(4.1) \quad \left| \frac{\operatorname{tr}(PB^*A) \operatorname{tr}(P|C|^2)}{\operatorname{tr}(PC^*A) \operatorname{tr}(PB^*C)} - 1 \right| \leq \frac{1}{4} \cdot \frac{|\gamma - \Gamma| |\delta - \Delta|}{\sqrt{\operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{Re}(\Delta\bar{\delta})}}.$$

Proof. We prove in the case that $P \in \mathcal{B}_+(H)$ and $A, B, C \in \mathcal{B}_2(H)$.

Making use of the Schwarz inequality for the nonnegative hermitian form $\langle \cdot, \cdot \rangle_{2,P}$ we have

$$\left| \langle A, B \rangle_{2,P} \right|^2 \leq \langle A, A \rangle_{2,P} \langle B, B \rangle_{2,P}$$

for any $A, B \in \mathcal{B}_2(H)$.

Let $C \in \mathcal{B}_2(H)$, $C \neq 0$. Define the mapping $[\cdot, \cdot]_{2,P,C} : \mathcal{B}_2(H) \times \mathcal{B}_2(H) \rightarrow \mathbb{C}$ by

$$[A, B]_{2,P,C} := \langle A, B \rangle_{2,P} \|C\|_{2,P}^2 - \langle A, C \rangle_{2,P} \langle C, B \rangle_{2,P}.$$

Observe that $[\cdot, \cdot]_{2,P,C}$ is a nonnegative Hermitian form on $\mathcal{B}_2(H)$ and by Schwarz inequality we also have

$$\begin{aligned} & \left| \langle A, B \rangle_{2,P} \|C\|_{2,P}^2 - \langle A, C \rangle_{2,P} \langle C, B \rangle_{2,P} \right|^2 \\ & \leq \left[\|A\|_{2,P}^2 \|C\|_{2,P}^2 - \left| \langle A, C \rangle_{2,P} \right|^2 \right] \left[\|B\|_{2,P}^2 \|C\|_{2,P}^2 - \left| \langle B, C \rangle_{2,P} \right|^2 \right] \end{aligned}$$

for any $A, B \in \mathcal{B}_2(H)$, namely

$$(4.2) \quad \begin{aligned} & \left| \operatorname{tr}(PB^*A) \operatorname{tr}(P|C|^2) - \operatorname{tr}(PC^*A) \operatorname{tr}(PB^*C) \right|^2 \\ & \leq \left[\operatorname{tr}(P|A|^2) \operatorname{tr}(P|C|^2) - |\operatorname{tr}(PC^*A)|^2 \right] \\ & \quad \times \left[\operatorname{tr}(P|B|^2) \operatorname{tr}(P|C|^2) - |\operatorname{tr}(PB^*C)|^2 \right], \end{aligned}$$

where for the last term we used the equality $\left| \langle B, C \rangle_{2,P} \right|^2 = \left| \langle C, B \rangle_{2,P} \right|^2$.

Since (A, C) has the trace P - (λ, Γ) -property and (B, C) has the trace P - (δ, Δ) -property, then by (3.13) we have

$$(4.3) \quad 0 \leq \operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |C|^2 \right) - |\operatorname{tr} (PC^*A)|^2 \leq \frac{1}{4} \cdot \frac{|\gamma - \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} |\operatorname{tr} (PC^*A)|^2$$

and

$$(4.4) \quad 0 \leq \operatorname{tr} \left(P |B|^2 \right) \operatorname{tr} \left(P |C|^2 \right) - |\operatorname{tr} (PB^*C)|^2 \leq \frac{1}{4} \cdot \frac{|\delta - \Delta|^2}{\operatorname{Re}(\Delta\bar{\delta})} |\operatorname{tr} (PB^*C)|^2.$$

If we multiply the inequalities (4.3) and (4.4) we get

$$(4.5) \quad \begin{aligned} & \left[\operatorname{tr} \left(P |A|^2 \right) \operatorname{tr} \left(P |C|^2 \right) - |\operatorname{tr} (PC^*A)|^2 \right] \\ & \times \left[\operatorname{tr} \left(P |B|^2 \right) \operatorname{tr} \left(P |C|^2 \right) - |\operatorname{tr} (PB^*C)|^2 \right] \\ & \leq \frac{1}{16} \cdot \frac{|\gamma - \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} \frac{|\delta - \Delta|^2}{\operatorname{Re}(\Delta\bar{\delta})} |\operatorname{tr} (PC^*A)|^2 |\operatorname{tr} (PB^*C)|^2. \end{aligned}$$

If we use (4.2) and (4.5) we get

$$(4.6) \quad \begin{aligned} & \left| \operatorname{tr} (PB^*A) \operatorname{tr} \left(P |C|^2 \right) - \operatorname{tr} (PC^*A) \operatorname{tr} (PB^*C) \right|^2 \\ & \leq \frac{1}{16} \cdot \frac{|\gamma - \Gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} \frac{|\delta - \Delta|^2}{\operatorname{Re}(\Delta\bar{\delta})} |\operatorname{tr} (PC^*A)|^2 |\operatorname{tr} (PB^*C)|^2. \end{aligned}$$

Since $P, A, B, C \neq 0$ then by (4.3) and (4.4) we get $\operatorname{tr} (PC^*A) \neq 0$ and $\operatorname{tr} (PB^*C) \neq 0$. Now, if we take the square root in (4.6) and divide by $|\operatorname{tr} (PC^*A) \operatorname{tr} (PB^*C)|$ we obtain the desired result (4.1). \square

Corollary 5. *Let, either $P \in \mathcal{B}_+(H)$, $A, B \in \mathcal{B}_2$ or $P \in \mathcal{B}_1^+(H)$, $A, B \in \mathcal{B}(H)$ with $P|A|^2, P|B|^2 \neq 0$ and $\lambda, \Gamma, \delta, \Delta \in \mathbb{C}$ with $\operatorname{Re}(\Gamma\bar{\gamma}), \operatorname{Re}(\Delta\bar{\delta}) > 0$. If A has the trace P - (λ, Γ) -property and B has the trace P - (δ, Δ) -property, then*

$$(4.7) \quad \left| \frac{\operatorname{tr} (PB^*A) \operatorname{tr} (P)}{\operatorname{tr} (PA) \operatorname{tr} (PB^*)} - 1 \right| \leq \frac{1}{4} \cdot \frac{|\gamma - \Gamma| |\delta - \Delta|}{\sqrt{\operatorname{Re}(\Gamma\bar{\gamma}) \operatorname{Re}(\Delta\bar{\delta})}}.$$

The case of selfadjoint operators is useful for applications.

Remark 4. *Assume that A, B, C are selfadjoint operators. If, either $P \in \mathcal{B}_+(H)$, $A, B, C \in \mathcal{B}_2(H)$ or $P \in \mathcal{B}_1^+(H)$, $A, B, C \in \mathcal{B}(H)$ with $PA^2, PB^2, PC^2 \neq 0$ and $m, M, n, N \in \mathbb{R}$ with $mM, nN > 0$. If (A, C) has the trace P - (m, M) -property and (B, C) has the trace P - (n, N) -property, then*

$$(4.8) \quad \left| \frac{\operatorname{tr} (PBA) \operatorname{tr} (PC^2)}{\operatorname{tr} (PCA) \operatorname{tr} (PBC)} - 1 \right| \leq \frac{1}{4} \cdot \frac{(M - m)(N - n)}{\sqrt{mnMN}}.$$

If A has the trace P - (k, K) -property and B has the trace P - (l, L) -property, then

$$(4.9) \quad \left| \frac{\operatorname{tr} (PBA) \operatorname{tr} (P)}{\operatorname{tr} (PA) \operatorname{tr} (PB)} - 1 \right| \leq \frac{1}{4} \cdot \frac{(K - k)(L - l)}{\sqrt{klKL}},$$

where $kK, lL > 0$.

We observe that, if $0 < k1_H \leq A \leq K1_H$ and $0 < l1_H \leq B \leq L1_H$, then by (4.10) we have

$$(4.10) \quad |\operatorname{tr}(PBA) \operatorname{tr}(P) - \operatorname{tr}(PA) \operatorname{tr}(PB)| \leq \frac{1}{4} \cdot \frac{(K-k)(L-l)}{\sqrt{k l K L}} \operatorname{tr}(PA) \operatorname{tr}(PB)$$

or, equivalently

$$(4.11) \quad \left| \frac{\operatorname{tr}(PBA)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(PB)}{\operatorname{tr}(P)} \right| \leq \frac{1}{4} \cdot \frac{(K-k)(L-l)}{\sqrt{k l K L}} \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(PB)}{\operatorname{tr}(P)}.$$

5. APPLICATIONS FOR CONVEX FUNCTIONS

In the paper [34] we obtained amongst other the following reverse of the Jensen trace inequality:

Let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If f is a continuously differentiable convex function on $[m, M]$ and $P \in \mathcal{B}_1(H) \setminus \{0\}$, $P \geq 0$, then we have

$$(5.1) \quad \begin{aligned} 0 &\leq \frac{\operatorname{tr}(Pf(A))}{\operatorname{tr}(P)} - f\left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)}\right) \\ &\leq \frac{\operatorname{tr}(Pf'(A)A)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \cdot \frac{\operatorname{tr}(Pf'(A))}{\operatorname{tr}(P)} \\ &\leq \begin{cases} \frac{1}{2} [f'(M) - f'(m)] \frac{\operatorname{tr}\left(P\left|A - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} 1_H\right|\right)}{\operatorname{tr}(P)} \\ \frac{1}{2} (M - m) \frac{\operatorname{tr}\left(P\left|f'(A) - \frac{\operatorname{tr}(Pf'(A))}{\operatorname{tr}(P)} 1_H\right|\right)}{\operatorname{tr}(P)} \end{cases} \\ &\leq \begin{cases} \frac{1}{2} [f'(M) - f'(m)] \left[\frac{\operatorname{tr}(PA^2)}{\operatorname{tr}(P)} - \left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)}\right)^2 \right]^{1/2} \\ \frac{1}{2} (M - m) \left[\frac{\operatorname{tr}(P[f'(A)]^2)}{\operatorname{tr}(P)} - \left(\frac{\operatorname{tr}(Pf'(A))}{\operatorname{tr}(P)}\right)^2 \right]^{1/2} \end{cases} \\ &\leq \frac{1}{4} [f'(M) - f'(m)] (M - m). \end{aligned}$$

Let $\mathcal{M}_n(\mathbb{C})$ be the space of all square matrices of order n with complex elements and $A \in \mathcal{M}_n(\mathbb{C})$ be a Hermitian matrix such that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If f is a continuously differentiable convex function on $[m, M]$,

then by taking $P = I_n$ in (5.1) we get

$$\begin{aligned}
(5.2) \quad 0 &\leq \frac{\operatorname{tr}(f(A))}{n} - f\left(\frac{\operatorname{tr}(A)}{n}\right) \\
&\leq \frac{\operatorname{tr}(f'(A)A)}{n} - \frac{\operatorname{tr}(A)}{n} \cdot \frac{\operatorname{tr}(f'(A))}{n} \\
&\leq \begin{cases} \frac{1}{2} [f'(M) - f'(m)] \frac{\operatorname{tr}\left(\left|A - \frac{\operatorname{tr}(A)}{n} 1_H\right|\right)}{n} \\ \frac{1}{2} (M - m) \frac{\operatorname{tr}\left(\left|f'(A) - \frac{\operatorname{tr}(f'(A))}{n} 1_H\right|\right)}{n} \end{cases} \\
&\leq \begin{cases} \frac{1}{2} [f'(M) - f'(m)] \left[\frac{\operatorname{tr}(A^2)}{n} - \left(\frac{\operatorname{tr}(A)}{n}\right)^2 \right]^{1/2} \\ \frac{1}{2} (M - m) \left[\frac{\operatorname{tr}([f'(A)]^2)}{n} - \left(\frac{\operatorname{tr}(f'(A))}{n}\right)^2 \right]^{1/2} \end{cases} \\
&\leq \frac{1}{4} [f'(M) - f'(m)] (M - m).
\end{aligned}$$

The following reverse inequality also holds:

Proposition 2. *Let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $0 < m < M$. If f is a continuously differentiable convex function on $[m, M]$ with $f'(m) > 0$ and $P \in \mathcal{B}_1(H) \setminus \{0\}$, $P \geq 0$, then we have*

$$\begin{aligned}
(5.3) \quad 0 &\leq \frac{\operatorname{tr}(Pf(A))}{\operatorname{tr}(P)} - f\left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)}\right) \\
&\leq \frac{\operatorname{tr}(Pf'(A)A)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \cdot \frac{\operatorname{tr}(Pf'(A))}{\operatorname{tr}(P)} \\
&\leq \frac{1}{4} \cdot \frac{(M - m) [f'(M) - f'(m)] \operatorname{tr}(PA) \operatorname{tr}(Pf'(A))}{\sqrt{mMf'(m)f'(M)} \operatorname{tr}(P) \operatorname{tr}(P)}.
\end{aligned}$$

The proof follows by the inequality (4.11) and the details are omitted.

Let $A \in \mathcal{M}_n(\mathbb{C})$ be a Hermitian matrix such that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If f is a continuously differentiable convex function on $[m, M]$ with $f'(m) > 0$ then by taking $P = I_n$ in (5.3) we get

$$\begin{aligned}
(5.4) \quad 0 &\leq \frac{\operatorname{tr}(f(A))}{n} - f\left(\frac{\operatorname{tr}(A)}{n}\right) \\
&\leq \frac{\operatorname{tr}(f'(A)A)}{n} - \frac{\operatorname{tr}(A)}{n} \cdot \frac{\operatorname{tr}(f'(A))}{n} \\
&\leq \frac{1}{4} \cdot \frac{(M - m) [f'(M) - f'(m)] \operatorname{tr}(A) \operatorname{tr}(f'(A))}{\sqrt{mMf'(m)f'(M)} \operatorname{tr}(A) \operatorname{tr}(f'(A))}.
\end{aligned}$$

We consider the power function $f : (0, \infty) \rightarrow (0, \infty)$, $f(t) = t^r$ with $t \in \mathbb{R} \setminus \{0\}$. For $r \in (-\infty, 0) \cup [1, \infty)$, f is convex while for $r \in (0, 1)$, f is concave.

Let $r \geq 1$ and A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $0 < m < M$. If $P \in \mathcal{B}_1^+(H) \setminus \{0\}$,

then

$$\begin{aligned}
 (5.5) \quad 0 &\leq \frac{\operatorname{tr}(PA^r)}{\operatorname{tr}(P)} - \left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right)^r \\
 &\leq r \left[\frac{\operatorname{tr}(PA^r)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \cdot \frac{\operatorname{tr}(PA^{r-1})}{\operatorname{tr}(P)} \right] \\
 &\leq \frac{1}{4^p} \frac{(M-m)(M^{p-1}-m^{p-1})}{m^{p/2}M^{p/2}} \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(PA^{p-1})}{\operatorname{tr}(P)}.
 \end{aligned}$$

If we take the first and last term in (5.5) we get the inequality:

$$\begin{aligned}
 (5.6) \quad 0 &\leq \frac{\operatorname{tr}(P)\operatorname{tr}(PA^r)}{\operatorname{tr}(PA)\operatorname{tr}(PA^{p-1})} - \frac{\operatorname{tr}(P)[\operatorname{tr}(PA)]^{r-1}}{\operatorname{tr}(PA^{p-1})[\operatorname{tr}(P)]^{r-1}} \\
 &\leq \frac{1}{4^p} \frac{(M-m)(M^{p-1}-m^{p-1})}{m^{p/2}M^{p/2}}.
 \end{aligned}$$

Consider the convex function $f : \mathbb{R} \rightarrow (0, \infty)$, $f(t) = \exp t$ and let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $0 < m < M$. If $P \in \mathcal{B}_1^+(H) \setminus \{0\}$, then using (5.3) we have

$$\begin{aligned}
 (5.7) \quad 0 &\leq \frac{\operatorname{tr}(P \exp A)}{\operatorname{tr}(P)} - \exp \left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right) \\
 &\leq \frac{\operatorname{tr}(PA \exp A)}{\operatorname{tr}(P)} - \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \cdot \frac{\operatorname{tr}(P \exp A)}{\operatorname{tr}(P)} \\
 &\leq \frac{1}{4} \cdot \frac{(M-m)(\exp M - \exp m)}{\sqrt{mM} \exp(m+M)} \frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \frac{\operatorname{tr}(P \exp A)}{\operatorname{tr}(P)}.
 \end{aligned}$$

If we take the first and last term in (5.7) we get the inequality:

$$\begin{aligned}
 (5.8) \quad 0 &\leq \frac{\operatorname{tr}(P)}{\operatorname{tr}(PA)} - \frac{[\operatorname{tr}(P)]^2 \exp \left(\frac{\operatorname{tr}(PA)}{\operatorname{tr}(P)} \right)}{\operatorname{tr}(PA)\operatorname{tr}(P \exp A)} \\
 &\leq \frac{1}{4} \cdot \frac{(M-m)(\exp M - \exp m)}{\sqrt{mM} \exp(m+M)}.
 \end{aligned}$$

REFERENCES

- [1] G. A. Anastassiou, Grüss type inequalities for the Stieltjes integral. *Nonlinear Funct. Anal. Appl.* **12** (2007), no. 4, 583–593.
- [2] G. A. Anastassiou, Chebyshev-Grüss type and comparison of integral means inequalities for the Stieltjes integral. *Panamer. Math. J.* **17** (2007), no. 3, 91–109.
- [3] G. A. Anastassiou, Chebyshev-Grüss type inequalities via Euler type and Fink identities. *Math. Comput. Modelling* **45** (2007), no. 9-10, 1189–1200
- [4] T. Ando, Matrix Young inequalities, *Oper. Theory Adv. Appl.* **75** (1995), 33–38.
- [5] R. Bellman, Some inequalities for positive definite matrices, in: E.F. Beckenbach (Ed.), *General Inequalities 2*, Proceedings of the 2nd International Conference on General Inequalities, Birkhäuser, Basel, 1980, pp. 89–90.
- [6] E. V. Belmega, M. Jungers and S. Lasaulce, A generalization of a trace inequality for positive definite matrices. *Aust. J. Math. Anal. Appl.* **7** (2010), no. 2, Art. 26, 5 pp.
- [7] N. G. de Bruijn, Problem 12, *Wisk. Opgaven*, **21** (1960), 12-14.
- [8] P. Cerone, On some results involving the Čebyšev functional and its generalisations. *J. Inequal. Pure Appl. Math.* **4** (2003), no. 3, Article 55, 17 pp.
- [9] P. Cerone, On Chebyshev functional bounds. *Differential & difference equations and applications*, 267–277, Hindawi Publ. Corp., New York, 2006.

- [10] P. Cerone, On a Čebyšev-type functional and Grüss-like bounds. *Math. Inequal. Appl.* **9** (2006), no. 1, 87–102.
- [11] P. Cerone and S. S. Dragomir, A refinement of the Grüss inequality and applications. *Tamkang J. Math.* **38** (2007), no. 1, 37–49.
- [12] P. Cerone and S. S. Dragomir, New bounds for the Čebyšev functional. *Appl. Math. Lett.* **18** (2005), no. 6, 603–611.
- [13] P. Cerone and S. S. Dragomir, Chebychev functional bounds using Ostrowski seminorms. *Southeast Asian Bull. Math.* **28** (2004), no. 2, 219–228.
- [14] D. Chang, A matrix trace inequality for products of Hermitian matrices, *J. Math. Anal. Appl.* **237** (1999) 721–725.
- [15] L. Chen and C. Wong, Inequalities for singular values and traces, *Linear Algebra Appl.* **171** (1992), 109–120.
- [16] I. D. Coop, On matrix trace inequalities and related topics for products of Hermitian matrix, *J. Math. Anal. Appl.* **188** (1994) 999–1001.
- [17] J. B. Diaz and F. T. Metcalf, Stronger forms of a class of inequalities of G. Pólya-G. Szegő and L.V. Kantorovich, *Bull. Amer. Math. Soc.*, **69** (1963), 415–418.
- [18] S. S. Dragomir, Some Grüss type inequalities in inner product spaces, *J. Inequal. Pure & Appl. Math.*, **4**(2003), No. 2, Article 42, [ON LINE: http://jipam.vu.edu.au/v4n2/032_03.html].
- [19] S. S. Dragomir, *A Survey on Cauchy-Bunyakovsky-Schwarz Type Discrete Inequalities*, RGMIA Monographs, Victoria University, 2002. (ON LINE: <http://rgmia.vu.edu.au/monographs/>).
- [20] S. S. Dragomir, A counterpart of Schwarz’s inequality in inner product spaces, *East Asian Math. J.*, **20**(1) (2004), 1-10. Zbl 1094.26017. Preprint *RGMIA Res. Rep. Coll.*, **6**(2003), Supplement, Article 18. [ON LINE: [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)].
- [21] S. S. Dragomir, Grüss inequality in inner product spaces, *The Australian Math Soc. Gazette*, **26** (1999), No. 2, 66-70.
- [22] S. S. Dragomir, A generalization of Grüss’ inequality in inner product spaces and applications, *J. Math. Anal. Appl.*, **237** (1999), 74-82.
- [23] S. S. Dragomir, Some discrete inequalities of Grüss type and applications in guessing theory, *Honam Math. J.*, **21**(1) (1999), 145-156.
- [24] S. S. Dragomir, Some integral inequalities of Grüss type, *Indian J. of Pure and Appl. Math.*, **31**(4) (2000), 397-415.
- [25] S. S. Dragomir, *Advances in Inequalities of the Schwarz, Grüss and Bessel Type in Inner Product Spaces*, Nova Science Publishers Inc., New York, 2005, x+249 pp.
- [26] S. S. Dragomir and G.L. Booth, On a Grüss-Lupaş type inequality and its applications for the estimation of p-moments of guessing mappings, *Mathematical Communications*, **5** (2000), 117-126.
- [27] S. S. Dragomir, A Grüss type integral inequality for mappings of r -Hölder’s type and applications for trapezoid formula, *Tamkang J. of Math.*, **31**(1) (2000), 43-47.
- [28] S. S. Dragomir and I. Fedotov, An inequality of Grüss’ type for Riemann-Stieltjes integral and applications for special means, *Tamkang J. of Math.*, **29**(4)(1998), 286-292.
- [29] S. S. Dragomir, Čebyšev’s type inequalities for functions of selfadjoint operators in Hilbert spaces, *Linear Multilinear Algebra* **58** (2010), no. 7-8, 805–814. Preprint *RGMIA Res. Rep. Coll.*, **11**(2008), Supp. Art. 9. [Online [http://rgmia.org/v11\(E\).php](http://rgmia.org/v11(E).php)]
- [30] S. S. Dragomir, Grüss’ type inequalities for functions of selfadjoint operators in Hilbert spaces, *Ital. J. Pure Appl. Math.* No. **28** (2011), 207–224. Preprint *RGMIA Res. Rep. Coll.* **11** (2008), Supp. Art. 11. [Online [http://rgmia.org/v11\(E\).php](http://rgmia.org/v11(E).php)]
- [31] S. S. Dragomir, New inequalities of the Kantorovich type for bounded linear operators in Hilbert spaces. *Linear Algebra Appl.* **428** (2008), no. 11-12, 2750–2760.
- [32] S. S. Dragomir, Some Čebyšev’s type trace inequalities for functions of selfadjoint operators in Hilbert spaces, Preprint *RGMIA Res. Rep. Coll.* **17** (2014), Art. 111. [Online <http://rgmia.org/papers/v17/v17a111.pdf>].
- [33] S. S. Dragomir, Some Grüss type inequalities for trace of operators in Hilbert spaces, Preprint *RGMIA Res. Rep. Coll.* **17** (2014), Art. 114. [Online <http://rgmia.org/papers/v17/v17a114.pdf>].
- [34] S. S. Dragomir, Reverse Jensen’s type trace inequalities for convex functions of selfadjoint operators in Hilbert spaces, Preprint *RGMIA Res. Rep. Coll.* **17** (2014), Art. 118. [Online <http://rgmia.org/papers/v17/v17a118.pdf>].

- [35] A. M. Fink, A treatise on Grüss' inequality, *Analytic and Geometric Inequalities*, 93-113, Math. Appl. 478, Kluwer Academic Publ., 1999.
- [36] S. Furuichi and M. Lin, Refinements of the trace inequality of Belmega, Lasaulce and Debbah. *Aust. J. Math. Anal. Appl.* **7** (2010), no. 2, Art. 23, 4 pp.
- [37] T. Furuta, J. Mičić Hot, J. Pečarić and Y. Seo, *Mond-Pečarić Method in Operator Inequalities. Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Element, Zagreb, 2005.
- [38] W. Greub and W. Rheinboldt, On a generalisation of an inequality of L.V. Kantorovich, *Proc. Amer. Math. Soc.*, **10** (1959), 407-415.
- [39] G. Grüss, Über das Maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx$, *Math. Z.*, **39**(1935), 215-226.
- [40] M. S. Klamkin and R. G. McLenaghan, An ellipse inequality, *Math. Mag.*, **50** (1977), 261-263.
- [41] H. D. Lee, On some matrix inequalities, *Korean J. Math.* **16** (2008), No. 4, pp. 565-571.
- [42] L. Liu, A trace class operator inequality, *J. Math. Anal. Appl.* **328** (2007) 1484-1486.
- [43] Z. Liu, Refinement of an inequality of Grüss type for Riemann-Stieltjes integral, *Soochow J. Math.*, **30**(4) (2004), 483-489.
- [44] S. Manjegani, Hölder and Young inequalities for the trace of operators, *Positivity* **11** (2007), 239-250.
- [45] A. Matković, J. Pečarić and I. Perić, A variant of Jensen's inequality of Mercer's type for operators with applications. *Linear Algebra Appl.* **418** (2006), no. 2-3, 551-564.
- [46] D. S. Mitrinović, J. E. Pečarić and A.M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [47] H. Neudecker, A matrix trace inequality, *J. Math. Anal. Appl.* **166** (1992) 302-303.
- [48] N. Ozeki, On the estimation of the inequality by the maximum, *J. College Arts, Chiba Univ.*, **5**(2) (1968), 199-203.
- [49] B. G. Pachpatte, A note on Grüss type inequalities via Cauchy's mean value theorem. *Math. Inequal. Appl.* **11** (2008), no. 1, 75-80.
- [50] J. Pečarić, J. Mičić and Y. Seo, Inequalities between operator means based on the Mond-Pečarić method. *Houston J. Math.* **30** (2004), no. 1, 191-207
- [51] G. Pólya and G. Szegő, *Aufgaben und Lehrsätze aus der Analysis*, Vol. 1, Berlin 1925, pp. 57 and 213-214.
- [52] M. B. Ruskai, Inequalities for traces on von Neumann algebras, *Commun. Math. Phys.* **26**(1972), 280-289.
- [53] K. Shebrawi and H. Albadawi, Operator norm inequalities of Minkowski type, *J. Inequal. Pure Appl. Math.* **9**(1) (2008), 1-10, article 26.
- [54] K. Shebrawi and H. Albadawi, Trace inequalities for matrices, *Bull. Aust. Math. Soc.* **87** (2013), 139-148.
- [55] O. Shisha and B. Mond, Bounds on differences of means, *Inequalities I*, New York-London, 1967, 293-308.
- [56] B. Simon, *Trace Ideals and Their Applications*, Cambridge University Press, Cambridge, 1979.
- [57] Z. Ulukök and R. Türkmen, On some matrix trace inequalities. *J. Inequal. Appl.* **2010**, Art. ID 201486, 8 pp.
- [58] G. S. Watson, Serial correlation in regression analysis I, *Biometrika*, **42** (1955), 327-342.
- [59] X. Yang, A matrix trace inequality, *J. Math. Anal. Appl.* **250** (2000) 372-374.
- [60] X. M. Yang, X. Q. Yang and K. L. Teo, A matrix trace inequality, *J. Math. Anal. Appl.* **263** (2001), 327-331.
- [61] Y. Yang, A matrix trace inequality, *J. Math. Anal. Appl.* **133** (1988) 573-574.
- [62] C.-J. Zhao and W.-S. Cheung, On multivariate Grüss inequalities. *J. Inequal. Appl.* **2008**, Art. ID 249438, 8 pp.

¹MATHEMATICS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.org/dragomir>

²SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA