

**NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR  
 $n$ -TIMES DIFFERENTIABLE CONVEX AND CONCAVE  
FUNCTIONS WITH APPLICATIONS**

M. A. LATIF AND S. S. DRAGOMIR<sup>1,2</sup>

ABSTRACT. In this paper, a new identity for  $n$ -times differentiable functions is established and by using the obtained identity, some new inequalities Hermite-Hadamard type are obtained for functions whose  $n$ th derivatives in absolute value are convex and concave functions. From our results, several inequalities of Hermite-Hadamard type can be derived in terms of functions whose first and second derivatives in absolute value are convex and concave functions as special cases. Our results may provide refinements of some results already exist in literature. Applications to trapezoidal formula and special means of established results are given.

1. INTRODUCTION

A function  $f : I \rightarrow \mathbb{R}$ ,  $\emptyset \neq I \subseteq \mathbb{R}$ , is said to be convex on  $I$  if the inequality

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y),$$

holds for all  $x, y \in I$  and  $t \in [0, 1]$ .

Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex mapping and  $a, b \in I$  with  $a < b$ . Then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}.$$

The double inequality (1.1) is known as the Hermite-Hadamard inequality (see [8]). The inequalities (1.1) hold in reversed direction if  $f$  is concave.

A number of papers have been written to extend, to generalize and to improve the inequality (1.1). Many papers can be found in literature to answer an important question of estimating the difference between the middle and the rightmost terms in (1.1) see for example [1, 2], [4, 5], [7], [9, 10, 11], [19], [21] and [22] and the references therein.

Dragomir and Agarwal [4], proved the following results to estimate the difference between the middle and the rightmost terms in (1.1).

**Theorem 1.** [4, Theorem 2.2] *Let  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I^\circ$  with  $a < b$ . If  $f' \in L([a, b])$  and  $|f'|$  is convex on  $[a, b]$ . Then*

$$(1.2) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{8} \left[ |f'(a)| + |f'(b)| \right].$$

---

*Date:* Today.

*2000 Mathematics Subject Classification.* Primary 26D07; Secondary 26D15.

*Key words and phrases.* Hermite-Hadamard's inequality, convex function, differentiable function, Hölder integral inequality.

This paper is in final form and no version of it will be submitted for publication elsewhere.

**Theorem 2.** [4, Theorem 2.3] *Let  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I^\circ$  with  $a < b$ . If  $f' \in L([a, b])$  and  $|f'|^{p/(p-1)}$  is convex on  $[a, b]$  for  $p > 1$ . Then*

$$(1.3) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{2(p+1)^{1/p}} \left[ \frac{|f'(a)|^{p/(p-1)} + |f'(b)|^{p/(p-1)}}{2} \right]^{(p-1)/p}.$$

Pearce and Pečarić [19], improved the inequality (1.3) as the following result.

**Theorem 3.** [19, Theorem 1] *Let  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I^\circ$  with  $a < b$ . If  $f' \in L([a, b])$  and  $|f'|^q$  is convex on  $[a, b]$  for  $q \geq 1$ . Then*

$$(1.4) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{4} \left[ \frac{|f'(a)|^q + |f'(b)|^q}{2} \right]^{1/q}.$$

Most recently, Hwang [9] established the following results for functions whose  $n$ th derivatives in absolute value are convex and concave functions.

**Theorem 4.** [9] *Suppose  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$ ,  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)}$  exists on  $I^\circ$ ,  $f^{(n)} \in L([a, b])$  and  $|f^{(n)}|^q$  is convex on  $[a, b]$  for  $q \geq 1$ ,  $n \in \mathbb{N}$ ,  $n \geq 2$ . Then*

$$(1.5) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=2}^{n-1} \frac{(k-1)(b-a)^k}{2(k+1)!} f^{(k)}(a) \right| = \frac{(n-1)^{1-\frac{1}{q}}(b-a)^n}{2(n+1)!} \left[ \frac{(n^2-2)|f^{(n)}(a)|^q + n|f^{(n)}(b)|^q}{n+2} \right]^{\frac{1}{q}}.$$

**Theorem 5.** [9] *Suppose  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$ ,  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)}$  exists on  $I^\circ$ ,  $f^{(n)} \in L([a, b])$  and  $|f^{(n)}|^q$  is concave on  $[a, b]$  for  $q \geq 1$ ,  $n \in \mathbb{N}$ ,  $n \geq 2$ . Then*

$$(1.6) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=2}^{n-1} \frac{(k-1)(b-a)^k}{2(k+1)!} f^{(k)}(a) \right| = \frac{(n-1)(b-a)^n}{2(n+1)!} \left| f^{(n)} \left( \frac{(n^2-2)a + nb}{(n-1)(n+2)} \right) \right|.$$

Following results can be derived directly from Theorem 4 and Theorem 5 for  $n = 2$ .

**Corollary 1.** [9] *Under the assumptions of Theorem 4, for  $n = 2$ , we have*

$$(1.7) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| = \frac{(b-a)^2}{12} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}.$$

**Corollary 2.** [9] *Under the assumptions of Theorem 4, for  $n = 2$ , we have*

$$(1.8) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| = \frac{(b-a)^2}{12} \left| f'' \left( \frac{a+b}{2} \right) \right|.$$

It can be noticed that the inequalities (1.7) and (1.8) may be better than those given in Theorem 1, Theorem 2 and Theorem 3.

For more recent results on Hermite-Hadamard type inequalities for functions whose  $n$ th derivatives in absolute value are  $s$ -convex functions and  $m$ -convex, we refer the interested reader to the works of Jiang *et. al* [11] and Wang *et al.* [21] and the references therein.

The main purpose of the present paper is to establish new Hermite-Hadamard type inequalities for functions whose  $n$ th derivatives in absolute value are convex and concave. We believe that the results presented in this paper are better than those established in [9] and hence better than those given in [4] and [19]. Applications of our results to trapezoidal formula and to special means are given in Section 3 and Section 4.

## 2. MAIN RESULTS

The following Lemma is essential in establishing our main results in this section:

**Lemma 1.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  for  $n \in \mathbb{N}$  and  $f^{(n)} \in L([a, b])$ , where  $a, b \in I^\circ$  with  $a < b$ , we have the identity*

$$(2.1) \quad \begin{aligned} & \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \\ &= \frac{(b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)} \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt \\ &+ \frac{(-1)^n (b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)} \left( \frac{1-t}{2} b + \frac{1+t}{2} a \right) dt, \end{aligned}$$

where an empty sum is understood to be nil.

*Proof.* Suppose

$$I_n = \frac{(b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)} \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt$$

and

$$J_n = \frac{(-1)^n (b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)} \left( \frac{1-t}{2} b + \frac{1+t}{2} a \right) dt.$$

For  $n = 1$ , we have

$$I_1 = \frac{b-a}{4} \int_0^1 t f' \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt$$

and

$$J_1 = \frac{(-1)(b-a)}{4} \int_0^1 t f' \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt.$$

By integration by parts and using the substitution  $x = \frac{1-t}{2}a + \frac{1+t}{2}b$  for  $I_1$  and  $x = \frac{1+t}{2}a + \frac{1-t}{2}b$  for  $J_1$ , we obtain

$$I_1 = \frac{1}{2}f(b) - \frac{1}{b-a} \int_{\frac{a+b}{2}}^b f(x)$$

and

$$J_1 = \frac{1}{2}f(a) - \frac{1}{b-a} \int_a^{\frac{a+b}{2}} f(x).$$

Hence

$$I_1 + J_1 = \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx,$$

which coincides with the L.H.S of (2.1) for  $n = 1$ .

Similarly for  $n = 2$ , and using similar arguments as above, we have

$$I_2 + J_2 = \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx$$

which coincides with the L.H.S of (2.1) for  $n = 2$ .

Suppose (2.1) holds for  $n = m - 1 \geq 3$ .

Now for  $n = m$ , we have

$$\begin{aligned} & \frac{(b-a)^m}{2^{m+1}m!} \int_0^1 (1-t)^{m-1} (m-1+t) f^{(m)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \\ & + \frac{(-1)^m (b-a)^m}{2^{m+1}m!} \int_0^1 (1-t)^{m-1} (m-1+t) f^{(m)} \left( \frac{1-t}{2}b + \frac{1+t}{2}a \right) dt \\ & = - \frac{(b-a)^{m-1} (m-1) [1 + (-1)^{m-1}]}{2^m m!} f^{(m-1)} \left( \frac{a+b}{2} \right) \\ & + \frac{(b-a)^{m-1}}{2^m (m-1)!} \int_0^1 (1-t)^{m-2} (m-2+t) f^{(m)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \\ & + \frac{(-1)^{m-1} (b-a)^{m-1}}{2^m (m-1)!} \int_0^1 (1-t)^{m-2} (m-2+t) f^{(m)} \left( \frac{1-t}{2}b + \frac{1+t}{2}a \right) dt \\ & = \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{m-2} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \\ & \quad - \frac{(b-a)^{m-1} (m-1) [1 + (-1)^{m-1}]}{2^m m!} f^{(m-1)} \left( \frac{a+b}{2} \right) \\ & = \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{m-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right). \end{aligned}$$

This completes the proof of the lemma.  $\square$

Now we state and prove some new Hermite-Hadamard type inequalities for functions whose  $n$ th derivatives in absolute value are convex and concave.

**Theorem 6.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  and  $f^{(n)} \in L([a, b])$  for  $n \in \mathbb{N}$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f^{(n)}|^q$  is convex on  $[a, b]$  for  $q \in [1, \infty)$ , we have the inequality

$$(2.2) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n n^{1-1/q}}{2^{n+\frac{1}{q}+1} (n+1)!} \left\{ \left( \frac{(n^2+n-1) |f^{(n)}(a)|^q + (n^2+3n+1) |f^{(n)}(b)|^q}{n+2} \right)^{1/q} \right. \\ \left. + \left( \frac{(n^2+n-1) |f^{(n)}(b)|^q + (n^2+3n+1) |f^{(n)}(a)|^q}{n+2} \right)^{1/q} \right\}.$$

*Proof.* From Lemma 1, the Hölder inequality and the convexity of  $|f^{(n)}|^q$  on  $[a, b]$ , we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n}{2^{n+1} n!} \left( \int_0^1 (1-t)^{n-1} (n-1+t) dt \right)^{1-1/q} \\ \times \left\{ \left( \int_0^1 (1-t)^{n-1} (n-1+t) \left[ \left(\frac{1-t}{2}\right) |f^{(n)}(a)|^q + \left(\frac{1+t}{2}\right) |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\ \left. + \left( \int_0^1 (1-t)^{n-1} (n-1+t) \left[ \left(\frac{1-t}{2}\right) |f^{(n)}(b)|^q + \left(\frac{1+t}{2}\right) |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\} \\ = \frac{(b-a)^n n^{1-1/q}}{2^{n+\frac{1}{q}+1} (n+1)!} \left\{ \left( \frac{n^2+n-1}{n+2} |f^{(n)}(a)|^q + \frac{n^2+3n+1}{n+2} |f^{(n)}(b)|^q \right)^{1/q} \right. \\ \left. + \left( \frac{n^2+n-1}{n+2} |f^{(n)}(b)|^q + \frac{n^2+3n+1}{n+2} |f^{(n)}(a)|^q \right)^{1/q} \right\}.$$

Which is the desired result. This completes the proof of the Theorem.  $\square$

**Corollary 3.** *Under the assumptions of Theorem 6, if  $q = 1$ , we have the inequality*

$$(2.3) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \leq \frac{n(b-a)^n}{2^{n+1} (n+1)!} \left[ |f^{(n)}(a)| + |f^{(n)}(b)| \right].$$

**Corollary 4.** *Under the assumptions of Theorem 6, if  $n = 1$ , we have the inequality*

$$(2.4) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{8} \left\{ \left( \frac{|f'(a)|^q + 5|f'(b)|^q}{6} \right)^{1/q} + \left( \frac{|f'(b)|^q + 5|f'(a)|^q}{6} \right)^{1/q} \right\}.$$

**Corollary 5.** *If we take  $q = 1$  in Corollary 4, we have*

$$(2.5) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{8} \left[ |f'(a)| + |f'(b)| \right].$$

**Corollary 6.** *Suppose the assumptions of Theorem 6 are fulfilled and if  $n = 2$ , we have*

$$(2.6) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{24} \left\{ \left( \frac{5|f''(a)|^q + 11|f''(b)|^q}{16} \right)^{1/q} + \left( \frac{5|f''(b)|^q + 11|f''(a)|^q}{16} \right)^{1/q} \right\}.$$

**Corollary 7.** *If  $q = 1$  in Corollary 6, we have*

$$(2.7) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{24} \left[ |f''(a)| + |f''(b)| \right].$$

**Remark 1.** *It is obvious that the inequality (2.7) gives better estimate than that of (1.7) for  $q = 1$ .*

**Theorem 7.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  and  $f^{(n)} \in L([a, b])$  for  $n \in \mathbb{N}$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f^{(n)}|^q$  is convex on  $[a, b]$*

for  $q \in (1, \infty)$ , we have the inequality

$$\begin{aligned}
 (2.8) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\
 & \quad \left. - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \\
 & \leq \frac{(b-a)^n \left[ n^{(2q-1)/(q-1)} - (n-1)^{(2q-1)/(q-1)} \right]^{1-1/q}}{2^{n+\frac{1}{q}+1} n!} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \\
 & \quad \times \left\{ \left[ P |f^{(n)}(a)|^q + Q |f^{(n)}(b)|^q \right]^{1/q} + \left[ P |f^{(n)}(b)|^q + Q |f^{(n)}(a)|^q \right]^{1/q} \right\},
 \end{aligned}$$

where

$$P = \frac{1}{nq - q + 2} \quad \text{and} \quad Q = \frac{nq - q + 3}{(nq - q + 1)(nq - q + 2)}.$$

*Proof.* Using Lemma 1, the Hölder inequality and the convexity of  $|f^{(n)}|^q$  on  $[a, b]$ , we have

$$\begin{aligned}
 (2.9) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\
 & \quad \left. - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \\
 & \leq \frac{(b-a)^n}{2^{n+1} n!} \left( \int_0^1 (n-1+t)^{q/(q-1)} dt \right)^{1-1/q} \\
 & \quad \times \left\{ \left( \int_0^1 (1-t)^{q(n-1)} \left[ \left( \frac{1-t}{2} \right) |f^{(n)}(a)|^q + \left( \frac{1+t}{2} \right) |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\
 & \quad \left. + \left( \int_0^1 (1-t)^{q(n-1)} \left[ \left( \frac{1-t}{2} \right) |f^{(n)}(b)|^q + \left( \frac{1+t}{2} \right) |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.
 \end{aligned}$$

From (2.9) we get the inequality (2.8), since

$$\int_0^1 (n-1+t)^{q/(q-1)} dt = \left( \frac{q-1}{2q-1} \right) \left[ n^{(2q-1)/(q-1)} - (n-1)^{(2q-1)/(q-1)} \right],$$

$$\int_0^1 (1-t)^{q(n-1)+1} dt = \frac{1}{nq - q + 2} = P$$

and

$$\int_0^1 (1-t)^{q(n-1)} (1+t) dt = \frac{nq - q + 3}{(nq - q + 1)(nq - q + 2)} = Q.$$

□

**Corollary 8.** *Suppose the conditions of Theorem 7 are satisfied and if  $n = 1$ , we have the inequality*

$$(2.10) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{4} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \\ \times \left\{ \left( \frac{|f'(a)|^q + 3|f'(b)|^q}{4} \right)^{1/q} + \left( \frac{|f'(b)|^q + 3|f'(a)|^q}{4} \right)^{1/q} \right\},$$

**Corollary 9.** *Under the assumptions of Theorem 7, if  $n = 2$ , we have the inequality*

$$(2.11) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2 [2^{(2q-1)/(q-1)} - 1]^{1-1/q}}{2^{1/q+4}} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \\ \times \left\{ [P|f''(a)|^q + Q|f''(b)|^q]^{1/q} + [P|f''(b)|^q + Q|f''(a)|^q]^{1/q} \right\},$$

where

$$P = \frac{1}{q+2} \text{ and } Q = \frac{q+3}{(q+1)(q+2)}.$$

**Theorem 8.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  and  $f^{(n)} \in L([a, b])$  for  $n \in \mathbb{N}$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f^{(n)}|^q$  is convex on  $[a, b]$  for  $q \in (1, \infty)$ , we have the inequality*

$$(2.12) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ \left. - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \leq \frac{(b-a)^n}{2^{n+\frac{1}{q}+1} n!} \left( \frac{q-1}{nq-1} \right)^{1-1/q} \\ \times \left\{ [R|f^{(n)}(a)|^q + S|f^{(n)}(b)|^q]^{1/q} + [R|f^{(n)}(b)|^q + S|f^{(n)}(a)|^q]^{1/q} \right\},$$

$$R = \frac{n^{q+2} - (n+q+1)(n-1)^{q+1}}{(q+1)(q+2)}$$

and

$$S = \frac{(2q-n+4)n^{q+1} - (q-n+3)(n-1)^{q+1}}{(q+1)(q+2)}.$$



*Proof.* Using Lemma 1, the Hölder inequality and the convexity of  $|f^{(n)}|^q$  on  $[a, b]$ , we have

$$(2.13) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \\ \leq \frac{(b-a)^n}{2^{n+1} n!} \left( \int_0^1 (1-t)^{q(n-1)/(q-1)} dt \right)^{1-1/q} \\ \times \left\{ \left( \int_0^1 (n-1+t)^q \left[ \left( \frac{1-t}{2} \right) |f^{(n)}(a)|^q + \left( \frac{1+t}{2} \right) |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\ \left. + \left( \int_0^1 (n-1+t)^q \left[ \left( \frac{1-t}{2} \right) |f^{(n)}(b)|^q + \left( \frac{1+t}{2} \right) |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.$$

Since

$$\int_0^1 (1-t)^{q(n-1)/(q-1)} dt = \frac{q-1}{nq-1},$$

$$\int_0^1 (n-1+t)^q (1-t) dt = \frac{n^{q+2} - (n+q+1)(n-1)^{q+1}}{(q+1)(q+2)}$$

and

$$\int_0^1 (n-1+t)^q (1+t) dt = \frac{(2q-n+4)n^{q+1} - (q-n+3)(n-1)^{q+1}}{(q+1)(q+2)}$$

and hence (2.12) follows from (2.13).  $\square$

**Corollary 10.** *Suppose the assumptions of Theorem 8 are satisfied and if  $n = 1$ , we have the inequality*

$$(2.14) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)}{2^{2+1/q}} \left\{ \left[ R |f'(a)|^q + S |f'(b)|^q \right]^{1/q} + \left[ R |f'(b)|^q + S |f'(a)|^q \right]^{1/q} \right\},$$

$$R = \frac{1}{(q+1)(q+2)} \text{ and } S = \frac{2q+3}{(q+1)(q+2)}.$$

**Corollary 11.** *Suppose the assumptions of Theorem 8 are satisfied and if  $n = 2$ , we have the inequality*

$$(2.15) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{2^{4+1/q}} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \\ \times \left\{ \left[ R |f''(a)|^q + S |f''(b)|^q \right]^{1/q} + \left[ R |f''(b)|^q + S |f''(a)|^q \right]^{1/q} \right\},$$

$$R = \frac{2^{q+2} - (q+3)}{(q+1)(q+2)} \text{ and } S = \frac{2^{q+2} - 1}{q+2}.$$

**Theorem 9.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  and  $f^{(n)} \in L([a, b])$  for  $n \in \mathbb{N}$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f^{(n)}|^q$  is convex on  $[a, b]$  for  $q \in (1, \infty)$ , we have the inequality

$$(2.16) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{n^{n+1-1/q} (b-a)^n}{2^{n+1+2/q} n!} \left[ B\left(\frac{1}{n}; \frac{nq-1}{q-1}, \frac{2q-1}{q-1}\right) \right]^{1-1/q} \\ \times \left\{ \left[ |f^{(n)}(a)|^q + 3 |f^{(n)}(b)|^q \right]^{1/q} + \left[ 3 |f^{(n)}(a)|^q + |f^{(n)}(b)|^q \right]^{1/q} \right\},$$

where

$$B(x; \alpha, \beta) = \int_0^x t^{\alpha-1} (1-t)^{1-\beta} dt, \quad 0 \leq x \leq 1, \alpha > 0, \beta > 0$$

is the incomplete beta function.

*Proof.* Using Lemma 1, the Hölder inequality and the convexity of  $|f^{(n)}|^q$  on  $[a, b]$ , we have

$$(2.17) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n}{2^{n+1} n!} \left( \int_0^1 (1-t)^{q(n-1)/(q-1)} (n-1+t)^{q/(q-1)} dt \right)^{1-1/q} \\ \times \left\{ \left( \int_0^1 \left[ \left(\frac{1-t}{2}\right) |f^{(n)}(a)|^q + \left(\frac{1+t}{2}\right) |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\ \left. + \left( \int_0^1 \left[ \left(\frac{1-t}{2}\right) |f^{(n)}(b)|^q + \left(\frac{1+t}{2}\right) |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.$$

The inequality (2.16) follows from the inequality (2.17) and using the fact that

$$\int_0^1 (1-t)^{q(n-1)/(q-1)} (n-1+t)^{q/(q-1)} dt = n^{\frac{nq+q-1}{q-1}} B\left(\frac{1}{n}; \frac{nq-1}{q-1}, \frac{2q-1}{q-1}\right).$$

This completes the proof of the theorem.  $\square$

**Corollary 12.** Suppose the assumptions of Theorem 9 are satisfied and if  $n = 1$ , the inequality (2.16) becomes the inequality (2.10), since for  $n = 1$ , we have

$$B\left(1; 1, \frac{2q-1}{q-1}\right) = B\left(1, \frac{2q-1}{q-1}\right) = \int_0^1 (1-t)^{\frac{2q-1}{q-1}-1} dt = \frac{q-1}{2q-1}.$$

**Corollary 13.** *Under the assumptions of Theorem 9 and  $n = 2$ , we have the following inequality*

$$(2.18) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2}{2} \left[ B\left(\frac{1}{2}; \frac{2q-1}{q-1}, \frac{2q-1}{q-1}\right) \right]^{1-1/q} \\ \times \left\{ \left[ \frac{|f''(a)|^q + 3|f''(b)|^q}{8} \right]^{1/q} + \left[ \frac{3|f''(a)|^q + |f''(b)|^q}{8} \right]^{1/q} \right\},$$

where  $B(x; \alpha, \beta)$  is as defined in Theorem 9.

A result for concave functions is given in the following theorem.

**Theorem 10.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f^{(n)}$  exists on  $I^\circ$  and  $f^{(n)} \in L([a, b])$  for  $n \in \mathbb{N}$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f^{(n)}|^q$  is convex on  $[a, b]$  for  $q \in [1, \infty)$ , we have the inequality*

$$(2.19) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ \left. - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{n(b-a)^n}{2^{n+1} (n+1)!} \left\{ \left| f^{(n)}\left(\frac{(n^2+n-1)a + (n^2+3n+1)b}{2n(n+2)}\right) \right| \right. \\ \left. + \left| f^{(n)}\left(\frac{(n^2+n-1)b + (n^2+3n+1)a}{2n(n+2)}\right) \right| \right\}.$$

*Proof.* By the concavity of  $|f^{(n)}|^q$  and the power-mean inequality, we have

$$\left| f^{(n)}(\lambda x + (1-\lambda)y) \right| \geq \lambda \left| f^{(n)}(x) \right| + (1-\lambda) \left| f^{(n)}(y) \right|$$

which shows that  $|f^{(n)}|$  is also concave.

Hence by using Lemma 1 and the Jensen integral inequality, we have

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ & \quad \left. - \sum_{k=1}^{n-1} \frac{k \left[ 1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left( \frac{a+b}{2} \right) \right| \\ & \leq \frac{(b-a)^n}{2^{n+1} n!} \left( \int_0^1 (1-t)^{n-1} (n-1+t) dt \right) \\ & \quad \times \left\{ \left| f^{(n)} \left( \frac{(1-t)^{n-1} (n-1+t) \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt}{\int_0^1 (1-t)^{n-1} (n-1+t) dt} \right) \right| \right. \\ & \quad \left. + \left| f^{(n)} \left( \frac{(1-t)^{n-1} (n-1+t) \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt}{\int_0^1 (1-t)^{n-1} (n-1+t) dt} \right) \right| \right\}. \end{aligned}$$

The above inequality gives the desired result.  $\square$

**Corollary 14.** *Suppose the assumptions of Theorem 10 are fulfilled and if  $n = 1$ , we have the inequality*

$$(2.20) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{8} \left[ \left| f' \left( \frac{a+5b}{6} \right) \right| + \left| f' \left( \frac{b+5a}{6} \right) \right| \right].$$

**Corollary 15.** *Suppose the assumptions of Theorem 10 are fulfilled and if  $n = 2$ , we have the inequality*

$$(2.21) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{24} \left[ \left| f'' \left( \frac{5a+11b}{16} \right) \right| + \left| f'' \left( \frac{5b+11a}{16} \right) \right| \right].$$

### 3. APPLICATIONS TO THE TRAPEZOIDAL FORMULA

Let  $d$  be a division of the interval  $[a, b]$ , i.e.  $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ , and consider the quadrature formula

$$\int_a^b f(x) dx = T(f, d) + E(f, d),$$

where

$$T(f, d) = \sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{f(x_i) + f(x_{i+1})}{2}$$

is the trapezoidal versions and  $E(f, d)$  is the associated error. Here, we derive some error estimates for the trapezoidal formula in terms of absolute values of the second derivative of  $f$  which may be better than those already exist in the literature.

**Theorem 11.** Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a twice differentiable function on  $I^\circ$  such that  $f'' \in L([a, b])$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f''|^q$  is convex on  $[a, b]$  for  $q \geq 1$ , then for every division  $d$  of  $[a, b]$ , we have

$$(3.1) \quad |E(f, d)| \leq \frac{1}{24} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left\{ \left( \frac{5 |f''(x_i)|^q + 11 |f''(x_{i+1})|^q}{16} \right)^{1/q} + \left( \frac{5 |f''(x_{i+1})|^q + 11 |f''(x_i)|^q}{16} \right)^{1/q} \right\}.$$

*Proof.* By applying Corollary 6 on the subinterval  $[x_i, x_{i+1}]$  ( $i = 0, 1, \dots, n-1$ ) of the division  $d$ , we have

$$(3.2) \quad |E(f, d)| = \left| \sum_{i=0}^{n-1} \left\{ (x_{i+1} - x_i) \frac{f(x_i) + f(x_{i+1})}{2} - \int_{x_i}^{x_{i+1}} f(x) dx \right\} \right| \\ \leq \sum_{i=0}^{n-1} (x_{i+1} - x_i) \left| \frac{f(x_i) + f(x_{i+1})}{2} - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right| \\ \leq \frac{1}{24} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left\{ \left( \frac{5}{16} |f''(x_i)|^q + \frac{11}{16} |f''(x_{i+1})|^q \right)^{1/q} + \left( \frac{5}{16} |f''(x_{i+1})|^q + \frac{11}{16} |f''(x_i)|^q \right)^{1/q} \right\}.$$

Hence the result is proved.  $\square$

**Corollary 16.** Under the assumptions of Theorem 11, for  $q = 1$ , we have

$$(3.3) \quad |E(f, d)| \leq \frac{1}{24} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left[ |f''(x_i)| + |f''(x_{i+1})| \right].$$

**Theorem 12.** Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function on  $I^\circ$  such that  $f'' \in L([a, b])$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f''|^q$  is convex on  $[a, b]$  for  $q > 1$ , then for every division  $d$  of  $[a, b]$ , we have

$$(3.4) \quad |E(f, d)| \leq \frac{[2^{(2q-1)/(q-1)} - 1]^{1-1/q}}{(2^{1/q}) \cdot 8} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \times \left\{ \left[ \frac{(q+1) |f''(x_i)|^q + (q+3) |f''(x_{i+1})|^q}{(q+1)(q+2)} \right]^{1/q} + \left[ \frac{(q+1) |f''(x_{i+1})|^q + (q+3) |f''(x_i)|^q}{(q+1)(q+2)} \right]^{1/q} \right\}.$$

*Proof.* It follows from Corollary 9.  $\square$

**Theorem 13.** Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a twice differentiable function on  $I^\circ$  such that  $f'' \in L([a, b])$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f''|^q$  is convex on  $[a, b]$  for  $q > 1$ , then for every division  $d$  of  $[a, b]$ , we have

$$(3.5) \quad |E(f, d)| \leq \frac{1}{2^{4+\frac{1}{q}}} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \times \left\{ \left[ \frac{(2^{q+2} - q - 3) |f''(x_i)|^q + (q+1)(2^{q+2} - 1) |f''(x_{i+1})|^q}{(q+1)(q+2)} \right]^{1/q} \right. \\ \left. \left[ \frac{(2^{q+2} - q - 3) |f''(x_{i+1})|^q + (q+1)(2^{q+2} - 1) |f''(x_i)|^q}{(q+1)(q+2)} \right]^{1/q} \right\}.$$

*Proof.* It is a direct consequence of Corollary 11.  $\square$

Error bound for trapezoidal formula when  $|f''|^q$  for  $q \geq 1$  is concave may be given as follows.

**Theorem 14.** Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a twice differentiable function on  $I^\circ$  such that  $f'' \in L([a, b])$ , where  $a, b \in I^\circ$  with  $a < b$ . If  $|f''|^q$  is concave on  $[a, b]$  for  $q \geq 1$ , then for every division  $d$  of  $[a, b]$ , we have

$$(3.6) \quad |E(f, d)| \\ \leq \frac{1}{24} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left[ \left| f'' \left( \frac{5x_i + 11x_{i+1}}{16} \right) \right| + \left| f'' \left( \frac{5x_{i+1} + 11x_i}{16} \right) \right| \right].$$

*Proof.* It can be proved easily by using Corollary 15.  $\square$

#### 4. APPLICATIONS TO THE SPECIAL MEANS

Now, we consider applications of our results to special means. We consider the means for positive real numbers  $a, b \in \mathbb{R}_+$ . We take

(1) The arithmetic mean:

$$A(a, b) = \frac{a+b}{2}, a, b > 0.$$

(2) The harmonic mean:

$$H(a, b) = \frac{2}{\frac{1}{a} + \frac{1}{b}}, a, b > 0.$$

(3) The logarithmic mean:

$$L(a, b) = \frac{b-a}{\ln b - \ln a}, a, b > 0, a \neq b.$$

(4) Generalized log-mean:

$$L_p(a, b) = \left[ \frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}}, a, b > 0, p \in \mathbb{R} \setminus \{-1, 0\}, a \neq b.$$

Now we apply our results to establish some inequalities for special means.

**Theorem 15.** For  $a, b \in \mathbb{R}_+$ ,  $a < b$  and  $p \in \mathbb{N}$ ,  $p > 1$ , we have

$$\left| A(a^{p+1}, b^{p+1}) - L_{p+1}^{p+1}(a^{p+1}, b^{p+1}) \right| \leq \frac{p(p+1)(b-a)^2}{12} A(a^{p-1}, b^{p-1}).$$

*Proof.* For the function  $f(x) = x^{p+1}$ ,  $x \in \mathbb{R}_+$  and  $p \in \mathbb{N}$ , then  $|f''(x)| = (p+1)px^{p-1}$  is a convex function on  $\mathbb{R}_+$ . Applying Corollary 7, we obtain the required result.  $\square$

**Theorem 16.** For  $a, b \in \mathbb{R}_+$ ,  $a < b$  and  $p, q \in \mathbb{N}$ ,  $p, q > 1$ , we have

$$(4.1) \quad \left| A(a^{p+1/q}, b^{p+1/q}) - L_{p+1/q}^{p+1/q}(a^{p+1/q}, b^{p+1/q}) \right| \\ \leq \frac{(b-a)^2 [2^{(2q-1)/(q-1)} - 1]^{1-1/q}}{16} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \left( n-1 + \frac{1}{q} \right) \left( n + \frac{1}{q} \right) \\ \times \left\{ \left[ \frac{A(a^{(p-2)q+1}, b^{(p-2)q+1})}{q+2} + \frac{b^{(p-2)q+1}}{(q+1)(q+2)} \right]^{1/q} \right. \\ \left. + \left[ \frac{A(a^{(p-2)q+1}, b^{(p-2)q+1})}{q+2} + \frac{a^{(p-2)q+1}}{(q+1)(q+2)} \right]^{1/q} \right\}.$$

*Proof.* Let  $f(x) = x^{p+1/q}$ ,  $x \in \mathbb{R}_+$ ,  $p, q \in \mathbb{N}$ ,  $p, q > 1$ , then

$$\left| f''(x) \right|^q = \left( p-1 + \frac{1}{q} \right)^q \left( p + \frac{1}{q} \right)^q x^{(p-2)q+1}$$

is convex on  $\mathbb{R}_+$  and the result follows directly from Corollary 9.  $\square$

**Theorem 17.** For  $a, b \in \mathbb{R}_+$ ,  $a < b$  and  $p, q \in \mathbb{N}$ ,  $p, q > 1$ , we have

$$(4.2) \quad \left| A(a^{p+1/q}, b^{p+1/q}) - L_{p+1/q}^{p+1/q}(a^{p+1/q}, b^{p+1/q}) \right| \\ \leq \frac{(b-a)^2}{2^{4+1/q}} \left( \frac{q-1}{2q-1} \right)^{1-1/q} \left( p-1 + \frac{1}{q} \right) \left( p + \frac{1}{q} \right) \\ \times \left\{ \left[ \frac{2^{q+2} - (q+3)}{(q+1)(q+2)} a^{(p-2)q+1} + \frac{2^{q+2} - 1}{q+2} b^{(p-2)q+1} \right]^{1/q} \right. \\ \left. + \left[ \frac{2^{q+2} - (q+3)}{(q+1)(q+2)} b^{(p-2)q+1} + \frac{2^{q+2} - 1}{q+2} a^{(p-2)q+1} \right]^{1/q} \right\}$$

*Proof.* Let  $f(x) = x^{p+1/q}$ ,  $x \in \mathbb{R}_+$ ,  $p, q \in \mathbb{N}$ ,  $p, q > 1$ , then

$$\left| f''(x) \right|^q = \left( p-1 + \frac{1}{q} \right)^q \left( p + \frac{1}{q} \right)^q x^{(p-2)q+1}$$

is convex on  $\mathbb{R}_+$  and the result follows directly from Corollary 11.  $\square$

**Theorem 18.** For  $a, b \in \mathbb{R}_+$ ,  $a < b$  and  $p, q \in \mathbb{N}$ ,  $p, q > 1$  with  $p < q$ , we have

$$(4.3) \quad \left| A(a^{p/q^2+2}, b^{p/q^2+2}) - L_{p/q^2+2}^{p/q^2+2}(a^{p/q^2+2}, b^{p/q^2+2}) \right| \\ \leq \frac{(b-a)^2}{12} \left( \frac{p}{q^2} + 2 \right) \left( \frac{p}{q^2} + 1 \right) A(a^{p/q^2}, b^{p/q^2}).$$

*Proof.* Let  $f(x) = x^{p/q^2+2}$ ,  $x \in \mathbb{R}_+$ , then  $|f''(x)| = \left(\frac{p}{q^2} + 2\right) \left(\frac{p}{q^2} + 1\right) x^{p/q^2}$ . It is obvious that  $|f''(x)|^q$  is concave on  $\mathbb{R}_+$ . Applying Corollary 15, we get the desired result.  $\square$

**Theorem 19.** For  $a, b \in \mathbb{R}_+$ ,  $a < b$ , we have

$$(4.4) \quad |H^{-1}(a, b) - L^{-1}(a, b)| \leq \frac{(b-a)^2}{6} A(a^{-3}, b^{-3})$$

*Proof.* Let  $f(x) = \frac{1}{x}$ ,  $x \in \mathbb{R}_+$ , then  $|f''(x)| = \frac{2}{x^3}$  is convex on  $\mathbb{R}_+$  so applying Corollary 7, we get the inequality (4.4).  $\square$

**Remark 2.** Many other interesting inequalities for means can be obtained by applying the other results to the function  $f(x) = \frac{1}{x}$ ,  $x \in \mathbb{R}_+$ , however the details are left to the interested reader.

#### REFERENCES

- [1] M. Alomari, M. Darus, S.S. Dragomir, Inequalities of Hermite Hadamard's type for functions whose derivatives absolute values are quasi-convex, RGMIA 12 (suppl. 14) (2009).
- [2] M. Alomari, M. Darus and U.S. Kirmaci, Refinements of Hadamard-type inequalities for quasi-convex functions with applications to trapezoidal formula and to special means, Comput. Math. Appl. 59 (2010) 225-232
- [3] S.S. Dragomir, Two mappings in connection to Hadamard's inequalities, J. Math. Anal. Appl. 167 (1992) 49-56.
- [4] S.S. Dragomir, R.P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, Appl. Math. Lett. 11 (1998) 91-95.
- [5] S.S. Dragomir, Y.J. Cho, S.S. Kim, Inequalities of Hadamard's type for Lipschitzian mappings and their applications, J. Math. Anal. Appl. 245 (2000) 489-501.
- [6] S.S. Dragomir, S. Wang, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and for some numerical quadrature rule, Appl. Math. Lett. 11 (1998) 105-109.
- [7] S.S. Dragomir, C.E.M. Pearce, Quasi-convex functions and Hadamard's inequality, Bull. Austr. Math. Soc. 57 (3) (1998) 377-385.
- [8] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, J. Math Pures Appl., 58 (1893), 171–215.
- [9] D.-Y. Hwang, Some Inequalities for  $n$ -time Differentiable Mappings and Applications, Kyungpook Math. J. 43(2003), 335-343
- [10] D.A. Ion, Some estimates on the Hermite Hadamard inequality through quasi-convex functions, Annals of University of Craiova, Math. Comp. Sci. Ser. 34 (2007) 82-87.
- [11] W.-D. Jiang, Da-Wei Niu, Yun Hua, and Feng Qi, Generalizations of Hermite-Hadamard inequality to  $n$ -time differentiable functions which are  $s$ -convex in the second sense, Analysis (Munich) 32 (2012), 1001–1012; Available online at <http://dx.doi.org/10.1524/anly.2012.1161>.
- [12] U.S. Kirmaci, Inequalities for differentiable mappings and applicatios to special means of real numbers to midpoint formula, Appl. Math. Comput. 147 (2004) 137-146.
- [13] U.S. Kirmaci, M.E. Özdemir, On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, Appl. Math. Comput. 153 (2004) 361-368.
- [14] M. A. Latif and S. S. Dragomir, New inequalities of Hermite-Hadamard type for functions whose derivatives in absolute value are convex with applications, Acta Univ. M. Belii, ser. Math. (2013), 24–39.
- [15] M. A. Latif and S. S. Dragomir, On Hermite-Hadamard type integral inequalities for  $n$ -times differentiable  $(\alpha, m)$ -logarithmically convex functions, RGMIA Research Report Collection, 17(2014), Article 14, 16 pp.



- [16] M. A. Latif and S. S. Dragomir, On Hermite-Hadamard type integral inequalities for  $n$ -times differentiable log-preinvex functions. (Accepted)
- [17] Generalization of Hermite-Hadamard type inequalities for  $n$ -times differentiable functions which are  $s$ -preinvex in the second sense with applications. (Accepted)
- [18] M.E. Özdemir, A theorem on mappings with bounded derivatives with applications to quadrature rules and means, *Appl. Math. Comput.* 138 (2003) 425-434.
- [19] C.E.M. Pearce, J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formula, *Appl. Math. Lett.* 13 (2000) 51-55.
- [20] C.E.M. Pearce, A.M. Rubinov,  $P$ -functions, quasiconvex functions and Hadamard-type inequalities, *J. Math. Anal. Appl.* 240 (1999) 92-104.
- [21] S.-H. Wang, Bo-Yan Xi and Feng Qi, Some new inequalities of Hermite-Hadamard type for  $n$ -times differentiable functions which are  $m$ -convex, *Analysis (Munich)* 32 (2012), no. 3, 247-262; Available online at <http://dx.doi.org/10.1524/anly.2012.1167>.
- [22] G.S. Yang, D.Y. Hwang, K.L. Tseng, Some inequalities for differentiable convex and concave mappings, *Comput. Math. Appl.* 47 (2004) 207-216

SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

*E-mail address:* `m.amer.latif@hotmail.com`

<sup>1</sup> SCHOOL OF ENGINEERING AND SCIENCE, VICTORIA UNIVERSITY, P. O. BOX 14428, MELBOURNE CITY, MC8001, AUSTRALIA

<sup>2</sup>SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

*E-mail address:* `sever.dragomir@vu.edu.au`