

Received 26/05/14

NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR n -TIMES DIFFERENTIABLE s -CONVEX FUNCTIONS WITH APPLICATIONS

M. A. LATIF, S. S. DRAGOMIR^{1,2}, AND E. MOMONIAT

ABSTRACT. In this paper, some new inequalities Hermite-Hadamard type are obtained for functions whose n th derivatives in absolute value are s -convex functions. From our results, several inequalities of Hermite-Hadamard type can be derived in terms of functions whose first and second derivatives in absolute value are s -convex functions as special cases. Our results may provide refinements of some results already exist in literature. Applications to trapezoidal rule and to special means of established results are given.

1. INTRODUCTION

A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if the inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

holds for all $x, y \in I$ and $t \in [0, 1]$.

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping and $a, b \in I$ with $a < b$. Then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

The double inequality (1.1) is known as the Hermite-Hadamard inequality (see [9]). The inequalities (1.1) hold in reversed direction if f is concave.

For recent results, refinements, counterparts, generalizations and new Hermite-Hadamard-type inequalities see [6, 7, 12, 15, 16, 17, 19, 21, 25, 29, 31] and the references therein.

In [9], Hudzik and Maligranda considered among others the class of functions which are s -convex in the second sense and is defined as follows.

Definition 1. Let $s \in (0, 1]$ be a fixed real number. A function $f : [0, \infty) \rightarrow [0, \infty)$ is said to be s -convex in the second sense, if

$$(1.2) \quad f(tx + (1-y)y) \leq t^s f(x) + (1-t)^s f(y)$$

holds for all $x, y \in [0, \infty)$ and $t \in [0, 1]$. The class of s -convex functions in the second sense is denoted by K_s^2 . If the inequality (1.2) holds in reversed direction, then f is to be an s -concave function in the second sense.

Date: Today.

2000 Mathematics Subject Classification. Primary 26D07; Secondary 26D15.

Key words and phrases. Hermite-Hadamard's inequality, convex function, s -convex, Hölder integral inequality, trapezoidal rule.

This paper is in final form and no version of it will be submitted for publication elsewhere.

It is clear that the definition of s -convexity (s -concavity) coincides with the definition of convexity (concavity) when $s = 1$.

In [5], Dragomir and Fitzpatrick proved a variant of Hadamard's inequality which holds for s -convex functions in the second sense.

Theorem 1. *Suppose that $f : [0, \infty) \rightarrow [0, \infty)$ is an s -convex function in the second sense, where $s \in (0, 1)$, $a, b \in [0, \infty)$ with $a < b$. If $a, b \in L([a, b])$, then the following inequalities hold*

$$(1.3) \quad 2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1}.$$

For more recent results on Hermite-Hadamard type inequalities for functions whose derivatives in absolute value are s -convex functions, we refer the interested reader to [1, 13, 14, 24] and the references therein.

The main purpose of the present paper is to establish new Hermite-Hadamard type inequalities for functions whose n th derivatives in absolute value are s -convex. We believe that the results presented in this paper are better than those already exist in the literature concerning the inequalities of Hermite-Hadamard type for s -convex functions. Applications of our results to trapezoidal formula and to special means are given in Section 3 and Section 4.

2. MAIN RESULTS

We will use the following Lemmas to establish our main results in this section.

Lemma 1. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f^{(n)}$ exists on I° and $f^{(n)} \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$, $n \in \mathbb{N}$, we have the identity*

$$(2.1) \quad \begin{aligned} & \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \\ &= \frac{(b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) dt \\ &+ \frac{(-1)^n (b-a)^n}{2^{n+1} n!} \int_0^1 (1-t)^{n-1} (n-1+t) f^{(n)}\left(\frac{1-t}{2}b + \frac{1+t}{2}a\right) dt, \end{aligned}$$

where an empty sum is understood to be nil.

Lemma 2. *Let $x \geq 0$, $y \geq 0$, the inequality*

$$(x+y)^\theta \leq x^\theta + y^\theta$$

holds for $0 < \theta \leq 1$ and the inequality

$$(x-y)^\theta \leq x^\theta - y^\theta$$

holds for $\theta \geq 1$.

Now we state and prove some new Hermite-Hadamard type inequalities for functions whose n th derivatives in absolute value are s -convex and s -concave.

Theorem 2. Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a function such that $f^{(n)}$ exists on I° and $f^{(n)} \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$, $n \in \mathbb{N}$. If $|f^{(n)}|^q$ is s -convex on $[a, b]$ for some fixed $s \in (0, 1]$ and $q \in [1, \infty)$, we have the inequality

$$(2.2) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left(\frac{a+b}{2} \right) \right| \leq \frac{(b-a)^n}{2^{n+s/q+1} n!} \left(\frac{n}{n+1} \right)^{1-1/q} \\ \times \left\{ \left(\frac{(n^2 + s(n-1)) |f^{(n)}(a)|^q}{(n+s)(n+s+1)} + \left(\frac{n}{n+1} + \frac{n(n+s)}{n+s+1} B(s+1, n) \right) |f^{(n)}(b)|^q \right)^{1/q} \right. \\ \left. \left(\frac{(n^2 + s(n-1)) |f^{(n)}(b)|^q}{(n+s)(n+s+1)} + \left(\frac{n}{n+1} + \frac{n(n+s)}{n+s+1} B(s+1, n) \right) |f^{(n)}(a)|^q \right)^{1/q} \right\},$$

where

$$B(\alpha, \beta) = \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt, \alpha, \beta > 0$$

is the Euler Beta function.

Proof. From Lemma 1, the Hölder inequality and s -convexity of $|f^{(n)}|^q$ on $[a, b]$, we have

$$(2.3) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left(\frac{a+b}{2} \right) \right| \\ \leq \frac{(b-a)^n}{2^{n+s/q+1} n!} \left(\int_0^1 (1-t)^{n-1} (n-1+t) dt \right)^{1-1/q} \\ \times \left\{ \left(\int_0^1 (1-t)^{n-1} (n-1+t) \left[(1-t)^s |f^{(n)}(a)|^q + (1+t)^s |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\ \left. + \left(\int_0^1 (1-t)^{n-1} (n-1+t) \left[(1-t)^s |f^{(n)}(b)|^q + (1+t)^s |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.$$

Since

$$\int_0^1 (1-t)^{n-1} (n-1+t) dt = \frac{n}{n+1}, \\ \int_0^1 (1-t)^{n-1} (n-1+t) (1-t)^s dt = \frac{n^2 + s(n-1)}{(n+s)(n+s+1)},$$

by using the property

$$B(x, y+1) = \frac{y}{x+y} B(x, y)$$

of the Euler Beta function and Lemma 2, we have

$$\begin{aligned}
& \int_0^1 (1-t)^{n-1} (n-1+t) (1+t)^s dt \\
&= \int_0^1 (1-t)^{n-1} (n-1+t) (1+t^s) dt \\
&= \frac{n}{n+1} + nB(s+1, n) - B(s+1, n+1) \\
&= \frac{n}{n+1} + nB(s+1, n) - \frac{n}{n+s+1} B(s+1, n) \\
&= \frac{n}{n+1} + \frac{n(n+s)}{n+s+1} B(s+1, n).
\end{aligned}$$

From the above facts and the inequality (2.3), we get the required inequality (2.2). This completes the proof of the Theorem. \square

The following corollaries are direct consequences of Theorem 2.

Corollary 1. *Under the assumptions of Theorem 2, if $q = 1$, we have the inequality*

$$\begin{aligned}
(2.4) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\
& \quad \left. - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left(\frac{a+b}{2} \right) \right| \leq \frac{(b-a)^n}{2^{n+s+1} n!} \\
& \times \left(\frac{n^2 + s(n-1)}{(n+s)(n+s+1)} + \frac{n}{n+1} + \frac{n(n+s)}{n+s+1} B(s+1, n) \right) \left[|f^{(n)}(a)| + |f^{(n)}(b)| \right],
\end{aligned}$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2.

Corollary 2. *Under the assumptions of Theorem 2, if $n = 1$, we have the inequality*

$$\begin{aligned}
(2.5) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
& \leq (b-a) \left(\frac{1}{2} \right)^{3+s/q-1/q} \left\{ \left[\frac{|f'(a)|^q}{(1+s)(2+s)} + \left(\frac{1}{2} + \frac{1}{2+s} \right) |f'(b)|^q \right]^{1/q} \right. \\
& \quad \left. + \left[\left(\frac{1}{2} + \frac{1}{2+s} \right) |f'(a)|^q + \frac{|f'(b)|^q}{(1+s)(2+s)} \right]^{1/q} \right\}.
\end{aligned}$$

Corollary 3. *If we take $q = 1$ in Corollary 2, we have*

$$\begin{aligned}
(2.6) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
& \leq \frac{(b-a)(s+3)}{s+1} \left(\frac{1}{2} \right)^{3+s} \left[|f'(a)| + |f'(b)| \right].
\end{aligned}$$

Corollary 4. *Suppose the assumptions of Theorem 2 are fulfilled and if $n = 2$, we have*

$$(2.7) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{2^{4+s/q}} \left(\frac{2}{3}\right)^{1-1/q} \\ \times \left\{ \left[\frac{(s+4)}{(s+2)(s+3)} |f''(a)|^q + \left(\frac{2}{3} + \frac{2}{(s+1)(s+3)}\right) |f''(b)|^q \right]^{1/q} \right. \\ \left. + \left[\frac{(s+4)}{(s+2)(s+3)} |f''(b)|^q + \left(\frac{2}{3} + \frac{2}{(s+1)(s+3)}\right) |f''(a)|^q \right]^{1/q} \right\}.$$

Corollary 5. *If $q = 1$ in Corollary 4, we have*

$$(2.8) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2}{2^{4+s}} \left(\frac{2}{3} + \frac{s^2 + 7s + 8}{(s+1)(s+2)(s+3)}\right) [|f''(a)| + |f''(b)|].$$

Theorem 3. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be an n -times differentiable function on I° , $n \in \mathbb{N}$. If $f^{(n)} \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f^{(n)}|^q$ is convex on $[a, b]$ for $q \in (1, \infty)$, we have the inequality*

$$(2.9) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ \left. - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n \left[n^{(2q-1)/(q-1)} - (n-1)^{(2q-1)/(q-1)} \right]^{1-1/q}}{2^{n+s/q+1} n!} \left(\frac{q-1}{2q-1}\right)^{1-1/q} \\ \times \left\{ \left[\frac{|f^{(n)}(a)|^q}{nq - q + s + 1} + \left(\frac{1}{nq - q + 1} + B(s+1, nq - q + 1)\right) |f^{(n)}(b)|^q \right]^{1/q} \right. \\ \left. + \left[\frac{|f^{(n)}(b)|^q}{nq - q + s + 1} + \left(\frac{1}{nq - q + 1} + B(s+1, nq - q + 1)\right) |f^{(n)}(a)|^q \right]^{1/q} \right\}.$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2.

Proof. Using Lemma 1, by using the first inequality in Lemma 2, the Hölder inequality and convexity of $|f^{(n)}|^q$ on $[a, b]$, we have

$$\begin{aligned}
(2.10) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\
& \quad \left. - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left(\frac{a+b}{2} \right) \right| \\
& \leq \frac{(b-a)^n}{2^{n+s/q+1} n!} \left(\int_0^1 (n-1+t)^{q/(q-1)} dt \right)^{1-1/q} \\
& \quad \times \left\{ \left(\int_0^1 (1-t)^{q(n-1)} \left[(1-t)^s |f^{(n)}(a)|^q + (1+t)^s |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\
& \quad \left. + \left(\int_0^1 (1-t)^{q(n-1)} \left[(1-t)^s |f^{(n)}(b)|^q + (1+t)^s |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.
\end{aligned}$$

By simple computations, we observe that

$$\int_0^1 (n-1+t)^{q/(q-1)} dt = \left(\frac{q-1}{2q-1} \right) \left[n^{(2q-1)/(q-1)} - (n-1)^{(2q-1)/(q-1)} \right],$$

$$\int_0^1 (1-t)^{q(n-1)+s} dt = \frac{1}{nq - q + s + 1}$$

and

$$\int_0^1 (1-t)^{q(n-1)} (1+t)^s dt = \frac{1}{nq - q + 1} + B(s+1, nq - q + 1).$$

Using the above results in (2.10), we obtain the required result. This completes the proof of the theorem. \square

Corollary 6. *Suppose the assumptions of Theorem 3 are satisfied and if $n = 1$, we have the inequality*

$$\begin{aligned}
(2.11) \quad & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{2^{2+s/q}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \left(\frac{1}{s+1} \right)^{1/q} \\
& \quad \left\{ \left[|f'(a)|^q + (s+2) |f'(b)|^q \right]^{1/q} + \left[|f'(b)|^q + (s+2) |f'(a)|^q \right]^{1/q} \right\}.
\end{aligned}$$

Corollary 7. *Under the assumptions of Theorem 3, if $n = 2$, we have the inequality*

$$(2.12) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2 [2^{(2q-1)/(q-1)} - 1]^{1-1/q}}{2^{4+s/q}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \\ \times \left\{ \left[\frac{1}{q+s+1} |f''(a)|^q + \left(\frac{1}{q+1} + B(s+1, q+1) \right) |f''(b)|^q \right]^{1/q} \right. \\ \left. + \left[\frac{1}{q+s+1} |f''(b)|^q + \left(\frac{1}{q+1} + B(s+1, q+1) \right) |f''(a)|^q \right]^{1/q} \right\},$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2.

Theorem 4. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be an n -times differentiable function on I° , $n \in \mathbb{N}$. If $f^{(n)} \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f^{(n)}|^q$ is convex on $[a, b]$ for $q \in (1, \infty)$, we have the inequality*

$$(2.13) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ \left. - \sum_{k=1}^{n-1} \frac{k [1 + (-1)^k] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)} \left(\frac{a+b}{2} \right) \right| \\ \leq \frac{(b-a)^n}{2^{n+\frac{s}{q}+1} n!} \left(\frac{q-1}{nq-1} \right)^{1-1/q} \left\{ \left[P |f^{(n)}(a)|^q + Q |f^{(n)}(b)|^q \right]^{1/q} \right. \\ \left. + \left[P |f^{(n)}(b)|^q + Q |f^{(n)}(a)|^q \right]^{1/q} \right\},$$

where

$$P = n^{q+s+1} B\left(\frac{1}{n}; s+1, q+1\right),$$

$$Q = n^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - B(s+1, q+1),$$

$B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2 and

$$B(x; \alpha, \beta) = \int_0^x t^{\alpha-1} (1-t)^{\beta-1} dt, \alpha, \beta > 0, 0 \leq x \leq 1$$

is the incomplete Beta function.

Proof. Using Lemma 1, the first inequality in Lemma 2, the Hölder inequality and convexity of $|f^{(n)}|^q$ on $[a, b]$, we have

$$(2.14) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n}{2^{n+s/q+1} n!} \left(\int_0^1 (1-t)^{q(n-1)/(q-1)} dt \right)^{1-1/q} \\ \times \left\{ \left(\int_0^1 (n-1+t)^q \left[(1-t)^s |f^{(n)}(a)|^q + (1+t^s) |f^{(n)}(b)|^q \right] dt \right)^{1/q} \right. \\ \left. + \left(\int_0^1 (n-1+t)^q \left[(1-t)^s |f^{(n)}(b)|^q + (1+t^s) |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.$$

By using the second inequality of Lemma 2 and simple computation, it is easy to observe that

$$\int_0^1 (1-t)^{q(n-1)/(q-1)} dt = \frac{q-1}{nq-1}, \\ \int_0^1 (n-1+t)^q (1-t)^s dt = n^{q+s+1} \int_0^{\frac{1}{n}} t^s (1-t)^q dt \\ = n^{q+s+1} B\left(\frac{1}{n}; s+1, q+1\right) = P$$

and

$$\int_0^1 (n-1+t)^q (1+t^s) dt \leq \int_0^1 (n^q - (1-t)^q) (1+t^s) dt \\ = n^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - B(s+1, q+1) = Q.$$

Hence (2.13) follows from (2.14) and using the above results. This completes the proof of the theorem. \square

Corollary 8. *Suppose the assumptions of Theorem 4 are satisfied and if $n = 1$, we have the inequality*

$$(2.15) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{2^{2+s/q}} \\ \times \left\{ \left[B(s+1, q+1) |f'(a)|^q + \left(\frac{s+2}{s+1} - \frac{1}{q+1} - B(s+1, q+1) \right) |f'(b)|^q \right]^{1/q} \right. \\ \left. + \left[B(s+1, q+1) |f'(b)|^q + \left(\frac{s+2}{s+1} - \frac{1}{q+1} - B(s+1, q+1) \right) |f'(a)|^q \right]^{1/q} \right\},$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2.

Corollary 9. *Suppose the assumptions of Theorem 4 are satisfied and if $n = 2$, we have the inequality*

$$(2.16) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2}{2^{4+\frac{s}{q}}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \left\{ \left[2^{q+s+1} B\left(\frac{1}{2}; s+1, q+1\right) |f''(a)|^q \right. \right. \\ \left. \left. + \left(2^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - B(s+1, q+1) \right) |f''(b)|^q \right]^{1/q} \right. \\ \left. + \left[2^{q+s+1} B\left(\frac{1}{2}; s+1, q+1\right) |f''(b)|^q \right. \right. \\ \left. \left. + \left(2^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - 2^{q+s+1} B(s+1, q+1) \right) |f''(a)|^q \right]^{1/q} \right\},$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2 and $B(x; \alpha, \beta)$ is the incomplete Beta function defined as in Theorem 4.

A different approach results in the following theorem.

Theorem 5. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f^{(n)}$ exists on I° and $f^{(n)} \in L([a, b])$ for $n \in \mathbb{N}$, where $a, b \in I^\circ$ with $a < b$. If $|f^{(n)}|^q$ is convex on $[a, b]$ for $q \in (1, \infty)$, we have the inequality*

$$(2.17) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right. \\ \left. - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k \right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{n^{n+1-1/q} (b-a)^n}{2^{n+s/q+1} n!} \left(\frac{1}{s+1} \right)^{1/q} \left[B\left(\frac{1}{n}; \frac{nq-1}{q-1}, \frac{2q-1}{q-1}\right) \right]^{1-1/q} \\ \times \left\{ \left[|f^{(n)}(a)|^q + (2^{s+1} - 1) |f^{(n)}(b)|^q \right]^{1/q} \right. \\ \left. + \left[(2^{s+1} - 1) |f^{(n)}(a)|^q + |f^{(n)}(b)|^q \right]^{1/q} \right\},$$

where

$$B(x; \alpha, \beta) = \int_0^x t^{\alpha-1} (1-t)^{1-\beta} dt, \quad 0 \leq x \leq 1, \alpha > 0, \beta > 0$$

is the incomplete beta function.

Proof. Using Lemma 1, the Hölder inequality and the convexity of $|f^{(n)}|^q$ on $[a, b]$, we have

$$(2.18) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx - \sum_{k=1}^{n-1} \frac{k \left[1 + (-1)^k\right] (b-a)^k}{2^{k+1} (k+1)!} f^{(k)}\left(\frac{a+b}{2}\right) \right| \\ \leq \frac{(b-a)^n}{2^{n+s/q+1} n!} \left(\int_0^1 (1-t)^{q(n-1)/(q-1)} (n-1+t)^{q/(q-1)} dt \right)^{1-1/q} \\ \times \left\{ \left(\int_0^1 \left[(1-t)^s |f^{(n)}(a)|^q + (1+t)^s |f^{(n)}(b)|^q \right] dt \right)^{1/q} + \left(\int_0^1 \left[(1-t)^s |f^{(n)}(b)|^q + (1+t)^s |f^{(n)}(a)|^q \right] dt \right)^{1/q} \right\}.$$

By simple computations of integrals, the inequality (2.17) follows from the inequality (2.18) and using the fact that

$$\int_0^1 (1-t)^{q(n-1)/(q-1)} (n-1+t)^{q/(q-1)} dt = n^{\frac{nq+q-1}{q-1}} B\left(\frac{1}{n}; \frac{nq-1}{q-1}, \frac{2q-1}{q-1}\right).$$

This completes the proof of the theorem. \square

Corollary 10. *Suppose the assumptions of Theorem 5 are satisfied and if $n = 1$, we have*

$$(2.19) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)}{2^{2+s/q}} \left(\frac{1}{s+1}\right)^{1/q} \left(\frac{q-1}{2q-1}\right)^{1-1/q} \\ \times \left\{ \left[|f'(a)|^q + (2^{s+1}-1) |f'(b)|^q \right]^{1/q} + \left[(2^{s+1}-1) |f'(a)|^q + |f'(b)|^q \right]^{1/q} \right\}.$$

Proof. Proof follows from the fact that

$$B\left(1; 1, \frac{2q-1}{q-1}\right) = B\left(1, \frac{2q-1}{q-1}\right) = \int_0^1 (1-t)^{\frac{2q-1}{q-1}-1} dt = \frac{q-1}{2q-1}.$$

\square

Corollary 11. *Under the assumptions of Theorem 5 and $n = 2$, we have the following inequality*

$$(2.20) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ \leq \frac{(b-a)^2}{2^{1+s/q+1/q}} \left(\frac{1}{s+1}\right)^{1/q} \left[B\left(\frac{1}{2}; \frac{2q-1}{q-1}, \frac{2q-1}{q-1}\right) \right]^{1-1/q} \\ \times \left\{ \left[|f''(a)|^q + (2^{s+1}-1) |f''(b)|^q \right]^{1/q} + \left[(2^{s+1}-1) |f''(a)|^q + |f''(b)|^q \right]^{1/q} \right\},$$

where $B(x; \alpha, \beta)$ is defined as in Theorem 5.

3. APPLICATIONS TO THE TRAPEZOIDAL FORMULA

Let d be a division of the interval $[a, b]$, i.e. $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$, and consider the quadrature formula

$$\int_a^b f(x)dx = T(f, d) + E(f, d),$$

where

$$T(f, d) = \sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{f(x_i) + f(x_{i+1})}{2}$$

is the trapezoidal versions and $E(f, d)$ is the associated error. Here, we derive some error estimates for the trapezoidal formula in terms of absolute values of the second derivative of f which may be better than those already exist in the literature.

Theorem 6. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f'' \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f''|^q$ is s -convex on $[a, b]$ for $q \geq 1$, then for every division d of $[a, b]$, we have*

$$(3.1) \quad |E(f, d)| \leq \frac{1}{2^{4+s/q}} \left(\frac{2}{3}\right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \times \left\{ \left[\frac{(s+4)}{(s+2)(s+3)} |f''(x_i)|^q + \left(\frac{2}{3} + \frac{2}{(s+1)(s+3)}\right) |f''(x_{i+1})|^q \right]^{1/q} \right. \\ \left. + \left[\frac{(s+4)}{(s+2)(s+3)} |f''(x_{i+1})|^q + \left(\frac{2}{3} + \frac{2}{(s+1)(s+3)}\right) |f''(x_i)|^q \right]^{1/q} \right\}.$$

Proof. By applying Corollary 4 on the subinterval $[x_i, x_{i+1}]$ ($i = 0, 1, \dots, n-1$) of the division d , we have

$$(3.2) \quad |E(f, d)| = \left| \sum_{i=0}^{n-1} \left\{ (x_{i+1} - x_i) \frac{f(x_i) + f(x_{i+1})}{2} - \int_{x_i}^{x_{i+1}} f(x) dx \right\} \right| \\ \leq \sum_{i=0}^{n-1} (x_{i+1} - x_i) \left| \frac{f(x_i) + f(x_{i+1})}{2} - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right| \\ \leq \frac{1}{2^{4+s/q}} \left(\frac{2}{3}\right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \times \left\{ \left[\frac{(s+4)}{(s+2)(s+3)} |f''(x_i)|^q + \left(\frac{2}{3} + \frac{2}{(s+1)(s+3)}\right) |f''(x_{i+1})|^q \right]^{1/q} \right. \\ \left. + \left[\frac{(s+4)}{(s+2)(s+3)} |f''(x_{i+1})|^q + \left(\frac{2}{3} + \frac{2}{s+1}\right) |f''(x_i)|^q \right]^{1/q} \right\}.$$

□

Corollary 12. *Under the assumptions of Theorem 6, for $q = 1$, we have*

$$(3.3) \quad |E(f, d)| \leq \frac{1}{2^{4+s}} \left(\frac{2}{3} + \frac{s^2 + 7s + 8}{(s+1)(s+2)(s+3)} \right) \\ \times \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left[|f''(x_i)| + |f''(x_{i+1})| \right].$$

Theorem 7. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f'' \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f''|^q$ is s -convex on $[a, b]$ for $q > 1$, then for every division d of $[a, b]$, we have*

$$(3.4) \quad |E(f, d)| \leq \frac{[2^{(2q-1)/(q-1)} - 1]^{1-1/q}}{2^{4+s/q}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \times \left\{ \left[\frac{1}{q+s+1} |f''(x_i)|^q + \left(\frac{1}{q+1} + B(s+1, q+1) \right) |f''(x_{i+1})|^q \right]^{1/q} \right. \\ \left. + \left[\frac{1}{q+s+1} |f''(x_{i+1})|^q + \left(\frac{1}{q+1} + B(s+1, q+1) \right) |f''(x_i)|^q \right]^{1/q} \right\},$$

where $B(\alpha, \beta)$ is the Euler Beta function defined as in Theorem 2.

Proof. It follows from Corollary 7. □

Theorem 8. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f'' \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f''|^q$ is s -convex on $[a, b]$ for $q > 1$, then for every division d of $[a, b]$, we have*

$$(3.5) \quad |E(f, d)| \leq \frac{1}{2^{4+\frac{1}{q}}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \\ \left\{ \left[2^{q+s+1} B\left(\frac{1}{2}; s+1, q+1\right) |f''(x_i)|^q \right. \right. \\ \left. \left. + \left(2^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - B(s+1, q+1) \right) |f''(x_{i+1})|^q \right]^{1/q} \right. \\ \left. + \left[2^{q+s+1} B\left(\frac{1}{2}; s+1, q+1\right) |f''(x_{i+1})|^q \right. \right. \\ \left. \left. + \left(2^q \left(\frac{s+2}{s+1} \right) - \frac{1}{q+1} - 2^{q+s+1} B(s+1, q+1) \right) |f''(x_i)|^q \right]^{1/q} \right\},$$

where $B(\alpha, \beta)$ is the Euler Beta functions and $B(x; \alpha, \beta)$ is the incomplete beta function defined as in Theorem 4.

Proof. It is a direct consequence of Corollary 9. □

Theorem 9. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f'' \in L([a, b])$, where $a, b \in I^\circ$ with $a < b$. If $|f''|^q$ is s -convex on $[a, b]$ for $q > 1$,*

then for every division d of $[a, b]$, we have

$$(3.6) \quad |E(f, d)| \leq \frac{1}{2^{1+s/q+1/q}} \left(\frac{1}{s+1} \right)^{1/q} \left[B \left(\frac{1}{2}; \frac{2q-1}{q-1}, \frac{2q-1}{q-1} \right) \right]^{1-1/q} \\ \times \sum_{i=0}^{n-1} (x_{i+1} - x_i)^3 \left\{ \left[|f''(x_i)|^q + (2^{s+1} - 1) |f''(x_{i+1})|^q \right]^{1/q} \right. \\ \left. + \left[(2^{s+1} - 1) |f''(x_i)|^q + |f''(x_{i+1})|^q \right]^{1/q} \right\},$$

where $B(x; \alpha, \beta)$ is the incomplete beta function defined as in Theorem 4.

Proof. The proof follows by using Corollary 9. \square

4. APPLICATIONS TO THE SPECIAL MEANS

Now, we consider applications of our results to special means. We consider the means for positive real numbers $a, b \in \mathbb{R}_+$. We consider

(1) The arithmetic mean:

$$A(a, b) = \frac{a+b}{2}; a, b > 0.$$

(2) Generalized log-mean:

$$L_p(a, b) = \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}}; a, b > 0, p \in \mathbb{R} \setminus \{-1, 0\}, a \neq b.$$

Now we apply our results from Section 2 to give some inequalities for special means.

It is shown in [20] that the function $f : [0, \infty) \rightarrow \mathbb{R}$ defined by

$$f(t) = \begin{cases} a, & t = 0, \\ bt^s + c, & t > 0, \end{cases}$$

where $s \in (0, 1)$, $a, b, c \in \mathbb{R}$ with $0 \leq c \leq a$, $b \geq 0$, is s -convex on $[0, \infty)$. Hence, for $a = c = 0$, $b \geq 0$, the function $f(t) = bt^s$ is an s -convex function on $[0, \infty)$, where $s \in (0, 1)$.

Theorem 10. For $a, b \in \mathbb{R}_+$, $a < b$, $0 < s < 1$ and $q \in [1, \infty)$, we have

$$(4.1) \quad \left| A(a^{s+2}, b^{s+2}) - L_{s+2}^{s+2}(a^{s+2}, b^{s+2}) \right| \\ \leq \frac{(b-a)^2}{2^{3+s}} \left(\frac{2}{3}(s+2)(s+1) + \frac{s^2+7s+8}{s+3} \right) A(a^s, b^s).$$

Proof. Let $f(x) = x^{s+2}$, $x \in \mathbb{R}_+$. Then $|f''(x)| = (s+2)(s+1)x^s$ is an s -convex function on \mathbb{R}_+ . Applying Corollary 5, we obtain the required result. \square

Theorem 11. For $a, b \in \mathbb{R}_+$, $a < b$ and $q \in (1, \infty)$, we have

$$(4.2) \quad \left| A \left(a^{s/q+1}, b^{s/q+1} \right) - L_{s/q+1}^{s/q+1} \left(a^{s/q+1}, b^{s/q+1} \right) \right| \\ \leq \frac{(b-a)}{2^{2+s/q}} \left(\frac{q-1}{2q-1} \right)^{1-1/q} \left(\frac{1}{s+1} \right)^{1/q} (s/q+1) \\ \times \left\{ [2A(a^s, b^s) + (s+1)b^s]^{1/q} + [2A(a^s, b^s) + (s+1)a^s]^{1/q} \right\}.$$

Proof. Let $f(x) = x^{s/q+1}$, $x \in \mathbb{R}_+$. Then $|f'(x)|^q = [(s/q+1)]^q x^s$ is an s -convex function on \mathbb{R}_+ . Applying Corollary 6, we obtain the required result. \square

Theorem 12. For $a, b \in \mathbb{R}_+$, $a < b$ and $n, q \in \mathbb{N}$, $n, q > 1$, we have

$$(4.3) \quad \left| A \left(a^{s/q+1}, b^{s/q+1} \right) - L_{s/q+1}^{s/q+1} \left(a^{s/q+1}, b^{s/q+1} \right) \right| \\ \leq \frac{(b-a)}{2^{2+s/q-1/q}} \left(\frac{1}{s+1} \right)^{1/q} \left(\frac{q-1}{2q-1} \right)^{1-1/q} (s/q+1) \\ \times \left\{ [A(a^s, b^s) + (2^s-1)b^s]^{1/q} + [(2^s-1)a^s + A(a^s, b^s)]^{1/q} \right\}.$$

Proof. Let $f(x) = x^{s/q+1}$, $x \in \mathbb{R}_+$. Then $|f'(x)|^q = [(s/q+1)]^q x^s$ is an s -convex function on \mathbb{R}_+ . Applying Corollary 10, we obtain the required result. \square

Remark 1. Many other interesting inequalities for means can be obtained by applying the other results to some suitable s -convex functions, however the details are left to the interested reader.

REFERENCES

- [1] M. W. Alomari, M. Darus and U. S. Kirmaci, Some inequalities of Hadamard type inequalities for s -convex, Acts Math. Sci. Ser. B Engl. Ed. 31 (2011), no. 4, 1643-1652.
- [2] A. O. Akdemir and M. Tunç, On some inegral inequalities for s -logarithmically convex and their applications, arXiv:1212.1584 [math.FA].
- [3] R.-F. Bai, F. Qi and B.-Y. Xi, Hermite-Hadamard type inequalities for the m - and (α, m) -logarithmically convex functions, Filomat 27:1 (2013), 1-7.
- [4] S. S. Dragomir, On some new inequalities of Hermite-Hadamard type for m -convex functions, Tamkang J. Math. 33 (2002) 45-55.
- [5] S. S. Dragomir and S. Fitzpatric, The Hadamard inequalities for s -convex functions in the second sense, Demonstratio Math. 32 (1999), no. 4 687-696.
- [6] S. S. Dragomir and C. E. M. Pearce, Selected Topics on Hermite-Hadamard Inequalities and Applications, RGMIA Monographs, Victoria University, 2000.
- [7] S. S. Dragomir, R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, Appl. Math. Lett. 11 (1998) 91-95.
- [8] J. Deng and J. R. Wang, Fractional Hermite-Hadamard inequalities for (α, m) -logarithmically convex functions, J Inequal Appl 2013, 2013:364.
- [9] H. Hudzik, L. Maligranda, Some remarks on s -convex functions, Aequationes Math. 48 (1994) 100-111
- [10] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, J. Math Pures Appl., 58 (1893), 171-215.
- [11] Ch. Hermite, Sur deux limites d'une intégrale définie, Mathesis 3 (1883), 82.
- [12] D.-Y. Hwang, Some inequalities for n -time differentiable mappings and applications, Kyug-pook Math. J. 43(2003), 335-343.

- [13] İ. İşcan, Generalization of different type integral inequalities for s -convex functions via fractional integrals, *Appl. Anal.* (2014). in press.
- [14] W.-D. Jiang, D.-W. Niu, Y. Hua, and F. Qi, Generalizations of Hermite-Hadamard inequality to n -time differentiable functions which are s -convex in the second sense, *Analysis (Munich)* 32 (2012), 1001–1012.
- [15] U.S. Kırmacı, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comp.*, 147 (2004), 137-146.
- [16] U.S. Kırmacı and M.E. Özdemir, On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comp.*, 153 (2004), 361-368.
- [17] U.S. Kırmacı, Improvement and further generalization of inequalities for differentiable mappings and applications, *Comp and Math. with Appl.*, 55 (2008), 485-493.
- [18] H. Kavurmaci and M. Tunç, On some inequalities for s -logarithmically convex in the second sense via fractional integrals, arXiv:1212.1604 [math.FA].
- [19] M. A. Latif and S. S. Dragomir, New inequalities of Hermite-Hadamard type for functions whose derivatives in absolute value are convex with applications, *Acta Univ. M. Belii, ser. Math.* (2013), 24–39.
- [20] M. A. Latif and S. S. Dragomir, On Hermite-Hadamard type integral inequalities for n -times differentiable (α, m) -logarithmically convex functions, *RGMIA Research Report Collection*, 17(2014), Article 14, 16 pp.
- [21] M. A. Latif and S. S. Dragomir, New inequalities of Hermite-Hadamard type for n -times differentiable convex and concave functions with applications. (Submitted)
- [22] M. A. Latif and S. S. Dragomir, On Hermite-Hadamard type integral inequalities for n -times differentiable log-preinvex functions. (Accepted)
- [23] M. A. Latif and S. S. Dragomir, Generalization of Hermite-Hadamard type inequalities for n -times differentiable functions which are s -preinvex in the second sense with applications. (Accepted)
- [24] M. Avci, H. Kavurmaci, M. E. Özdemir, New inequalities of Hermite-Hadamard type via s -convex functions in the second sense with applications, *Appl. Math. Comp.*, 217 (2011) 5171–5176.
- [25] D. S. Mitrinović and I. B. Lacković, Hermite and convexity, *Aequationes Math.* 28 (1985), 229–232.
- [26] C.E.M. Pearce and J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formulae, *Appl. Math. Lett.*, 13(2) (2000), 51–55.
- [27] M. A. Noor, F. Qi and M. U. Awan, Some Hermite-Hadamard type inequalities for log- h -convex functions, *Analysis* 12/2013, 33(4):367–375. DOI:10.1524/anly.2013.1223
- [28] M. Z. Sarikaya, A. Saglam and H. Yildirim, On some Hadamard-type inequalities for h -convex functions, *J Math Inequal*, Vol. 2, No. 3 (2008), 335-341.
- [29] M. Z. Sarikaya and N. Aktan, On the generalization some integral inequalities and their applications *Mathematical and Computer Modelling*, Volume 54, Issues 9-10, November 2011, Pages 2175-2182.
- [30] M. Z. Sarikaya, E. Set and M. E. Ozdemir, On some new inequalities of Hadamard type involving h -convex functions, *Acta Mathematica Universitatis Comenianae*, Vol. LXXIX, 2(2010), pp. 265-272.
- [31] A. Saglam, M. Z. Sarikaya and H. Yildirim, Some new inequalities of Hermite-Hadamard's type, *Kyungpook Mathematical Journal*, 50(2010), 399-410.
- [32] M. Tunç and A. O. Akdemir, Ostrowski type inequalities for s -logarithmically convex functions in the second sense with applications, arXiv:1301.3041 [math.CA].
- [33] J. Wang, J. Deng, M. Fečkan, Exploring s -e-condition and applications to some Ostrowski type inequalities via Hadamard fractional integrals. *Math. Slovaca* (2013, in press).
- [34] S.-H. Wang, B.-Y. Xi, F. Qi, Some new inequalities of Hermite-Hadamard type for n -time differentiable functions which are m -convex, *Analysis (Munich)* 32 (2012) 247–262.
- [35] B.-Y. Xi, R.-F. Bai and F. Qi, Hermite-Hadamard type inequalities for the m - and (α, m) -geometrically convex functions, *Aequationes Math.* 39 (2012), in press.
- [36] B.-Y. Xi and F. Qi, Some integral inequalities of Hermite-Hadamard type for s -logarithmically convex functions. (Accepted)
- [37] T.-Y. Zhang, A.-P. Ji, F. Qi, On integral inequalities of Hermite-Hadamard type for s -geometrically convex functions, *Abstr. Appl. Anal.*, 2012 (2012).

- [38] T.-Y. Zhang, M. Tunç, A.-P. Ji and B.-Y. Xi, Erratum to “On integral inequalities of Hermite-Hadamard type for s -geometrically convex functions”, *Abstr. Appl. Anal.*, Volume 2014, Article ID 294739, 5 pages <http://dx.doi.org/10.1155/2014/294739>.

SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: `m.amer.latif@hotmail.com`

¹ SCHOOL OF ENGINEERING AND SCIENCE, VICTORIA UNIVERSITY, P. O. BOX 14428, MELBOURNE CITY, MC8001, AUSTRALIA

²SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: `sever.dragomir@vu.edu.au`

SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: `Ebrahim.Momoniata@wits.ac.za`