

## ON THE WEIGHTED INTEGRAL INEQUALITIES FOR CONVEX FUNCTION

MEHMET ZEKI SARIKAYA AND SAMET ERDEN

ABSTRACT. In this paper, we establish several weighted inequalities for some differentiable mappings that are connected with the celebrated Hermite-Hadamard-Fejér type and Ostrowski type integral inequalities. The results presented here would provide extensions of those given in earlier works.

### 1. INTRODUCTION

The following result is known in the literature as Ostrowski's inequality [10]:

**Theorem 1.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  whose derivative  $f' : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ , i.e.,  $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$ . Then, the inequality holds:*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for all  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is the best possible.

Inequality (1.1) has wide applications in numerical analysis and in the theory of some special means; estimating error bounds for some special means, some midpoint, trapezoid and Simpson rules and quadrature rules, etc. Hence inequality (1.1) has attracted considerable attention and interest from mathematicians and researchers. Due to this, over the years, the interested reader is also referred to ([1]-[7],[12]-[17]) for integral inequalities in several independent variables. In addition, the current approach of obtaining the bounds, for a particular quadrature rule, have depended on the use of Peano kernel. The general approach in the past has involved the assumption of bounded derivatives of degree greater than one.

If  $f : [a, b] \rightarrow \mathbb{R}$  is differentiable on  $[a, b]$  with the first derivative  $f'$  integrable on  $[a, b]$ , then Montgomery identity holds:

$$(1.2) \quad f(x) = \frac{1}{b-a} \int_a^b f(t) dt + \int_a^b P(x, t) f'(t) dt,$$

---

2000 *Mathematics Subject Classification.* 26D07, 26D15.

*Key words and phrases.* Ostrowski's inequality, Montgomery's identities, convex function, Hölder inequality.

where  $P(x, t)$  is the Peano kernel defined by

$$P(x, t) := \begin{cases} \frac{t-a}{b-a}, & a \leq t < x \\ \frac{t-b}{b-a}, & x \leq t \leq b. \end{cases}$$

**Definition 1.** The function  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ , is said to be convex if the following inequality holds

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$

for all  $x, y \in [a, b]$  and  $\lambda \in [0, 1]$ . We say that  $f$  is concave if  $(-f)$  is convex.

The following inequality is well known in the literature as the Hermite-Hadamard integral inequality (see, [11]):

$$(1.3) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}$$

where  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  is a convex function on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ .

The most well-known inequalities related to the integral mean of a convex function are the Hermite Hadamard inequalities or its weighted versions, the so-called Hermite-Hadamard-Fejér inequalities (see, [18]-[22]). In [8], Fejer gave a weighted generalization of the inequalities (1.3) as the following:

**Theorem 2.**  $f : [a, b] \rightarrow \mathbb{R}$ , be a convex function, then the inequality

$$(1.4) \quad f\left(\frac{a+b}{2}\right) \int_a^b w(x)dx \leq \frac{1}{b-a} \int_a^b f(x)w(x)dx \leq \frac{f(a) + f(b)}{2} \int_a^b w(x)dx$$

holds, where  $w : [a, b] \rightarrow \mathbb{R}$  is nonnegative, integrable, and symmetric about  $x = \frac{a+b}{2}$ .

In [18], some inequalities of Hermite-Hadamard-Fejer type for differentiable convex mappings were proved using the following lemma.

**Lemma 1.** Let  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ , and  $w : [a, b] \rightarrow [0, \infty)$  be a differentiable mapping. If  $f' \in L[a, b]$ , then the following equality holds:

$$(1.5) \quad \frac{1}{b-a} \int_a^b f(x)w(x)dx - \frac{1}{b-a} f\left(\frac{a+b}{2}\right) \int_a^b w(x)dx = (b-a) \int_0^1 k(t)f'(ta+(1-t)b)dt$$

for each  $t \in [0, 1]$ , where

$$k(t) = \begin{cases} \int_0^t w(as + (1-s)b)ds, & t \in [0, \frac{1}{2}) \\ -\int_t^1 w(as + (1-s)b)ds, & t \in [\frac{1}{2}, 1]. \end{cases}$$

The main result in [18] is as follows:

**Theorem 3.** Let  $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ , and  $w : [a, b] \rightarrow [0, \infty)$  be a differentiable mapping and symmetric to  $\frac{a+b}{2}$ . If

$|f'|$  is convex on  $[a, b]$ , then the following inequality holds:

$$(1.6) \quad \left| \frac{1}{b-a} \int_a^b f(x)w(x)dx - \frac{1}{b-a} f\left(\frac{a+b}{2}\right) \int_a^b w(x)dx \right| \\ \leq \left( \frac{1}{(b-a)^2} \int_{\frac{a+b}{2}}^b w(x) [(x-a)^2 - (b-x)^2] dx \right) \left( \frac{|f'(a)| + |f'(b)|}{2} \right).$$

In this article, using functions whose derivatives absolute values are convex, we obtained new inequalities of Fejer-Hermite-Hadamard type and Ostrowski type. The results presented here would provide extensions of those given in earlier works.

## 2. MAIN RESULTS

We will establish some new results connected with the left-hand side of (1.4) and Ostrowski type inequalities used the following Lemma. Now, we give the following new Lemma for our results:

**Lemma 2.** Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$  and let  $w : [a, b] \rightarrow \mathbb{R}$ . If  $f', w \in L[a, b]$ , then, for all  $x \in [a, b]$ , the following equality holds:

$$(2.1) \quad \int_a^x \left( \int_a^t w(s)ds \right)^\alpha f'(t)dt - \int_x^b \left( \int_t^b w(s)ds \right)^\alpha f'(t)dt \\ = \left[ \left( \int_a^x w(s)ds \right)^\alpha + \left( \int_x^b w(s)ds \right)^\alpha \right] f(x) \\ - \alpha \int_a^x \left( \int_a^t w(s)ds \right)^{\alpha-1} w(t)f(t)dt - \alpha \int_x^b \left( \int_t^b w(s)ds \right)^{\alpha-1} w(t)f(t)dt$$

*Proof.* By integration by parts, we have the following equalities:

$$(2.2) \quad \int_a^x \left( \int_a^t w(s)ds \right)^\alpha f'(t)dt \\ = \left( \int_a^t w(s)ds \right)^\alpha f(t) \Big|_a^x - \alpha \int_a^x \left( \int_a^t w(s)ds \right)^{\alpha-1} w(t)f(t)dt \\ = \left( \int_a^x w(s)ds \right)^\alpha f(x) - \alpha \int_a^x \left( \int_a^t w(s)ds \right)^{\alpha-1} w(t)f(t)dt$$

and

$$\begin{aligned}
(2.3) \quad & \int_x^b \left( \int_t^b w(s) ds \right)^\alpha f'(t) dt \\
&= \left( \int_t^b w(s) ds \right)^\alpha f(t) \Big|_x^b - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \\
&= - \left( \int_x^b w(s) ds \right)^\alpha f(x) + \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt
\end{aligned}$$

Subtracting to (2.2) from (2.3), we obtain (2.1)

$$\begin{aligned}
& \int_a^x \left( \int_a^t w(s) ds \right)^\alpha f'(t) dt - \int_x^b \left( \int_t^b w(s) ds \right)^\alpha f'(t) dt \\
&= \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \\
&\quad - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt.
\end{aligned}$$

This completes the proof.  $\square$

**Corollary 1.** *Under the same assumptions of Lemma 2 with  $\alpha = 1$ , then the following identity holds:*

$$(2.4) \quad \left( \int_a^b w(s) ds \right) f(x) - \int_a^b w(t) f(t) dt = \int_a^x \left( \int_a^t w(s) ds \right) f'(t) dt - \int_x^b \left( \int_t^b w(s) ds \right) f'(t) dt$$

**Remark 1.** *If we take  $w(s) = 1$  in (2.4), the identity (2.4) reduces to the identity (1.2).*

**Definition 2.** *Let  $f \in L_1[a, b]$ . The Riemann-Liouville integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by*

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function and  $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$ .

**Corollary 2.** *Under the same assumptions of Lemma 2 with  $w(s) = 1$ , then the following equality holds:*

$$(2.5) \quad \begin{aligned} & [(x-a)^\alpha + (b-x)^\alpha] f(x) - \Gamma(\alpha+1) J_{x^-}^\alpha f(a) - \Gamma(\alpha+1) J_{x^+}^\alpha f(b) \\ &= \int_a^x (t-a)^\alpha f'(t) dt - \int_x^b (b-t)^\alpha f'(t) dt. \end{aligned}$$

**Corollary 3.** *Under the same assumptions of Corollary 2 with  $x = \frac{a+b}{2}$ , the identity (2.5) becomes to the following identity*

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{2^{1-\alpha}(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \\ &= \frac{1}{2^{1-\alpha}(b-a)^\alpha} \left\{ \int_a^{\frac{a+b}{2}} (t-a)^\alpha f'(t) dt - \int_{\frac{a+b}{2}}^b (b-t)^\alpha f'(t) dt \right\}. \end{aligned}$$

Now, by using the above lemma, we prove our main theorems:

**Theorem 4.** *Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ . If  $|f'|$  is convex on  $[a, b]$ , then the following inequality holds:*

$$\begin{aligned} & \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\ & \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\ & \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{b-a} \left\{ \left( \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} - \frac{(x-a)^{\alpha+2}}{\alpha+2} \right) |f'(a)| + \frac{(x-a)^{\alpha+2}}{\alpha+2} |f'(b)| \right\} \\ & \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{b-a} \left\{ \frac{(b-x)^{\alpha+2}}{\alpha+2} |f'(a)| + \left( \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} - \frac{(b-x)^{\alpha+2}}{\alpha+2} \right) |f'(b)| \right\} \\ & \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{b-a} \left\{ \left( \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} + \frac{(b-x)^{\alpha+2} - (x-a)^{\alpha+2}}{\alpha+2} \right) |f'(a)| \right. \\ & \quad \left. + \left( \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} + \frac{(x-a)^{\alpha+2} - (b-x)^{\alpha+2}}{\alpha+2} \right) |f'(b)| \right\} \end{aligned}$$

where  $\alpha > 0$  and  $\|w\|_{[a,b],\infty} = \sup_{t \in [a,b]} |w(t)|$ .

*Proof.* We take absolute value of (2.1), we find that

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \int_a^x \left( \int_a^t w(s) ds \right)^\alpha |f'(t)| dt + \int_x^b \left( \int_t^b w(s) ds \right)^\alpha |f'(t)| dt \\
& \leq \|w\|_{[a,x],\infty}^\alpha \int_a^x (t-a)^\alpha |f'(t)| dt + \|w\|_{[x,b],\infty}^\alpha \int_x^b (b-t)^\alpha |f'(t)| dt \\
& = \|w\|_{[a,x],\infty}^\alpha \int_a^x (t-a)^\alpha \left| f' \left( \frac{b-t}{b-a} a + \frac{t-a}{b-a} b \right) \right| dt \\
& \quad + \|w\|_{[x,b],\infty}^\alpha \int_x^b (b-t)^\alpha \left| f' \left( \frac{b-t}{b-a} a + \frac{t-a}{b-a} b \right) \right| dt
\end{aligned}$$

Since  $|f'|$  is convex on  $[a,b]$ , it follows that

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \|w\|_{[a,x],\infty}^\alpha \int_a^x (t-a)^\alpha \left[ \frac{b-t}{b-a} |f'(a)| + \frac{t-a}{b-a} |f'(b)| \right] dt \\
& \quad + \|w\|_{[x,b],\infty}^\alpha \int_x^b (b-t)^\alpha \left[ \frac{b-t}{b-a} |f'(a)| + \frac{t-a}{b-a} |f'(b)| \right] dt \\
& = \frac{\|w\|_{[a,x],\infty}^\alpha}{b-a} \left\{ \left( \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} - \frac{(x-a)^{\alpha+2}}{\alpha+2} \right) |f'(a)| + \frac{(x-a)^{\alpha+2}}{\alpha+2} |f'(b)| \right\} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{b-a} \left\{ \frac{(b-x)^{\alpha+2}}{\alpha+2} |f'(a)| + \left( \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} - \frac{(b-x)^{\alpha+2}}{\alpha+2} \right) |f'(b)| \right\} \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{b-a} \left\{ \left( \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} + \frac{(b-x)^{\alpha+2} - (x-a)^{\alpha+2}}{\alpha+2} \right) |f'(a)| \right. \\
& \quad \left. + \left( \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} + \frac{(x-a)^{\alpha+2} - (b-x)^{\alpha+2}}{\alpha+2} \right) |f'(b)| \right\}.
\end{aligned}$$

Hence, the proof of theorem is completed.  $\square$

**Corollary 4.** *Under the same assumptions of Theorem 4 with  $w(s) = 1$ , then the following inequality holds:*

$$(2.6) \quad \begin{aligned} & \left| [(x-a)^\alpha + (b-x)^\alpha] f(x) - \Gamma(\alpha+1) [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\ & \leq \frac{1}{b-a} \left\{ \left( \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} + \frac{(b-x)^{\alpha+2} - (x-a)^{\alpha+2}}{\alpha+2} \right) |f'(a)| \right. \\ & \quad \left. + \left( \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} + \frac{(x-a)^{\alpha+2} - (b-x)^{\alpha+2}}{\alpha+2} \right) |f'(b)| \right\}. \end{aligned}$$

**Remark 2.** *If we take  $x = \frac{a+b}{2}$  in (2.6), we get*

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)+}^\alpha f(b) \right] \right| \\ & \leq \frac{(b-a)}{4(\alpha+1)} \left( |f'(a)| + |f'(b)| \right) \end{aligned}$$

which is proved by Sarikaya and Yildirim in [19].

**Corollary 5.** *Under the same assumptions of Theorem 4 with  $\alpha = 1$ , then the following inequality holds:*

$$\begin{aligned} & \left| \left( \int_a^b w(s) ds \right) f(x) - \int_a^b w(t) f(t) dt \right| \\ & \leq \frac{\|w\|_{[a,b],\infty}}{b-a} \left\{ \left( \frac{(b-a)(x-a)^2}{2} + \frac{(b-x)^3 - (x-a)^3}{3} \right) |f'(a)| \right. \\ & \quad \left. + \left( \frac{(b-a)(b-x)^2}{\alpha+1} + \frac{(x-a)^3 - (b-x)^3}{\alpha+2} \right) |f'(b)| \right\}. \end{aligned}$$

**Corollary 6.** *Under the same assumptions of Corollary 5 with  $x = \frac{a+b}{2}$ , we get*

$$(2.7) \quad \left| \left( \int_a^b w(s) ds \right) f\left(\frac{a+b}{2}\right) - \int_a^b w(t) f(t) dt \right| \leq \frac{(b-a)^2 \|w\|_{[a,b],\infty}^\alpha}{4} \left( \frac{|f'(a)| + |f'(b)|}{2} \right).$$

**Remark 3.** *If we take  $w(s) = 1$  in (2.7), we have*

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)}{4} \left( \frac{|f'(a)| + |f'(b)|}{2} \right)$$

which is proved by Kırmacı in [9].

**Corollary 7.** Under the same assumptions of Theorem 4 with  $|f'(a)| = |f'(b)|$  in (2.4), then the following inequality holds:

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \frac{|f'(a)| \|w\|_{[a,x],\infty}^\alpha}{b-a} \left\{ \frac{(b-a)(x-a)^{\alpha+1}}{\alpha+1} \right\} + \frac{|f'(a)| \|w\|_{[x,b],\infty}^\alpha}{b-a} \left\{ \frac{(b-a)(b-x)^{\alpha+1}}{\alpha+1} \right\} \\
& \leq \frac{|f'(a)| \|w\|_{[a,b],\infty}^\alpha}{\alpha+1} [(x-a)^{\alpha+1} + (b-x)^{\alpha+1}]
\end{aligned}$$

**Theorem 5.** Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ . If  $|f'|^q$  is convex on  $[a, b]$ ,  $q > 1$ , then the following inequality holds:

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left( \frac{(x-a)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left( \frac{(b-x)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left\{ \left( \frac{(x-a)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \frac{(b-x)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right\}
\end{aligned}$$

where  $\alpha > 0$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $\|w\|_{[a,b],\infty} = \sup_{t \in [a,b]} |w(t)|$ .



*Proof.* We take absolute value of (2.1). Using Holder's inequality, we find that

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \int_a^x \left| \int_a^t w(s) ds \right|^\alpha |f'(t)| dt + \int_x^b \left| \int_t^b w(s) ds \right|^\alpha |f'(t)| dt \\
& \leq \int_a^x \left( \left| \int_a^t w(s) ds \right|^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_a^x |f'(t)|^q dt \right)^{\frac{1}{q}} + \int_x^b \left( \left| \int_t^b w(s) ds \right|^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_x^b |f'(t)|^q dt \right)^{\frac{1}{q}} \\
& \leq \|w\|_{[a,x],\infty}^\alpha \left( \int_a^x |t-a|^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_a^x |f'(t)|^q dt \right)^{\frac{1}{q}} \\
& \quad + \|w\|_{[x,b],\infty}^\alpha \left( \int_x^b |b-t|^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_x^b |f'(t)|^q dt \right)^{\frac{1}{q}}
\end{aligned}$$

Since  $|f'(t)|^q$  is convex on  $[a, b]$

$$(2.9) \quad \left| f' \left( \frac{b-t}{b-a} a + \frac{t-a}{b-a} b \right) \right|^q \leq \frac{b-t}{b-a} |f'(a)|^q + \frac{t-a}{b-a} |f'(b)|^q$$

From (2.9), it follows that

$$\begin{aligned}
& \left| \left[ \left( \int_a^x w(s) ds \right)^\alpha + \left( \int_x^b w(s) ds \right)^\alpha \right] f(x) \right. \\
& \quad \left. - \alpha \int_a^x \left( \int_a^t w(s) ds \right)^{\alpha-1} w(t) f(t) dt - \alpha \int_x^b \left( \int_t^b w(s) ds \right)^{\alpha-1} w(t) f(t) dt \right| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left( \frac{(x-a)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left( \frac{(b-x)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{(b-a)^{\frac{1}{q}}} \left\{ \left( \frac{(x-a)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \frac{(b-x)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right\}
\end{aligned}$$

which this completes the proof.  $\square$

**Corollary 8.** *Under the same assumptions of Theorem 4 with  $w(s) = 1$ , then the following inequality holds:*

$$\begin{aligned}
& |[(x-a)^\alpha + (b-x)^\alpha] f(x) - \Gamma(\alpha+1) [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)]| \\
& \leq \frac{1}{(b-a)^{\frac{1}{q}}} \left\{ \left( \frac{(x-a)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \frac{(b-x)^{\alpha p+1}}{\alpha p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right\}.
\end{aligned}$$

**Remark 4.** *If we take  $x = \frac{a+b}{2}$  in (2.10), we have*

$$\begin{aligned}
& \left| f\left(\frac{a+b}{2}\right) - \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)+}^\alpha f(b) \right] \right| \\
& \leq \frac{2^{\alpha-1}(b-a)}{(\alpha p+1)^{\frac{1}{p}}} \left[ \left( \frac{3|f'(a)|^q + |f'(b)|^q}{8} \right)^{\frac{1}{q}} + \left( \frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right)^{\frac{1}{q}} \right]
\end{aligned}$$

which is proved by Sarikaya and Yildirim in [19].

**Corollary 9.** *Let the conditions of Theorem 5 hold. If we take  $\alpha = 1$  in (2.8), then the following inequality holds:*

$$\begin{aligned} & \left| \left( \int_a^b w(s) ds \right) f(x) - \int_a^b w(t) f(t) dt \right| \\ & \leq \frac{\|w\|_{[a,b],\infty}}{(b-a)^{\frac{1}{q}}} \left\{ \left( \frac{(x-a)^{p+1}}{p+1} \right)^{\frac{1}{p}} \left( \frac{(b-a)^2 - (b-x)^2}{2} |f'(a)|^q + \frac{(x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left( \frac{(b-x)^{p+1}}{p+1} \right)^{\frac{1}{p}} \left( \frac{(b-x)^2}{2} |f'(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f'(b)|^q \right)^{\frac{1}{q}} \right\} \end{aligned}$$

**Corollary 10.** *Under the same assumptions of Corollary 9 with  $x = \frac{a+b}{2}$ , we get*

$$\begin{aligned} (2.11) \quad & \left| \left( \int_a^b w(s) ds \right) f\left(\frac{a+b}{2}\right) - \int_a^b w(t) f(t) dt \right| \\ & \leq \frac{(b-a) \|w\|_{[a,b],\infty}}{2^{2+\frac{1}{q}}(p+1)^{\frac{1}{p}}} \left\{ \left( \frac{3|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}} + \left( \frac{|f'(a)|^q + 3|f'(b)|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 5.** *If we take  $w(s) = 1$  in (2.11), we have*

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(b-a)}{2^{2+\frac{1}{q}}(p+1)^{\frac{1}{p}}} \left\{ \left( \frac{3|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}} + \left( \frac{|f'(a)|^q + 3|f'(b)|^q}{2} \right)^{\frac{1}{q}} \right\} \end{aligned}$$

which is proved by Kirmaci in [9].

#### REFERENCES

- [1] F. Ahmad, N. S. Barnett and S. S. Dragomir, *New Weighted Ostrowski and Cebysev Type Inequalities*, *Nonlinear Analysis: Theory, Methods & Appl.*, 71 (12), (2009), 1408-1412.
- [2] F. Ahmad, A. Rafiq, N. A. Mir, *Weighted Ostrowski type inequality for twice differentiable mappings*, *Global Journal of Research in Pure and Applied Math.*, 2 (2) (2006), 147-154.
- [3] N. S. Barnett and S. S. Dragomir, *An Ostrowski type inequality for double integrals and applications for cubature formulae*, *Soochow J. Math.*, 27(1), (2001), 109-114.
- [4] N. S. Barnett, S. S. Dragomir and C.E.M. Pearce, *A Quasi-trapezoid inequality for double integrals*, *ANZIAM J.*, 44(2003), 355-364.
- [5] S. S. Dragomir, P. Cerone, N. S. Barnett and J. Roumeliotis, *An inequality of the Ostrowski type for double integrals and applications for cubature formulae*, *Tamsui Oxf. J. Math.*, 16(1), (2000), 1-16.
- [6] S. Hussain, M.A.Latif and M. Alomari, *Generalized duple-integral Ostrowski type inequalities on time scales*, *Appl. Math. Letters*, 24(2011), 1461-1467.
- [7] M. E. Kiris and M. Z. Sarikaya, *On the new generalization of Ostrowski type inequality for double integrals*, *International Journal of Modern Mathematical Sciences*, 2014, 9(3): 221-229.
- [8] L. Fejer, *Über die Fourierreihen*, II. *Math. Naturwiss. Anz Ungar. Akad. Wiss.*, 24 (1906), 369-390. (Hungarian).

- [9] U.S. Kırmacı, *Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, Appl. Math. Comp., 147 (2004), 137-146.
- [10] A. M. Ostrowski, *Über die absolutabweichung einer differentiebaren funktion von ihrem integralmittelwert*, Comment. Math. Helv. 10(1938), 226-227.
- [11] J. Pečarić, F. Proschan and Y.L. Tong, *Convex functions, partial ordering and statistical applications*, Academic Press, New York, 1991.
- [12] A. Qayyum, *A weighted Ostrowski-Grüss type inequality and applications*, Proceeding of the World Cong. on Engineering, Vol:2, 2009, 1-9.
- [13] A. Rafiq and F. Ahmad, *Another weighted Ostrowski-Grüss type inequality for twice differentiable mappings*, Kragujevac Journal of Mathematics, 31 (2008), 43-51.
- [14] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [15] M. Z. Sarikaya *On the Ostrowski type integral inequality for double integrals*, Demonstratio Mathematica, Vol. XLV, No 3, pp:533-540, 2012.
- [16] M. Z. Sarikaya and H. Ogunmez, *On the weighted Ostrowski type integral inequality for double integrals*, The Arabian Journal for Science and Engineering (AJSE)-Mathematics, (2011) 36:1153-1160.
- [17] M. Z. Sarikaya, *On the generalized weighted integral inequality for double integrals*, Annals of the Alexandru Ioan Cuza University-Mathematics, accepted.
- [18] M. Z. Sarikaya, *On new Hermite Hadamard Fejer Type integral inequalities*, Studia Universitatis Babes-Bolyai Mathematica., 57(2012), No. 3, 377-386.
- [19] M. Z. Sarikaya and H. Yildirim, *On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals*, Submitted
- [20] K-L. Tseng, G-S. Yang and K-C. Hsu, *Some inequalities for differentiable mappings and applications to Fejer inequality and weighted trapezoidal formula*, Taiwanese J. Math. 15(4), pp:1737-1747, 2011.
- [21] C.-L. Wang, X.-H. Wang, *On an extension of Hadamard inequality for convex functions*, Chin. Ann. Math. 3 (1982) 567-570.
- [22] S.-H. Wu, *On the weighted generalization of the Hermite-Hadamard inequality and its applications*, The Rocky Mountain J. of Math., vol. 39, no. 5, pp. 1741-1749, 2009.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, KONURALP CAMPUS, DÜZCE-TURKEY  
E-mail address: sarikayamz@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BARTIN UNIVERSITY, BARTIN-TURKEY  
E-mail address: erdem1627@gmail.com