

**GENERALIZED HERMITE -HADAMARD TYPE INTEGRAL  
INEQUALITIES FOR FRACTIONAL INTEGRALS**

MEHMET ZEKI SARIKAYA AND HÜSEYİN BUDAK

ABSTRACT. In this paper, we have established Hermite-Hadamard type inequalities for fractional integrals depending on a parameter.

1. INTRODUCTION

**Definition 1.** *The function  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ , is said to be convex if the following inequality holds*

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$

for all  $x, y \in [a, b]$  and  $\lambda \in [0, 1]$ . We say that  $f$  is concave if  $(-f)$  is convex.

The inequalities discovered by C. Hermite and J. Hadamard for convex functions are very important in the literature (see, e.g., [15, p.137], [9]). These inequalities state that if  $f : I \rightarrow \mathbb{R}$  is a convex function on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ , then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}.$$

Both inequalities hold in the reversed direction if  $f$  is concave. We note that Hadamard's inequality may be regarded as a refinement of the concept of convexity and it follows easily from Jensen's inequality. Hadamard's inequality for convex functions has received renewed attention in recent years and a remarkable variety of refinements and generalizations have been found (see, for example, [1, 2, 9, 10, 15]) and the references cited therein.

In [10], Dragomir and Agarwal proved the following results connected with the right part of (1.1).

**Lemma 1.** *Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ . If  $f' \in L[a, b]$ , then the following equality holds:*

$$(1.2) \quad \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)dx = \frac{b-a}{2} \int_0^1 (1-2t)f'(ta + (1-t)b)dt.$$

**Theorem 1.** *Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable mapping on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ . If  $|f'|$  is convex on  $[a, b]$ , then the following inequality holds:*

$$(1.3) \quad \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)dx \right| \leq \frac{(b-a)}{8} (|f'(a)| + |f'(b)|).$$

---

2000 *Mathematics Subject Classification.* 26D07, 26D10, 26D15, 26A33.

*Key words and phrases.* Hermite-Hadamard's inequalities, Riemann-Liouville fractional integral, integral inequalities.

Meanwhile, Sarikaya et al.[18] presented the following important integral identity including the first-order derivative of  $f$  to establish many interesting Hermite-Hadamard type inequalities for convexity functions via Riemann-Liouville fractional integrals of the order  $\alpha > 0$ .

**Lemma 2.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $a < b$ . If  $f' \in L[a, b]$ , then the following equality for fractional integrals holds:*

$$(1.4) \quad \frac{f(a) + f(b)}{2} - \frac{\Gamma(\alpha + 1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] = \frac{b-a}{2} \int_0^1 [(1-t)^\alpha - t^\alpha] f'(ta + (1-t)b) dt.$$

It is remarkable that Sarikaya et al.[18] first give the following interesting integral inequalities of Hermite-Hadamard type involving Riemann-Liouville fractional integrals.

**Theorem 2.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a positive function with  $0 \leq a < b$  and  $f \in L_1[a, b]$ . If  $f$  is a convex function on  $[a, b]$ , then the following inequalities for fractional integrals hold:*

$$(1.5) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha + 1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2}$$

with  $\alpha > 0$ .

In the following we will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper. More details, one can consult [11, 12, 14, 16].

**Definition 2.** *Let  $f \in L_1[a, b]$ . The Riemann-Liouville integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by*

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function and  $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$ .

For some recent results connected with fractional integral inequalities see ([3, 4, 5, 6, 7, 8],[13],[17],[19],[20],[21])

The aim of this paper is to establish generalized Hermite-Hadamard type integral inequalities for Riemann-Liouville fractional integral and some other integral inequalities using the generalized identity is obtained for fractional integrals. The results presented in this paper provide extensions of those given in earlier works.

## 2. MAIN RESULTS

We give a important fractional integral identity for differentiable convex functions:

**Lemma 3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $0 \leq a < b$ . If  $f' \in L[a, b]$ , then the following equality for fractional integrals hold:

$$\begin{aligned}
& -\frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} + \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \\
& \times \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \\
(2.1) \quad & = \int_0^1 [(1-t)^\alpha - t^\alpha] f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt
\end{aligned}$$

where  $\lambda \in [0, 1] \setminus \{\frac{1}{2}\}$  and  $\alpha > 0$ .

*Proof.* It suffices to note that

$$\begin{aligned}
I & = \int_0^1 [(1-t)^\alpha - t^\alpha] f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
& = \int_0^1 (1-t)^\alpha f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
& \quad - \int_0^1 t^\alpha f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
& = I_1 - I_2
\end{aligned}$$

Integrating by parts

$$\begin{aligned}
I_1 & = \int_0^1 (1-t)^\alpha f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
& = \frac{(1-t)^\alpha f [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]}{(1-2\lambda)(b-a)} \Big|_0^1 \\
& \quad + \frac{\alpha}{(1-2\lambda)(b-a)} \int_0^1 (1-t)^{\alpha-1} f [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt
\end{aligned}$$

$$\begin{aligned}
&= -\frac{f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} \\
&\quad + \frac{\alpha}{(1-2\lambda)(b-a)} \int_0^1 (1-t)^{\alpha-1} f[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
&= -\frac{f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} + \frac{\alpha}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \int_{\lambda b + (1-\lambda)a}^{\lambda a + (1-\lambda)b} [(\lambda a + (1-\lambda)b) - x] f(x) dx \\
&= -\frac{f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} + \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b)
\end{aligned}$$

and similarly we get

$$\begin{aligned}
I_2 &= \int_0^1 t^\alpha f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)] dt \\
&= \frac{f(\lambda a + (1-\lambda)b)}{(1-2\lambda)(b-a)} - \frac{\alpha}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \int_{\lambda b + (1-\lambda)a}^{\lambda a + (1-\lambda)b} [x - (\lambda b + (1-\lambda)a)] f(x) dx \\
&= \frac{f(\lambda a + (1-\lambda)b)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a)
\end{aligned}$$

From  $I_1$  and  $I_2$ , it follows that

$$\begin{aligned}
I &= I_1 - I_2 \\
&= -\frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} \\
&\quad + \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right].
\end{aligned}$$

This completes the proof.  $\square$

**Remark 1.** If we take  $\lambda = 0$  in Lemma 3, then the identity (2.1) reduces the identity (1.4) which is proved in [18]. Similarly, if we take  $\lambda = 1$  in Lemma 3, then (2.2)

$$\frac{f(b) + f(a)}{(b-a)} + \frac{\Gamma(\alpha+1)}{(-1)^{\alpha+1}(b-a)^{\alpha+1}} [J_{b^+}^\alpha f(a) + J_{a^-}^\alpha f(b)] = \int_0^1 [(1-t)^\alpha - t^\alpha] f'(ta + (1-t)b) dt.$$

By using  $J_{b^+}^\alpha f(a) + J_{a^-}^\alpha f(b) = (-1)^\alpha [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)]$  in (2.2), it follows that (2.3)

$$\frac{f(b) + f(a)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] = \frac{(b-a)}{2} \int_0^1 [(1-t)^\alpha - t^\alpha] f'(ta + (1-t)b) dt.$$

If we choose  $\alpha = 1$  in (2.3), it follows that (2.3) reduces to (1.2).

**Theorem 3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $0 \leq a < b$ . If  $|f'|^q$ ,  $q \geq 1$  is convex on  $[a, b]$ , then the following inequality for fractional integrals holds:

$$\begin{aligned}
& \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\
& \quad \left. \times \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\
(2.4) \leq & 2^{\frac{q-1}{q}} \left[ \frac{1}{\alpha+1} \left( 1 - \frac{1}{2^\alpha} \right) \right] \left[ |f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q \right]^{\frac{1}{q}}.
\end{aligned}$$

where  $\lambda \in [0, 1] \setminus \{\frac{1}{2}\}$  and  $\alpha > 0$ .

*Proof.* Firstly, we suppose that  $q = 1$ . Using Lemma 3 and convexity of  $|f'|^q$ , we find that

$$\begin{aligned}
& \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\
& \quad \left. \times \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\
& \leq \int_0^1 |(1-t)^\alpha - t^\alpha| |f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]| dt \\
& \leq \int_0^1 |(1-t)^\alpha - t^\alpha| [t |f'(\lambda a + (1-\lambda)b)| + (1-t) |f'(\lambda b + (1-\lambda)a)|] dt \\
& = \int_0^{\frac{1}{2}} [(1-t)^\alpha - t^\alpha] [t |f'(\lambda a + (1-\lambda)b)| + (1-t) |f'(\lambda b + (1-\lambda)a)|] dt \\
& \quad + \int_{\frac{1}{2}}^1 [t^\alpha - (1-t)^\alpha] [t |f'(\lambda a + (1-\lambda)b)| + (1-t) |f'(\lambda b + (1-\lambda)a)|] dt \\
(2.5) = & K_1 + K_2.
\end{aligned}$$

Hence, conculating  $K_1$  ve  $K_2$ , we have

$$\begin{aligned}
K_1 &= \int_0^{\frac{1}{2}} [(1-t)^\alpha - t^\alpha] [t |f'(\lambda a + (1-\lambda)b)| + (1-t) |f'(\lambda b + (1-\lambda)a)|] dt \\
&= |f'(\lambda a + (1-\lambda)b)| \int_0^{\frac{1}{2}} [(1-t)^\alpha t - t^{\alpha+1}] dt \\
&\quad + |f'(\lambda b + (1-\lambda)a)| \int_0^{\frac{1}{2}} [(1-t)^{\alpha+1} - t^\alpha(1-t)] dt \\
&= |f'(\lambda a + (1-\lambda)b)| \left[ \frac{1}{(\alpha+1)(\alpha+2)} - \frac{1}{2^{\alpha+1}(\alpha+1)} \right] \\
(2.6) \quad &+ |f'(\lambda b + (1-\lambda)a)| \left[ \frac{1}{\alpha+2} - \frac{1}{2^{\alpha+1}(\alpha+1)} \right]
\end{aligned}$$

and

$$\begin{aligned}
K_2 &= \int_{\frac{1}{2}}^1 [t^\alpha - (1-t)^\alpha] [t |f'(\lambda a + (1-\lambda)b)| + (1-t) |f'(\lambda b + (1-\lambda)a)|] dt \\
&= |f'(\lambda a + (1-\lambda)b)| \left[ \frac{1}{\alpha+2} - \frac{1}{2^{\alpha+1}(\alpha+1)} \right] \\
(2.7) \quad &+ |f'(\lambda b + (1-\lambda)a)| \left[ \frac{1}{(\alpha+2)(\alpha+1)} - \frac{1}{2^{\alpha+1}(\alpha+1)} \right].
\end{aligned}$$

Using (2.6) and (2.7) in (2.5), it follows that

$$\begin{aligned}
&\left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\
&\quad \times \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \Big| \\
&\leq \frac{1}{\alpha+1} \left[ 1 - \frac{1}{2^\alpha} \right] [|f'(\lambda a + (1-\lambda)b)| + |f'(\lambda b + (1-\lambda)a)|].
\end{aligned}$$

Secondly, we suppose that  $q > 1$ . Using Lemma 3 and power mean inequality, we obtain

$$\begin{aligned}
 & \int_0^1 |(1-t)^\alpha - t^\alpha| |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]| dt \\
 & \leq \left( \int_0^1 |(1-t)^\alpha - t^\alpha| dt \right)^{1-\frac{1}{q}} \\
 (2.8) \quad & \times \left( \int_0^1 |(1-t)^\alpha - t^\alpha| |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right)^{\frac{1}{q}}.
 \end{aligned}$$

Hence, using convexity of  $|f'|^q$  and (2.8) we obtain

$$\begin{aligned}
 & \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\
 & \times \left. \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\
 & \leq \left( \int_0^1 |(1-t)^\alpha - t^\alpha| dt \right)^{1-\frac{1}{q}} \\
 & \times \left( \int_0^1 |(1-t)^\alpha - t^\alpha| |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right)^{\frac{1}{q}} \\
 & \leq \left( \int_0^{\frac{1}{2}} [(1-t)^\alpha - t^\alpha] dt + \int_{\frac{1}{2}}^1 [t^\alpha - (1-t)^\alpha] dt \right)^{1-\frac{1}{q}} \\
 & \times \left( \int_0^1 |(1-t)^\alpha - t^\alpha| [t |f'(\lambda a + (1-\lambda)b)|^q + (1-t) |f'(\lambda b + (1-\lambda)a)|^q] dt \right)^{\frac{1}{q}} \\
 & \leq \left( \frac{2}{\alpha+1} \left[ 1 - \frac{1}{2^\alpha} \right] \right)^{1-\frac{1}{q}} \left( \frac{1}{\alpha+1} \left[ 1 - \frac{1}{2^\alpha} \right] \right)^{\frac{1}{q}} \\
 & \times [ |f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q ]^{\frac{1}{q}} \\
 & \leq 2^{\frac{q-1}{q}} \left[ \frac{1}{\alpha+1} \left( 1 - \frac{1}{2^\alpha} \right) \right] [ |f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q ]^{\frac{1}{q}}.
 \end{aligned}$$

This completes the proof.  $\square$

**Corollary 1.** *Under assumption Theorem 3 with  $\lambda = 0$  and  $\lambda = 1$ , we have*

$$(2.9) \quad \left| \frac{f(a) + f(b)}{(b-a)} - \frac{\Gamma(\alpha+1)}{(b-a)^{\alpha+1}} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \right| \\ \leq 2^{\frac{q-1}{q}} \left[ \frac{1}{\alpha+1} \left( 1 - \frac{1}{2^\alpha} \right) \right] [|f'(a)|^q + |f'(b)|^q]^{\frac{1}{q}}.$$

where  $q \geq 1$ .

*Proof.* By using  $J_{b^+}^\alpha f(a) + J_{a^-}^\alpha f(b) = (-1)^\alpha [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)]$  in Theorem 3 with  $\lambda = 1$ , we obtain the inequality (2.9).  $\square$

**Remark 2.** *If we take  $\alpha = 1$  in Corollary 1, we have*

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{8} 2^{\frac{q-1}{q}} [|f'(a)|^q + |f'(b)|^q]^{\frac{1}{q}}$$

where  $q \geq 1$ . Choosing  $q = 1$  in last inequality, it follows that

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{8} [|f'(a)| + |f'(b)|]$$

which are proved Dragomir and Agarwal in [10].

**Theorem 4.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $0 \leq a < b$ . If  $|f'|^q$  convex on  $[a, b]$  for some fixed  $q > 1$ , then the following inequality for fractional integrals holds:*

$$\left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\ \left. \times [J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a)] \right| \\ \leq \left[ \frac{2}{\alpha p + 1} \left( 1 - \frac{1}{2^{\alpha p}} \right) \right]^{\frac{1}{p}} \left( \frac{|f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q}{2} \right)^{\frac{1}{q}}$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\alpha > 0$  and  $\lambda \in [0, 1] \setminus \{\frac{1}{2}\}$ .



*Proof.* Proof: Using Lemma 3, convexity of  $|f|^q$  and well-known Hölder's inequality, we obtain

$$\begin{aligned}
 & \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\
 & \quad \left. \times \left[ J_{(\lambda b+(1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a+(1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\
 & \leq \int_0^1 |(1-t)^\alpha - t^\alpha| |f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]| dt \\
 & \leq \left( \int_0^1 |(1-t)^\alpha - t^\alpha|^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f' [t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right)^{\frac{1}{q}} \\
 & \leq \left( \int_0^{\frac{1}{2}} [(1-t)^\alpha - t^\alpha]^p dt + \int_{\frac{1}{2}}^1 [t^\alpha - (1-t)^\alpha]^p dt \right)^{\frac{1}{p}} \\
 & \quad \times \left( \int_0^1 [t |f'(\lambda a + (1-\lambda)b)|^q + (1-t) |f'(\lambda b + (1-\lambda)a)|^q] dt \right)^{\frac{1}{q}} \\
 & \leq \left( \int_0^{\frac{1}{2}} [(1-t)^{\alpha p} - t^{\alpha p}] dt + \int_{\frac{1}{2}}^1 [t^{\alpha p} - (1-t)^{\alpha p}] dt \right)^{\frac{1}{p}} \\
 & \quad \times \left( \frac{|f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q}{2} \right)^{\frac{1}{q}} \\
 & \leq \left[ \frac{2}{\alpha p + 1} \left( 1 - \frac{1}{2^{\alpha p}} \right) \right]^{\frac{1}{p}} \left( \frac{|f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q}{2} \right)^{\frac{1}{q}}.
 \end{aligned}$$

Here, we use

$$(A - B)^p \leq A^p - B^p,$$

for any  $A > B \geq 0$  and  $p \geq 1$ .  $\square$

**Corollary 2.** Under assumption Theorem 4 with  $\lambda = 0$  and  $\lambda = 1$ , we have

$$\begin{aligned}
 (2.10) \quad & \left| \frac{f(a) + f(b)}{(b-a)} - \frac{\Gamma(\alpha+1)}{(b-a)^{\alpha+1}} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \right| \\
 & \leq \left[ \frac{2}{\alpha p + 1} \left( 1 - \frac{1}{2^{\alpha p}} \right) \right]^{\frac{1}{p}} \left( \frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}.
 \end{aligned}$$

*Proof.* By using  $J_{b^+}^\alpha f(a) + J_{a^-}^\alpha f(b) = (-1)^\alpha [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)]$  in Theorem 4 with  $\lambda = 1$ , we obtain the inequality (2.10).  $\square$

**Remark 3.** If we take  $\alpha = 1$  in Corollary 2, we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{2} \left[ \frac{2}{p+1} \left( 1 - \frac{1}{2^p} \right) \right]^{\frac{1}{p}} \left( \frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}$$

which are proved Dragomir and Agarwal in [10].

**Theorem 5.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $0 \leq a < b$ . If  $|f'|^q$  convex on  $[a, b]$  for some fixed  $q \geq 1$ , then the following inequality for fractional integrals holds:

$$\begin{aligned} & \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\ & \quad \times \left. \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\ & \leq \left[ \frac{1}{q\alpha+1} \left( 1 - \frac{1}{2^{q\alpha+1}} \right) \right]^{\frac{1}{q}} \left( \frac{|f'(\lambda a + (1-\lambda)b)|^q + |f'(\lambda b + (1-\lambda)a)|^q}{2} \right)^{\frac{1}{q}} \end{aligned}$$

where  $\lambda \in [0, 1] \setminus \{\frac{1}{2}\}$  and  $\alpha > 0$ .

*Proof.* Using Lemma 3, convexity of  $|f'|^q$ , and well-known Hölder's inequality, we have

$$\begin{aligned} & \left| \frac{f(\lambda a + (1-\lambda)b) + f(\lambda b + (1-\lambda)a)}{(1-2\lambda)(b-a)} - \frac{\Gamma(\alpha+1)}{(1-2\lambda)^{\alpha+1}(b-a)^{\alpha+1}} \right. \\ & \quad \times \left. \left[ J_{(\lambda b + (1-\lambda)a)^+}^\alpha f(\lambda a + (1-\lambda)b) + J_{(\lambda a + (1-\lambda)b)^-}^\alpha f(\lambda b + (1-\lambda)a) \right] \right| \\ & \leq \int_0^1 |(1-t)^\alpha - t^\alpha| |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]| dt \\ & \leq \left( \int_0^1 1^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |(1-t)^\alpha - t^\alpha|^q |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right)^{\frac{1}{q}} \\ & = \left( \int_0^{\frac{1}{2}} [(1-t)^\alpha - t^\alpha]^q |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 [t^\alpha - (1-t)^\alpha]^q |f'[t(\lambda a + (1-\lambda)b) + (1-t)(\lambda b + (1-\lambda)a)]|^q dt \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned} &\leq \left( |f'(\lambda a + (1 - \lambda)b)|^q \int_0^{\frac{1}{2}} [(1-t)^{q\alpha}t - t^{q\alpha+1}] dt \right. \\ &\quad + |f'(\lambda b + (1 - \lambda)a)|^q \int_0^{\frac{1}{2}} [(1-t)^{q\alpha+1} - t^{q\alpha}(1-t)] dt \\ &\quad + |f'(\lambda a + (1 - \lambda)b)|^q \int_{\frac{1}{2}}^1 [t^{q\alpha+1} - (1-t)^{q\alpha}t] dt \\ &\quad \left. + |f'(\lambda b + (1 - \lambda)a)|^q \int_{\frac{1}{2}}^1 [t^{q\alpha}(1-t) - (1-t)^{q\alpha+1}] dt \right)^{\frac{1}{q}} \\ &= \left( \frac{1}{\alpha+1} \left[ 1 - \frac{1}{2^{\alpha}} \right] \right)^{\frac{1}{q}} [|f'(\lambda a + (1 - \lambda)b)| + |f'(\lambda b + (1 - \lambda)a)|]^{\frac{1}{q}}. \end{aligned}$$

Here, we use  $(A - B)^p \leq A^p - B^p$ , for any  $A > B \geq 0$  and  $q \geq 1$ . □

**Corollary 3.** *Under assumption Theorem 5 with  $\lambda = 0$  and  $\lambda = 1$ , we have*

$$\begin{aligned} (2.11) \quad &\left| \frac{f(a) + f(b)}{(b-a)} - \frac{\Gamma(\alpha+1)}{(b-a)^{\alpha+1}} [J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)] \right| \\ &\leq \left[ \frac{1}{q\alpha+1} \left( 1 - \frac{1}{2^{q\alpha+1}} \right) \right]^{\frac{1}{q}} \left( \frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}. \end{aligned}$$

*Proof.* By using  $J_{b^+}^{\alpha} f(a) + J_{a^-}^{\alpha} f(b) = (-1)^{\alpha} [J_{a^+}^{\alpha} f(b) + J_{b^-}^{\alpha} f(a)]$  in Theorem 5 with  $\lambda = 1$ , we obtain the inequality (2.11). □

**Remark 4.** *If we take  $\alpha = 1$  in Corollary 3, we have*

$$\begin{aligned} &\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ &\leq \frac{b-a}{2} \left[ \frac{1}{q+1} \left( 1 - \frac{1}{2^{q+1}} \right) \right]^{\frac{1}{q}} \left( \frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}. \end{aligned}$$

REFERENCES

[1] A.G. Azpeitia, *Convex functions and the Hadamard inequality*, Rev. Colombiana Math., 28 (1994), 7-12.  
 [2] M. K. Bakula and J. Pečarić, *Note on some Hadamard-type inequalities*, Journal of Inequalities in Pure and Applied Mathematics, vol. 5, no. 3, article 74, 2004.  
 [3] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.  
 [4] Z. Dahmani, *New inequalities in fractional integrals*, International Journal of Nonlinear Science, 9(4) (2010), 493-497.

- [5] Z. Dahmani, *On Minkowski and Hermite-Hadamard integral inequalities via fractional integration*, Ann. Funct. Anal. 1(1) (2010), 51-58.
- [6] Z. Dahmani, L. Tabharit, S. Taf, *Some fractional integral inequalities*, Nonl. Sci. Lett. A, 1(2) (2010), 155-160.
- [7] Z. Dahmani, L. Tabharit, S. Taf, *New generalizations of Gruss inequality using Riemann-Liouville fractional integrals*, Bull. Math. Anal. Appl., 2(3) (2010), 93-99.
- [8] J. Deng and J. Wang, *Fractional Hermite-Hadamard inequalities for  $(\alpha, m)$ -logarithmically convex functions*. J. Inequal. Appl. 2013, Article ID 364 (2013)
- [9] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.
- [10] S. S. Dragomir and R.P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. Lett., 11(5) (1998), 91-95.
- [11] R. Gorenflo, F. Mainardi, *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien (1997), 223-276.
- [12] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204, Elsevier Sci. B.V., Amsterdam, 2006.
- [13] M.A. Latif, S.S. Dragomir and A.E. Matouk, *New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals*, J. Fract. Calc. Appl. 2, 1-15 (2012)
- [14] S. Miller and B. Ross, *An introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, USA, 1993, p.2.
- [15] J.E. Pečarić, F. Proschan and Y.L. Tong, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, Boston, 1992.
- [16] I. Podlubni, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [17] M.Z. Sarikaya and H. Ogunmez, *On new inequalities via Riemann-Liouville fractional integration*, Abstract and Applied Analysis, Volume 2012 (2012), Article ID 428983, 10 pages.
- [18] M.Z. Sarikaya, E. Set, H. Yaldiz and N., Basak, *Hermite -Hadamard's inequalities for fractional integrals and related fractional inequalities*, Mathematical and Computer Modelling, DOI:10.1016/j.mcm.2011.12.048, 57 (2013) 2403-2407.
- [19] M.Z. Sarikaya, H. Filiz and M.E. Kiris, *On some generalized integral inequalities for Riemann-Liouville fractional integrals*, Filomat, accepted.
- [20] M. Tunc, *On new inequalities for  $h$ -convex functions via Riemann-Liouville fractional integration*, Filomat 27:4 (2013), 559-565.
- [21] Y. Zhang and J. Wang, *On some new Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals*, J. Inequal. Appl. 2013, Article ID 220 (2013)

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

*E-mail address:* sarikayamz@gmail.com

*E-mail address:* hsyn.budak@gmail.com