

ON POWER POMPEIU'S TYPE INEQUALITIES FOR DOUBLE INTEGRALS

MEHMET ZEKI SARIKAYA, SAMET ERDEN, AND HATICE YALDIZ

ABSTRACT. The main of this paper is to derive some new inequalities of Ostrowski type using Pompeiu's mean value theorem for double integrals involving functions of two independent variables.

1. INTRODUCTION

In 1938, the classical integral inequality established by Ostrowski [12] as follows:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative $f' : (a, b) \rightarrow \mathbb{R}$ is bounded on (a, b) , i.e., $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then, the inequality holds:*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ is the best possible.

Inequality (1.1) has wide applications in numerical analysis and in the theory of some special means; estimating error bounds for some special means, some midpoint, trapezoid and Simpson rules and quadrature rules, etc. Hence inequality (1.1) has attracted considerable attention and interest from mathematicians and researchers. Due to this, over the years, the interested reader is also referred to ([4]-[6], [11], [13]-[17]) for integral inequalities in several independent variables.. In addition, the current approach of obtaining the bounds, for a particular quadrature rule, have depended on the use of Peano kernel. The general approach in the past has involved the assumption of bounded derivatives of degree greater than one.

In 1946, Pompeiu [8] derived a variant of Lagrange's mean value theorem, now known as *Pompeiu's mean value theorem*.

Theorem 2. *For every real valued function f differentiable on an interval $[a, b]$ not containing 0 and for all pairs $x_1 \neq x_2$ in $[a, b]$, there exist a point ξ between x_1 and x_2 such that*

$$\frac{x_1 f(x_2) - x_2 f(x_1)}{x_1 - x_2} = f(\xi) - \xi f'(\xi).$$

In [9], E.C. Popa using a mean value theorem obtained following theorem.

2000 *Mathematics Subject Classification.* 26D10, 26D15, 41A55.

Key words and phrases. Ostrowski's inequality, Pompeiu's inequalities, absolutely continuous function, double integrals.

Theorem 3. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) . Assume that $\alpha \notin [a, b]$. Then for any $x \in [a, b]$, we have the inequality

$$\begin{aligned} & \left| \left(\frac{a+b}{2} - \alpha \right) f(x) + \frac{\alpha - x}{b-a} \int_a^b f(t) dt \right| \\ & \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f - l_\alpha f'\|_\infty. \end{aligned}$$

In [7], the authors have proved the following Ostrowski type inequality:

Theorem 4. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) with $0 < a < b$. Then for $1 \leq p, q \leq \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$ we have the inequality

$$\left| \frac{a+b}{2} \cdot \frac{f(x)}{x} + \frac{1}{b-a} \int_a^b f(t) dt \right| \leq PU(x, p) \|f - l f'\|_p,$$

for $x \in [a, b]$, where

$$\begin{aligned} PU(x, p) & : = (b-a)^{\frac{1}{p}-1} \left[\left(\frac{a^{2-q} - x^{2-q}}{(1-2q)(2-q)} + \frac{x^{2-q} - a^{1+q}x^{1-2q}}{(1-2q)(1+q)} \right)^{\frac{1}{p}} \right. \\ & \quad \left. + \left(\frac{b^{2-q} - x^{2-q}}{(1-2q)(2-q)} + \frac{x^{2-q} - b^{1+q}x^{1-2q}}{(1-2q)(1+q)} \right)^{\frac{1}{q}} \right]. \end{aligned}$$

In the cases $(p, q) = (1, \infty), (\infty, 1)$ and $(2, 2)$ the quantity $PU(x, p)$ has to be taken as the limit as $p \rightarrow 1, \infty$ and 2 , respectively.

In [10], Dragomir has proved the Ostrowski type inequalities for complex valued absolutely continuous functions as follows:

Theorem 5. Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous function on the interval $[a, b]$ with $b > a > 0$. If $r \in \mathbb{R}$, $r \neq 0$, and $f'l - rf \in L_\infty[a, b]$, then for any $x \in [a, b]$ we have

$$\begin{aligned} & \left| \frac{b^{r+1} - a^{r+1}}{r+1} f(x) - x^r \int_a^b f(t) dt \right| \\ & \leq \frac{1}{|r|} \|f'l - rf\|_\infty \\ & \quad \times \begin{cases} \frac{2rx^{r+1} - x^r(a+b)(r+1) + b^{r+1} + a^{r+1}}{r+1} & , \quad \text{if } r > 0 \\ \frac{x^r(a+b)(r+1) - 2rx^{r+1} - b^{r+1} - a^{r+1}}{r+1} & , \quad \text{if } r \in (-\infty, 1) \setminus \{-1\}. \end{cases} \end{aligned}$$

Also, for $r = -1$, we have

$$\left| f(x) \ln \frac{b}{a} - \frac{1}{x} \int_a^b f(t) dt \right| \leq \frac{1}{|r|} \|f'l + f\|_\infty \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right)$$

for any $x \in [a, b]$, provided $f'l + f \in L_\infty[a, b]$.

The interested reader is also referred to ([1]-[3], [7], [9], [10], [18]-[21]) for integral inequalities by using Pompeiu's mean value theorem. The main aim of this paper is to establish some Pompeiu's type inequality for complex valued absolutely continuous functions with double integrals involving functions of two independent variables.

2. MAIN RESULT

To prove our theorems, we need the following lemma:

Lemma 1. $f : \Delta = [a, b] \times [c, d] \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ with $0 < a < b$, $0 < c < d$. If $r \in \mathbb{R}$, $r \neq 0$, then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have

$$\begin{aligned} & x^r y^r s^r t^r \int_t^x \int_s^y \frac{1}{u^{r+1} v^{r+1}} [uvf_{uv}(u, v) - ru f_u(u, v) - rv f_v(u, v) + r^2 f(u, v)] dv du \\ &= s^r t^r f(x, y) - y^r t^r f(x, s) - x^r s^r f(t, y) + x^r y^r f(t, s). \end{aligned}$$

Proof. Since f is continuously differentiable function, $\frac{f(u, v)}{u^r v^r}$ is an absolutely continuous function on Δ . Then we get

$$\begin{aligned} (2.1) \quad \int_t^x \int_s^y \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{u^r v^r} \right] dv du &= \int_t^x \frac{\partial}{\partial u} \left(\frac{f(u, y)}{u^r y^r} - \frac{f(u, s)}{u^r s^r} \right) du \\ &= \frac{f(x, y)}{x^r y^r} - \frac{f(x, s)}{x^r s^r} - \frac{f(t, y)}{t^r y^r} + \frac{f(t, s)}{t^r s^r}, \end{aligned}$$

for all $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$. On the other hand,

$$\begin{aligned} (2.2) \quad & \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{u^r v^r} \right] \\ &= \frac{1}{u^{r+1} v^{r+1}} [uvf_{uv}(u, v) - ru f_u(u, v) - rv f_v(u, v) + r^2 f(u, v)] \end{aligned}$$

for all $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$. Thus, from (2.1) and (2.2), it follows that

$$\begin{aligned} & \int_t^x \int_s^y \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{u^r v^r} \right] dv du \\ &= \int_t^x \int_s^y \frac{1}{u^{r+1} v^{r+1}} [uvf_{uv}(u, v) - ru f_u(u, v) - rv f_v(u, v) + r^2 f(u, v)] dv du \\ &= \frac{f(x, y)}{x^r y^r} - \frac{f(x, s)}{x^r s^r} - \frac{f(t, y)}{t^r y^r} + \frac{f(t, s)}{t^r s^r} \end{aligned}$$

which this completes the proof. \square

Theorem 6. $f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ with $0 < a < b$, $0 < c < d$. If $r \in \mathbb{R}$,

Now, consider the case $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, by using Hölder's inequality we have

$$\begin{aligned}
& \left| \int_t^x \int_s^y \frac{1}{u^{r+1}v^{r+1}} |uvf_{uv}(u, v) - ruf_u(u, v) - rvf_v(u, v) + r^2f(u, v)| dvdu \right| \\
& \leq \left| \int_t^x \int_s^y |uvf_{uv}(u, v) - ruf_u(u, v) - rvf_v(u, v) + r^2f(u, v)|^p dvdu \right|^{\frac{1}{p}} \left| \int_t^x \int_s^y \left(\frac{1}{u^{r+1}v^{r+1}} \right)^q dvdu \right|^{\frac{1}{q}} \\
& = \begin{cases} \|l_1f_{uv} - rl_2f_u - rl_3f_v + r^2f\|_p \frac{\left| x^{\frac{1}{q}-r-1} - t^{\frac{1}{q}-r-1} \right| \left| y^{\frac{1}{q}-r-1} - s^{\frac{1}{q}-r-1} \right|}{(1-q(r+1))^{\frac{2}{q}}} & \text{if } r \neq -\frac{1}{p} \\ \|l_1f_{uv} - rl_2f_u - rl_3f_v + r^2f\|_p |\ln x - \ln t| |\ln y - \ln s| & \text{if } r = -\frac{1}{p} \end{cases}
\end{aligned}$$

Finally, we consider the case $p = 1$ and $q = \infty$. Then, we get

$$\begin{aligned}
& \left| \int_t^x \int_s^y \frac{1}{u^{r+1}v^{r+1}} |uvf_{uv}(u, v) - ruf_u(u, v) - rvf_v(u, v) + r^2f(u, v)| dvdu \right| \\
& \leq \left| \int_t^x \int_s^y |uvf_{uv}(u, v) - ruf_u(u, v) - rvf_v(u, v) + r^2f(u, v)| dvdu \right| \sup_{(u,v) \in [t,x] \times [s,y]} \left(\frac{1}{u^{r+1}v^{r+1}} \right) \\
& = \|l_1f_{uv} - rl_2f_u - rl_3f_v + r^2f\|_1 \frac{1}{\min\{t^{r+1}, x^{r+1}\}} \frac{1}{\min\{s^{r+1}, y^{r+1}\}}.
\end{aligned}$$

This completes the proof. \square

Theorem 7. $f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ with $0 < a < b$, $0 < c < d$. If $r \in \mathbb{R}$, $r \neq 0$ and $l_1f_{uv} - rl_2f_u - rl_3f_v + r^2f \in L_\infty(\Delta)$, then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have

$$\begin{aligned}
(2.5) \quad & \left| \frac{(b^{r+1} - a^{r+1})(d^{r+1} - c^{r+1})}{(r+1)^2} f(x, y) - \frac{b^{r+1} - a^{r+1}}{r+1} y^r \int_c^d f(x, s) ds \right. \\
& \left. - \frac{d^{r+1} - c^{r+1}}{r+1} x^r \int_a^b f(t, y) dt + x^r y^r \int_a^b \int_c^d f(t, s) ds dt \right| \\
& \leq \frac{1}{r^2} \|l_1f_{uv} - rl_2f_u - rl_3f_v + r^2f\|_\infty \\
& \quad \times \left[\frac{2rx^{r+1} - (a+b)(r+1)x^r + (a^{r+1} + b^{r+1})}{r+1} \right] \\
& \quad \times \left[\frac{2ry^{r+1} - (c+d)(r+1)y^r + (c^{r+1} + d^{r+1})}{r+1} \right]
\end{aligned}$$

for $r \neq -1$. Also, for $r = -1$, we have

$$(2.6) \quad \left| \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) f(x, y) - y^{-1} \left(\ln \frac{b}{a} \right) \int_c^d f(x, s) ds \right. \\ \left. - x^{-1} \left(\ln \frac{d}{c} \right) \int_a^b f(t, y) dt + x^{-1} y^{-1} \int_a^b \int_c^d f(t, s) ds dt \right| \\ \leq 4 \|l_1 f_{uv} + l_2 f_u + l_3 f_v + f\|_\infty \\ \times \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right) \left(\ln \frac{y}{\sqrt{cd}} + \frac{\frac{c+d}{2} - y}{y} \right)$$

for $(x, y) \in \Delta$, provided $l_1 f_{uv} + l_2 f_u + l_3 f_v + f \in L_\infty(\Delta)$. The constant 4 is best possible in (2.6).

Proof. We observe that

$$\int_a^b \int_c^d [s^r t^r f(x, y) - y^r t^r f(x, s) - x^r s^r f(t, y) + x^r y^r f(t, s)] ds dt \\ = \frac{(b^{r+1} - a^{r+1})(d^{r+1} - c^{r+1})}{(r+1)^2} f(x, y) - \frac{b^{r+1} - a^{r+1}}{r+1} y^r \int_c^d f(x, s) ds \\ - \frac{d^{r+1} - c^{r+1}}{r+1} x^r \int_a^b f(t, y) dt + x^r y^r \int_a^b \int_c^d f(t, s) ds dt.$$

By using the first inequality in (2.3) for $r \neq -1$, it follows that

$$(2.7) \quad \left| \frac{(b^{r+1} - a^{r+1})(d^{r+1} - c^{r+1})}{(r+1)^2} f(x, y) - \frac{b^{r+1} - a^{r+1}}{r+1} y^r \int_c^d f(x, s) ds \right. \\ \left. - \frac{d^{r+1} - c^{r+1}}{r+1} x^r \int_a^b f(t, y) dt + x^r y^r \int_a^b \int_c^d f(t, s) ds dt \right| \\ \leq \int_a^b \int_c^d |s^r t^r f(x, y) - y^r t^r f(x, s) - x^r s^r f(t, y) + x^r y^r f(t, s)| ds dt \\ \leq \int_a^b \int_c^d \frac{x^r y^r s^r t^r}{r^2} \|l_1 f_{uv} - r l_2 f_u - r l_3 f_v + r^2 f\|_\infty \left| \frac{1}{x^r} - \frac{1}{t^r} \right| \left| \frac{1}{y^r} - \frac{1}{s^r} \right| ds dt \\ = \frac{1}{r^2} \|l_1 f_{uv} - r l_2 f_u - r l_3 f_v + r^2 f\|_\infty \left[\int_a^b |t^r - x^r| dt \right] \left[\int_c^d |s^r - y^r| ds \right].$$

Then for $r > 0$, we have

$$\begin{aligned} \int_a^b |t^r - x^r| dt &= \int_a^x (x^r - t^r) dt + \int_x^b (t^r - x^r) dt \\ &= \left[\frac{2rx^{r+1}}{r+1} - (a+b)x^r + \frac{(a^{r+1} + b^{r+1})}{r+1} \right] \end{aligned}$$

and similarly,

$$\begin{aligned} \int_c^d |s^r - y^r| ds &= \int_c^y (y^r - s^r) ds + \int_y^d (s^r - y^r) ds \\ &= \left[\frac{2ry^{r+1}}{r+1} - (c+d)y^r + \frac{(c^{r+1} + d^{r+1})}{r+1} \right]. \end{aligned}$$

If we take $r \in (-\infty, 0) \setminus \{-1\}$, then we obtain

$$\int_a^b |t^r - x^r| dt = - \left[\frac{2rx^{r+1}}{r+1} - (a+b)x^r + \frac{(a^{r+1} + b^{r+1})}{r+1} \right]$$

and similarly,

$$\int_c^d |s^r - y^r| ds = - \left[\frac{2ry^{r+1}}{r+1} - (c+d)y^r + \frac{(c^{r+1} + d^{r+1})}{r+1} \right].$$

Making use of (2.7), we get (2.5).

By using the first inequality in (2.3) for $r = -1$, it follows that

$$\begin{aligned} &|s^{-1}t^{-1}f(x, y) - y^{-1}t^{-1}f(x, s) - x^{-1}s^{-1}f(t, y) + x^{-1}y^{-1}f(t, s)| \\ &\leq \|l_1f_{uv} + l_2f_u + l_3f_v + f\|_\infty |x^{-1} - t^{-1}| |y^{-1} - s^{-1}| \end{aligned}$$

if $l_1f_{uv} + l_2f_u + l_3f_v + f \in L_\infty$. Integrating this inequality, we get

$$\begin{aligned} (2.8) \quad &\left| \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) f(x, y) - y^{-1} \left(\ln \frac{b}{a} \right) \int_c^d f(x, s) ds \right. \\ &\quad \left. - x^{-1} \left(\ln \frac{d}{c} \right) \int_a^b f(t, y) dt + x^{-1} y^{-1} \int_a^b \int_c^d f(t, s) ds dt \right| \\ &\leq \int_a^b \int_c^d |s^{-1}t^{-1}f(x, y) - y^{-1}t^{-1}f(x, s) - x^{-1}s^{-1}f(t, y) + x^{-1}y^{-1}f(t, s)| ds dt \\ &\leq \|l_1f_{uv} + l_2f_u + l_3f_v + f\|_\infty \int_a^b \int_c^d |x^{-1} - t^{-1}| |y^{-1} - s^{-1}| ds dt. \end{aligned}$$

By simple computation,

$$\begin{aligned}
(2.9) \quad \int_a^b |x^{-1} - t^{-1}| dt &= \int_a^b \left| \frac{1}{x} - \frac{1}{t} \right| dt \\
&= \int_a^x \left(\frac{1}{t} - \frac{1}{x} \right) dt + \int_x^b \left(\frac{1}{x} - \frac{1}{t} \right) dt \\
&= 2 \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right)
\end{aligned}$$

and similarly

$$(2.10) \quad \int_c^d |y^{-1} - s^{-1}| ds = 2 \left(\ln \frac{y}{\sqrt{cd}} + \frac{\frac{c+d}{2} - y}{y} \right).$$

By using (2.9) and (2.10) in (2.8), we get the desired inequality (2.6).

Now, we assume that the inequality (2.6) holds with a constant $C > 0$, i.e.

$$\begin{aligned}
(2.11) \quad & \left| \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) f(x, y) - y^{-1} \left(\ln \frac{b}{a} \right) \int_c^d f(x, s) ds \right. \\
& \left. - x^{-1} \left(\ln \frac{d}{c} \right) \int_a^b f(t, y) dt + x^{-1} y^{-1} \int_a^b \int_c^d f(t, s) ds dt \right| \\
& \leq C \|l_1 f_{uv} + l_2 f_u + l_3 f_v + f\|_\infty \\
& \quad \times \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right) \left(\ln \frac{y}{\sqrt{cd}} + \frac{\frac{c+d}{2} - y}{y} \right)
\end{aligned}$$

for any $(x, y) \in \Delta$. If we choose in (2.11), $f(x, s) = f(t, y) = f(t, y) = f(t, s) = 1$, $(t, s) \in \Delta$, then it follows that

$$\begin{aligned}
(2.12) \quad & \left| \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) - \left(\ln \frac{b}{a} \right) \frac{(d-c)}{y} \right. \\
& \left. - \left(\ln \frac{d}{c} \right) \frac{(b-a)}{x} + \frac{(b-a)(d-c)}{xy} \right| \\
& \leq C \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right) \left(\ln \frac{y}{\sqrt{cd}} + \frac{\frac{c+d}{2} - y}{y} \right)
\end{aligned}$$

for any $(x, y) \in \Delta$. Making $x = a$ and $y = c$ in (2.12) produces the inequality

$$\begin{aligned}
& \left| \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) - \left(\ln \frac{b}{a} \right) \frac{(d-c)}{c} \right. \\
& \left. - \left(\ln \frac{d}{c} \right) \frac{(b-a)}{a} + \frac{(b-a)(d-c)}{ac} \right| \\
& \leq C \left(\frac{1}{4} \left(\ln \frac{b}{a} \right) \left(\ln \frac{d}{c} \right) - \left(\ln \frac{b}{a} \right) \frac{(d-c)}{4c} - \left(\ln \frac{d}{c} \right) \frac{(b-a)}{4a} + \frac{(b-a)(d-c)}{4ac} \right)
\end{aligned}$$

which implies that $C \geq 4$. This proves the sharpness of the constant 4 in (2.6). \square

Theorem 8. $f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ with $0 < a < b$, $0 < c < d$. If $r \in \mathbb{R}$, $r \neq 0$ and $l_1 f_{uv} - rl_2 f_u - rl_3 f_v + r^2 f \in L_\infty(\Delta)$, then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have

$$(2.13) \quad \left| \frac{f(x, y)}{x^r y^r} (b-a)(d-c) - \frac{(b-a)}{x^r} \int_c^d \frac{f(x, s)}{s^r} ds \right. \\ \left. - \frac{(d-c)}{y^r} \int_a^b \frac{f(t, y)}{t^r} dt + \int_a^b \int_c^d \frac{f(t, s)}{s^r t^r} ds dt \right| \\ \leq \frac{1}{r^2} \|l_1 f_{uv} - rl_2 f_u - rl_3 f_v + r^2 f\|_\infty \left[\frac{2x^{1-r} - a^{1-r} - b^{1-r}}{1-r} - \frac{(a+b-2x)}{x^r} \right] \\ \times \left[\frac{2y^{1-r} - c^{1-r} - d^{1-r}}{1-r} - \frac{(c+d-2y)}{y^r} \right]$$

for $r \neq 1$. Also, for $r = 1$, we have

$$(2.14) \quad \left| \frac{f(x, y)}{xy} (b-a)(d-c) - \frac{(b-a)}{x} \int_c^d \frac{f(x, s)}{s} ds \right. \\ \left. - \frac{(d-c)}{y} \int_a^b \frac{f(t, y)}{t} dt + \int_a^b \int_c^d \frac{f(t, s)}{st} ds dt \right| \\ \leq 4 \|l_1 f_{uv} - l_2 f_u - l_3 f_v + f\|_\infty \\ \times \left(\ln \frac{x}{\sqrt{ab}} + \frac{\frac{a+b}{2} - x}{x} \right) \left(\ln \frac{y}{\sqrt{cd}} + \frac{\frac{c+d}{2} - y}{y} \right)$$

for $(x, y) \in \Delta$, provided $l_1 f_{uv} - l_2 f_u - l_3 f_v + f \in L_\infty(\Delta)$. The constant 4 is best possible in (2.14).

Proof. By using the inequality (2.4), we get

$$\left| \frac{f(x, y)}{x^r y^r} - \frac{f(x, s)}{x^r s^r} - \frac{f(t, y)}{y^r t^r} + \frac{f(t, s)}{s^r t^r} \right| \\ \leq \left| \int_t^x \int_s^y \frac{1}{u^{r+1} v^{r+1}} |uv f_{uv}(u, v) - ru f_u(u, v) - rv f_v(u, v) + r^2 f(u, v)| dv du \right| \\ \leq \sup_{(u, v) \in [t, x] \times [s, y]} |uv f_{uv}(u, v) - ru f_u(u, v) - rv f_v(u, v) + r^2 f(u, v)| \left| \int_t^x \int_s^y \frac{1}{u^{r+1} v^{r+1}} dv du \right| \\ = \frac{1}{r^2} \|l_1 f_{uv} - rl_2 f_u - rl_3 f_v + r^2 f\|_\infty \left| \frac{1}{x^r} - \frac{1}{t^r} \right| \left| \frac{1}{y^r} - \frac{1}{s^r} \right|$$

for $(x, y), (t, s) \in \Delta$. Integrating this inequality with respect to $(t, s) \in \Delta$, we get

$$\begin{aligned}
(2.15) \quad & \left| \frac{f(x, y)}{x^r y^r} (b-a)(d-c) - \frac{(b-a)}{x^r} \int_c^d \frac{f(x, s)}{s^r} ds \right. \\
& \left. - \frac{(d-c)}{y^r} \int_a^b \frac{f(t, y)}{t^r} dt + \int_a^b \int_c^d \frac{f(t, s)}{s^r t^r} ds dt \right| \\
& \leq \int_a^b \int_c^d \left| \frac{f(x, y)}{x^r y^r} - \frac{f(x, s)}{x^r s^r} - \frac{f(t, y)}{y^r t^r} + \frac{f(t, s)}{s^r t^r} \right| ds dt \\
& \leq \frac{1}{r^2} \|l_1 f_{uv} - r l_2 f_u - r l_3 f_v + r^2 f\|_\infty \int_a^b \int_c^d \left| \frac{1}{x^r} - \frac{1}{t^r} \right| \left| \frac{1}{y^r} - \frac{1}{s^r} \right| ds dt
\end{aligned}$$

for $r \in \mathbb{R}, r \neq 0$. By simple computation, if we take $r \in (0, \infty) \setminus \{1\}$, then we obtain

$$\begin{aligned}
(2.16) \quad & \int_a^b \left| \frac{1}{x^r} - \frac{1}{t^r} \right| dt \\
& = \int_a^x \left(\frac{1}{t^r} - \frac{1}{x^r} \right) dt + \int_x^b \left(\frac{1}{x^r} - \frac{1}{t^r} \right) dt \\
& = \left[\frac{2x^{1-r} - a^{1-r} - b^{1-r}}{1-r} - \frac{(a+b-2x)}{x^r} \right]
\end{aligned}$$

and similarly

$$(2.17) \quad \int_c^d \left| \frac{1}{y^r} - \frac{1}{s^r} \right| ds = \left[\frac{2y^{1-r} - c^{1-r} - d^{1-r}}{1-r} - \frac{(c+d-2y)}{y^r} \right].$$

for any $(x, y) \in \Delta$. On the other hand, for $r > 0$, we have

$$(2.18) \quad \int_a^b \left| \frac{1}{x^r} - \frac{1}{t^r} \right| dt = - \left[\frac{2x^{1-r} - a^{1-r} - b^{1-r}}{1-r} - \frac{(a+b-2x)}{x^r} \right]$$

and

$$(2.19) \quad \int_c^d \left| \frac{1}{y^r} - \frac{1}{s^r} \right| ds = - \left[\frac{2y^{1-r} - c^{1-r} - d^{1-r}}{1-r} - \frac{(c+d-2y)}{y^r} \right]$$

for any $(x, y) \in \Delta$. By using (2.16)-(2.19) in (2.15), we get the desired inequality (2.13).

For $r = 1$, proceeding as in the proof of Theorem 7, we also obtain the inequality (2.14). \square

REFERENCES

- [1] A. M. Acu, A. Babos and F. D. Sofonea, *The mean value theorems and inequalities of Ostrowski type*. Sci. Stud. Res. Ser. Math. Inform. 21 (2011), no. 1, 5–16.

- [2] S.S. Dragomir, *An inequality of Ostrowski type via Pompeiu's mean value theorem*, J. of Inequal. in Pure and Appl. Math., 6(3) (2005), Art. 83.
- [3] I. Muntean, *Extensions of some mean value theorems*, Babes-Bolyai University, Faculty of Mathematics, Research Seminars on Mathematical Analysis, Preprint Nr. 7, 1991, 7-24.
- [4] F. Ahmad, N. S. Barnett and S. S. Dragomir, *New Weighted Ostrowski and Cebyshev Type Inequalities*, Nonlinear Analysis: Theory, Methods & Appl., 71 (12), (2009), 1408-1412.
- [5] F. Ahmad, A. Rafiq, N. A. Mir, *Weighted Ostrowski type inequality for twice differentiable mappings*, Global Journal of Research in Pure and Applied Math., 2 (2) (2006), 147-154.
- [6] N. S. Barnett and S. S. Dragomir, *An Ostrowski type inequality for double integrals and applications for cubature formulae*, Soochow J. Math., 27(1), (2001), 109-114.
- [7] J. Pečarić and Š. Ungar, *On an inequality of Ostrowski type*, J. Ineq. Pure & Appl. Math. 7 (2006). No. 4. Art. 151.
- [8] D. Pompeiu, *Sur une proposition analogue au théorème des accroissements finis*, Mathematica (Cluj, Romania), 22(1946), 143-146.
- [9] E. C. Popa, *An inequality of Ostrowski type via a mean value theorem*, General Mathematics Vol. 15, No. 1, 2007, 93-100.
- [10] S. S. Dragomir, *Power Pompeiu's type inequalities for absolutely continuous functions with applications to Ostrowski's inequality*,
- [11] S. Hussain, M.A.Latif and M. Alomari, *Generalized duple-integral Ostrowski type inequalities on time scales*, Appl. Math. Letters, 24(2011), 1461-1467.
- [12] A. M. Ostrowski, *Über die absolutabweichung einer differentiebaren funktion von ihrem integralmittelwert*, Comment. Math. Helv. 10(1938), 226-227.
- [13] A. Qayyum, *A weighted Ostrowski-Grüss type inequality and applications*, Proceeding of the World Cong. on Engineering, Vol:2, 2009, 1-9.
- [14] A. Rafiq and F. Ahmad, *Another weighted Ostrowski-Grüss type inequality for twice differentiable mappings*, Kragujevac Journal of Mathematics, 31 (2008), 43-51.
- [15] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [16] M. Z. Sarikaya *On the Ostrowski type integral inequality for double integrals*, Demonstratio Mathematica, Vol. XLV, No 3, 2012, pp. 533-540.
- [17] M. Z. Sarikaya and H. Ogunmez, *On the weighted Ostrowski type integral inequality for double integrals*, The Arabian Journal for Science and Engineering (AJSE)-Mathematics, (2011) 36:1153-1160.
- [18] F. Ahmad, N. A. Mir and M.Z. Sarikaya, *An inequality of Ostrowski type via variant of Pompeiu's mean value theorem*, J. Basic. Appl. Sci. Res., 4(4)204-211, 2014.
- [19] M. Z. Sarikaya and H. Budak, *On an inequality of Ostrowski type via variant of Pompeiu's mean value theorem*, RGMIA Research Report Collection, 17(2014), Article 78, 11 pp.
- [20] M. Z. Sarikaya, *On an inequality of Grüss type via variant of Pompeiu's mean value theorem*, RGMIA Research Report Collection, 17(2014), Article 77, 9 pp.
- [21] M. Z. Sarikaya, *Some new integral inequalities via variant of Pompeiu's mean value theorem*, RGMIA Research Report Collection, 17(2014), Article 76, 7 pp.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, KUNURRALP CAMPUS, DÜZCE-TURKEY
E-mail address: sarikayamz@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BARTIN UNIVERSITY, BARTIN-TURKEY
E-mail address: erdem1627@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, KUNURRALP CAMPUS, DÜZCE-TURKEY
E-mail address: yaldizhatice@gmail.com