

**ON NEW INEQUALITIES OF HERMITE-HADAMARD-FEJER
TYPE FOR HARMONICALLY CONVEX FUNCTIONS VIA
FRACTIONAL INTEGRALS**

MEHMET KUNT AND İMDAT İŞCAN

ABSTRACT. In this paper, firstly, new Hermite-Hadamard type inequality for harmonically convex function in fractional integral forms was given. Secondly, Hermite-Hadamard-Fejer inequality for harmonically convex functions in fractional integral forms was built. Finally, an integral identity and some Hermite-Hadamard-Fejer type integral inequalities for harmonically convex functions in fractional integral forms were obtained. The some results presented here would provide extensions of those given in earlier works.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \quad (1.1)$$

is well known in the literature as Hermite-Hadamard's inequality [5].

The most well-known inequalities related to the integral mean of a convex function f are the Hermite Hadamard inequalities or its weighted versions, the so-called Hermite-Hadamard-Fejér inequalities.

In [4], Fejér established the following Fejér inequality which is the weighted generalization of Hermite-Hadamard inequality (1.1):

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be convex function. Then the inequality*

$$f\left(\frac{a+b}{2}\right) \int_a^b g(x)dx \leq \int_a^b f(x)g(x)dx \leq \frac{f(a)+f(b)}{2} \int_a^b g(x)dx \quad (1.2)$$

holds, where $g : [a, b] \rightarrow \mathbb{R}$ is nonnegative, integrable and symmetric to $(a+b)/2$.

For some results which generalize, improve, and extend the inequalities (1.1) and (1.2) see [1, 6, 7, 15, 17].

We recall the following inequality and special functions which are known as Beta and hypergeometric function respectively

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad x, y > 0,$$

$${}_2F_1(a, b; c; z) = \frac{1}{\beta(b, c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-zt)^{-a} dt,$$

$$c > b > 0, |z| < 1 \text{ (see [12])}.$$

2000 *Mathematics Subject Classification.* 26A51, 26A33, 26D10.

Key words and phrases. Hermite-Hadamard inequality, Hermite-Hadamard-Fejer inequality, Riemann-Liouville fractional integral, harmonically convex function.

Lemma 1. [14, 19]. For $0 < \alpha \leq 1$ and $0 \leq a < b$ we have

$$|a^\alpha - b^\alpha| \leq (b - a)^\alpha.$$

Following definitions and mathematical preliminaries of fractional calculus theory are used further in this paper.

Definition 1. [12]. Let $f \in L[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively, where $\Gamma(\alpha)$ is the Gamma function defined by $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$ and $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

Because of the wide application of Hermite-Hadamard type inequalities and fractional integrals, many researchers extend their studies to Hermite-Hadamard type inequalities involving fractional integrals not limited to integer integrals. Recently, more and more Hermite-Hadamard inequalities involving fractional integrals have been obtained for different classes of functions; see [3, 8, 9, 16, 18, 19].

In [11], İşcan gave definition of harmonically convex functions and established following Hermite-Hadamard type inequality for harmonically convex functions as follows:

Definition 2. Let $I \subset \mathbb{R} \setminus \{0\}$ be a real interval. A function $f : I \rightarrow \mathbb{R}$ is said to be harmonically convex, if

$$f\left(\frac{xy}{tx + (1-t)y}\right) \leq tf(y) + (1-t)f(x) \quad (1.3)$$

for all $x, y \in I$ and $t \in [0, 1]$. If the inequality in (1.3) is reversed, then f is said to be harmonically concave.

Theorem 2. [11]. Let $f : I \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a harmonically convex function and $a, b \in I$ with $a < b$. If $f \in L[a, b]$ then the following inequalities hold:

$$f\left(\frac{2ab}{a+b}\right) \leq \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \leq \frac{f(a) + f(b)}{2}. \quad (1.4)$$

In [13] Latif et. al. gave the following definition:

Definition 3. A function $g : [a, b] \subseteq \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ is said to be harmonically symmetric with respect to $2ab/a+b$ if

$$g(x) = g\left(\frac{1}{\frac{1}{a} + \frac{1}{b} - \frac{1}{x}}\right)$$

holds for all $x \in [a, b]$.

In [2] Chan and Wu represented Hermite-Hadamard-Fejér inequality for harmonically convex functions as follows:

Theorem 3. Let $f : I \subseteq \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a harmonically convex function and $a, b \in I$ with $a < b$. If $f \in L[a, b]$ and $g : [a, b] \subseteq \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ is nonnegative, integrable and harmonically symmetric with respect to $\frac{2ab}{a+b}$, then

$$\begin{aligned} f\left(\frac{2ab}{a+b}\right) \int_a^b \frac{g(x)}{x^2} dx &\leq \int_a^b \frac{f(x)g(x)}{x^2} dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b \frac{g(x)}{x^2} dx. \end{aligned} \quad (1.5)$$

In this paper, we gave new Hermite-Hadamard type inequality for harmonically convex function in fractional integral forms. We established new Hermite-Hadamard-Fejer inequality for harmonically convex function in fractional integral forms. We obtained an integral identity and some Hermite-Hadamard-Fejer type integral inequalities for harmonically convex functions in fractional integral forms.

2. MAIN RESULTS

Throughout this section, we take $\|g\|_\infty = \sup_{t \in [a, b]} |g(x)|$, for the continuous function $g : [a, b] \rightarrow \mathbb{R}$.

Lemma 2. If $g : [a, b] \subseteq \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ is integrable and harmonically symmetric with respect to $\frac{2ab}{a+b}$ with $a < b$, then

$$J_{\frac{2ab}{2ab}+}^\alpha (g \circ h)(1/a) = J_{\frac{2ab}{2ab}-}^\alpha (g \circ h)(1/b) = \frac{1}{2} \left[\begin{array}{l} J_{\frac{2ab}{2ab}+}^\alpha (g \circ h)(1/a) \\ + J_{\frac{2ab}{2ab}-}^\alpha (g \circ h)(1/b) \end{array} \right]$$

with $\alpha > 0$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. Since g is harmonically symmetric with respect to $\frac{2ab}{a+b}$, using Definition 3 we have $g(\frac{1}{x}) = g(\frac{1}{(\frac{1}{a})+(\frac{1}{b})-x})$, for all $x \in [\frac{1}{b}, \frac{1}{a}]$. Hence, in the following integral setting $t = (\frac{1}{a}) + (\frac{1}{b}) - x$ and $dt = -dx$ gives

$$\begin{aligned} J_{\frac{2ab}{2ab}+}^\alpha (g \circ h)(1/a) &= \frac{1}{\Gamma(\alpha)} \int_{\frac{1}{a}}^{\frac{1}{b}} \left(\frac{1}{a} - t\right)^{\alpha-1} g\left(\frac{1}{t}\right) dt \\ &= \frac{1}{\Gamma(\alpha)} \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b}\right)^{\alpha-1} g\left(\frac{1}{(1/a) + (1/b) - x}\right) dx \\ &= \frac{1}{\Gamma(\alpha)} \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b}\right)^{\alpha-1} g\left(\frac{1}{x}\right) dx = J_{\frac{2ab}{2ab}-}^\alpha (g \circ h)(1/b). \end{aligned}$$

This completes the proof. \square

Theorem 4. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a function such that $f \in L[a, b]$, where $a, b \in I$ with $a < b$. If f is a harmonically convex function on $[a, b]$, then the following inequalities for fractional integrals hold:

$$\begin{aligned} f\left(\frac{2ab}{a+b}\right) &\leq \frac{\Gamma(\alpha+1)}{2^{1-\alpha}} \left(\frac{ab}{b-a}\right)^\alpha \left\{ \begin{array}{l} J_{\frac{2ab}{2ab}+}^\alpha (f \circ h)(1/a) \\ + J_{\frac{2ab}{2ab}-}^\alpha (f \circ h)(1/b) \end{array} \right\} \\ &\leq \frac{f(a)+f(b)}{2} \end{aligned} \quad (2.1)$$

with $\alpha > 0$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. Since f is a harmonically convex function on $[a, b]$, we have for all $t \in [0, 1]$

$$\begin{aligned} f\left(\frac{2ab}{a+b}\right) &= f\left(\frac{2\left(\frac{ab}{ta+(1-t)b}\right)\left(\frac{ab}{tb+(1-t)a}\right)}{\left(\frac{ab}{ta+(1-t)b}\right) + \left(\frac{ab}{tb+(1-t)a}\right)}\right) \\ &\leq \frac{f\left(\frac{ab}{ta+(1-t)b}\right) + f\left(\frac{ab}{tb+(1-t)a}\right)}{2}. \end{aligned} \quad (2.2)$$

Multiplying both sides of (2.2) by $2t^{\alpha-1}$ then integrating the resulting inequality with respect to t over $[0, \frac{1}{2}]$, we obtain

$$\begin{aligned} &2f\left(\frac{2ab}{a+b}\right) \int_0^{\frac{1}{2}} t^{\alpha-1} dt \\ &\leq \int_0^{\frac{1}{2}} t^{\alpha-1} \left[f\left(\frac{ab}{ta+(1-t)b}\right) + f\left(\frac{ab}{tb+(1-t)a}\right) \right] dt \\ &= \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{ta+(1-t)b}\right) dt + \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{tb+(1-t)a}\right) dt. \end{aligned}$$

Setting $x = \frac{tb+(1-t)a}{ab}$, and $dx = \left(\frac{b-a}{ab}\right) dt$ gives

$$\begin{aligned} &\frac{2^{1-\alpha}}{\alpha} f\left(\frac{2ab}{a+b}\right) \\ &\leq \left(\frac{ab}{b-a}\right)^\alpha \left\{ \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} (x - \frac{1}{b})^{\alpha-1} f\left(\frac{1}{\frac{1}{a} + \frac{1}{b} - x}\right) dx \right. \\ &\quad \left. + \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} (x - \frac{1}{b})^{\alpha-1} f\left(\frac{1}{x}\right) dx \right\} \\ &= \left(\frac{ab}{b-a}\right)^\alpha \left\{ \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} (\frac{1}{a} - x)^{\alpha-1} f\left(\frac{1}{x}\right) dx \right. \\ &\quad \left. + \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} (x - \frac{1}{b})^{\alpha-1} f\left(\frac{1}{x}\right) dx \right\} \\ &= \left(\frac{ab}{b-a}\right)^\alpha \Gamma(\alpha) \left[J_{\frac{a+b}{2ab}+}^\alpha (f \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (f \circ h)(1/b) \right] \end{aligned}$$

and the first inequality is proved.

For the proof of the second inequality in (2.1), we first note that, if f is a harmonically convex function, then, for all $t \in [0, 1]$, it yields

$$f\left(\frac{ab}{ta+(1-t)b}\right) + f\left(\frac{ab}{tb+(1-t)a}\right) \leq f(a) + f(b). \quad (2.3)$$

Then multiplying both sides of (2.3) by $t^{\alpha-1}$ and integrating the resulting inequality with respect to t over $[0, \frac{1}{2}]$, we obtain

$$\begin{aligned} &\int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{ta+(1-t)b}\right) dt + \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{tb+(1-t)a}\right) dt \\ &\leq [f(a) + f(b)] \int_0^{\frac{1}{2}} t^{\alpha-1} dt = \frac{2^{1-\alpha}}{\alpha} \frac{f(a) + f(b)}{2} \end{aligned}$$

i.e.

$$\begin{aligned} &\left(\frac{ab}{b-a}\right)^\alpha \Gamma(\alpha) \left[J_{\frac{a+b}{2ab}+}^\alpha (f \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (f \circ h)(1/b) \right] \\ &\leq \frac{2^{1-\alpha}}{\alpha} \left(\frac{f(a) + f(b)}{2} \right) \end{aligned}$$

The proof is completed. \square

Theorem 5. Let $f : [a, b] \rightarrow \mathbb{R}$ be harmonically convex function with $a < b$ and $f \in L[a, b]$. If $g : [a, b] \rightarrow \mathbb{R}$ is nonnegative, integrable and harmonically symmetric with respect to $\frac{2ab}{a+b}$, then the following inequalities for fractional integrals hold:

$$\begin{aligned} & f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{2ab}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \\ & \leq \left[J_{\frac{2ab}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \\ & \leq \frac{f(a) + f(b)}{2} \left[J_{\frac{2ab}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \end{aligned} \quad (2.4)$$

with $\alpha > 0$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. Since f is a harmonically convex function on $[a, b]$, multiplying both sides of (2.2) by $2t^{\alpha-1}g\left(\frac{ab}{tb+(1-t)a}\right)$ then integrating the resulting inequality with respect to t over $[0, \frac{1}{2}]$, we obtain

$$\begin{aligned} & 2f\left(\frac{2ab}{a+b}\right) \int_0^{\frac{1}{2}} t^{\alpha-1} g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & \leq \int_0^{\frac{1}{2}} t^{\alpha-1} \left[f\left(\frac{ab}{ta+(1-t)b}\right) + f\left(\frac{ab}{tb+(1-t)a}\right) \right] g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & = \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{ta+(1-t)b}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & \quad + \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{tb+(1-t)a}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt. \end{aligned}$$

Since g is harmonically symmetric with respect to $\frac{2ab}{a+b}$, using Definition 3 we have $g\left(\frac{1}{x}\right) = g\left(\frac{1}{\left(\frac{1}{a}\right) + \left(\frac{1}{b}\right) - x}\right)$, for all $x \in [\frac{1}{b}, \frac{1}{a}]$. Setting $x = \frac{tb+(1-t)a}{ab}$, and $dx = \left(\frac{b-a}{ab}\right) dt$ gives

$$\begin{aligned} & 2\left(\frac{ab}{b-a}\right)^{\alpha} f\left(\frac{2ab}{a+b}\right) \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(x - \frac{1}{b}\right)^{\alpha-1} g\left(\frac{1}{x}\right) dx \\ & \leq \left(\frac{ab}{b-a}\right)^{\alpha} \left\{ \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(x - \frac{1}{b}\right)^{\alpha-1} f\left(\frac{1}{\left(\frac{1}{a}\right) + \left(\frac{1}{b}\right) - x}\right) g\left(\frac{1}{x}\right) dx \right. \\ & \quad \left. + \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(x - \frac{1}{b}\right)^{\alpha-1} f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx \right\} \\ & = \left(\frac{ab}{b-a}\right)^{\alpha} \left\{ \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{1}{a} - x\right)^{\alpha-1} f\left(\frac{1}{x}\right) g\left(\frac{1}{\left(\frac{1}{a}\right) + \left(\frac{1}{b}\right) - x}\right) dx \right. \\ & \quad \left. + \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(x - \frac{1}{b}\right)^{\alpha-1} f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx \right\} \\ & = \left(\frac{ab}{b-a}\right)^{\alpha} \left\{ \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{1}{a} - x\right)^{\alpha-1} f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx \right. \\ & \quad \left. + \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(x - \frac{1}{b}\right)^{\alpha-1} f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx \right\} \end{aligned}$$

Therefore, by Lemma 2 we have

$$\begin{aligned} & \left(\frac{ab}{b-a}\right)^{\alpha} \Gamma(\alpha) f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{2ab}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \\ & \leq \left(\frac{ab}{b-a}\right)^{\alpha} \Gamma(\alpha) \left[J_{\frac{2ab}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \end{aligned}$$

and the first inequality is proved.

For the proof of the second inequality in (2.4) we first note that if f is a harmonically convex function, then, multiplying both sides of (2.3) by $t^{\alpha-1}g\left(\frac{ab}{tb+(1-t)a}\right)$

and integrating the resulting inequality with respect to t over $[0, \frac{1}{2}]$, we obtain

$$\begin{aligned} & \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{ta+(1-t)b}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & + \int_0^{\frac{1}{2}} t^{\alpha-1} f\left(\frac{ab}{tb+(1-t)a}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & \leq [f(a) + f(b)] \int_0^{\frac{1}{2}} t^{\alpha-1} g\left(\frac{ab}{tb+(1-t)a}\right) dt \end{aligned}$$

i.e.

$$\begin{aligned} & \left(\frac{ab}{b-a}\right)^\alpha \Gamma(\alpha) \left[J_{\frac{a+b}{2ab}+}^\alpha (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (fg \circ h)(1/b) \right] \\ & \leq \left(\frac{ab}{b-a}\right)^\alpha \Gamma(\alpha) \left(\frac{f(a) + f(b)}{2}\right) \left[J_{\frac{a+b}{2ab}+}^\alpha (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (g \circ h)(1/b) \right] \end{aligned}$$

The proof is completed. \square

Remark 1. In Theorem 5,

- (i) if we take $\alpha = 1$, then inequality (2.4) becomes inequality (1.5) of Theorem 3.
- (ii) if we take $g(x) = 1$, then inequality (2.4) becomes inequality (2.1) of Theorem 4.
- (iii) if we take $\alpha = 1$ and $g(x) = 1$, then inequality (2.4) becomes inequality (1.4) of Theorem 2.

Lemma 3. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ and $a < b$. If $g : [a, b] \rightarrow \mathbb{R}$ is integrable and harmonically symmetric with respect to $\frac{2ab}{a+b}$, then the following equality for fractional integrals hold:

$$\begin{aligned} & f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{a+b}{2ab}+}^\alpha (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (g \circ h)(1/b) \right] \\ & - \left[J_{\frac{a+b}{2ab}+}^\alpha (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^\alpha (fg \circ h)(1/b) \right] \\ & = \frac{1}{\Gamma(\alpha)} \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_{\frac{1}{b}}^t (s - \frac{1}{b})^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)'(t) dt \right. \\ & \quad \left. - \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} (\frac{1}{a} - s)^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)'(t) dt \right] \quad (2.5) \end{aligned}$$

with $\alpha > 0$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. It suffices to note that

$$\begin{aligned} I & = \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_{\frac{1}{b}}^t \left(s - \frac{1}{b}\right)^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ & \quad - \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(\frac{1}{a} - s\right)^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ & = I_1 - I_2. \end{aligned}$$

By integration by parts and Lemma 2 we get

$$\begin{aligned}
I_1 &= \left(\int_{\frac{1}{b}}^t \left(s - \frac{1}{b} \right)^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)(t) \Big|_{\frac{1}{b}}^{\frac{a+b}{2ab}} \\
&\quad - \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(t - \frac{1}{b} \right)^{\alpha-1} (g \circ h)(t) (f \circ h)(t) dt \\
&= \left(\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(s - \frac{1}{b} \right)^{\alpha-1} (g \circ h)(s) ds \right) f\left(\frac{2ab}{a+b}\right) \\
&\quad - \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(t - \frac{1}{b} \right)^{\alpha-1} (g \circ h)(t) (f \circ h)(t) dt \\
&= \Gamma(\alpha) \left[f\left(\frac{2ab}{a+b}\right) J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) - J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \\
&= \Gamma(\alpha) \left[\frac{f\left(\frac{2ab}{a+b}\right)}{2} \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\
&\quad \left. - J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right]
\end{aligned}$$

and similarly

$$\begin{aligned}
I_2 &= \left(\int_t^{\frac{1}{a}} \left(\frac{1}{a} - s \right)^{\alpha-1} (g \circ h)(s) ds \right) (f \circ h)(t) \Big|_{\frac{a+b}{2ab}}^{\frac{1}{a}} \\
&\quad + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{1}{a} - t \right)^{\alpha-1} (g \circ h)(t) (f \circ h)(t) dt \\
&= - \left(\int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{1}{a} - s \right)^{\alpha-1} (g \circ h)(s) ds \right) f\left(\frac{2ab}{a+b}\right) \\
&\quad + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{1}{a} - t \right)^{\alpha-1} (g \circ h)(t) (f \circ h)(t) dt \\
&= \Gamma(\alpha) \left[-f\left(\frac{2ab}{a+b}\right) J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) \right] \\
&= \Gamma(\alpha) \left[-\frac{f\left(\frac{2ab}{a+b}\right)}{2} \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\
&\quad \left. + J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) \right].
\end{aligned}$$

Thus, we can write

$$I = I_1 - I_2 = \Gamma(\alpha) \left\{ \begin{aligned} & f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \\ & - \left[J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \end{aligned} \right\}.$$

Multiplying the both sides by $(\Gamma(\alpha))^{-1}$ we obtain (2.5), which completes the proof. \square

Theorem 6. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ and $a < b$. If $|f'|$ is harmonically convex on $[a, b]$, $g : [a, b] \rightarrow \mathbb{R}$ is continuous and harmonically symmetric with respect to $\frac{2ab}{a+b}$, then

the following inequality for fractional integrals hold:

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\ & \quad \left. - \left[J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \right| \\ & \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} [C_1(\alpha) |f'(a)| + C_2(\alpha) |f'(b)|] \end{aligned} \quad (2.6)$$

where

$$\begin{aligned} C_1(\alpha) &= \left[\begin{aligned} & \frac{b^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; 1-\frac{a}{b}\right) \\ & - \frac{(a+b)^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; \frac{b-a}{b+a}\right) \end{aligned} \right] \\ C_2(\alpha) &= \left[\begin{aligned} & \frac{b^{-2}}{\alpha+2} {}_2F_1\left(2, \alpha+2; \alpha+3; 1-\frac{a}{b}\right) \\ & - \frac{2(a+b)^{-2}}{\alpha+1} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{b-a}{b+a}\right) \\ & + \frac{(a+b)^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; \frac{b-a}{b+a}\right) \end{aligned} \right] \end{aligned}$$

with $0 < \alpha \leq 1$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. From Lemma 3 we have

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\ & \quad \left. - \left[J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \right| \\ & \leq \frac{1}{\Gamma(\alpha)} \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_{\frac{1}{b}}^t \left(s - \frac{1}{b}\right)^{\alpha-1} |(g \circ h)(s)| ds \right) |(f \circ h)'(t)| dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(\frac{1}{a} - s\right)^{\alpha-1} |(g \circ h)(s)| ds \right) |(f \circ h)'(t)| dt \right] \\ & \leq \frac{\|g\|_{\infty}}{\Gamma(\alpha)} \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_{\frac{1}{b}}^t \left(s - \frac{1}{b}\right)^{\alpha-1} ds \right) |(f \circ h)'(t)| dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(\frac{1}{a} - s\right)^{\alpha-1} ds \right) |(f \circ h)'(t)| dt \right] \\ & = \frac{\|g\|_{\infty}}{\Gamma(\alpha)} \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \frac{\left(t - \frac{1}{b}\right)^{\alpha}}{\alpha} \frac{1}{t^2} |f'(\frac{1}{t})| dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \frac{\left(\frac{1}{a} - t\right)^{\alpha}}{\alpha} \frac{1}{t^2} |f'(\frac{1}{t})| dt \right] \end{aligned}$$

Setting $t = \frac{ub+(1-u)a}{ab}$, and $dt = \left(\frac{b-a}{ab}\right) du$ gives

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{a+b}{2ab}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\ & \quad \left. - \left[J_{\frac{a+b}{2ab}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{a+b}{2ab}-}^{\alpha} (fg \circ h)(1/b) \right] \right| \\ & \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \left[\int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \right] \end{aligned} \quad (2.7)$$

Since $|f'|$ is harmonically convex on $[a, b]$, we have

$$\left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| \leq u |f'(a)| + (1-u) |f'(b)| \quad (2.8)$$

If we use (2.8) in (2.7), we have

$$\begin{aligned}
& \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{2ab}{a+b}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{2ab}{a+b}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\
& \quad \left. - \left[J_{\frac{2ab}{a+b}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{2ab}{a+b}-}^{\alpha} (fg \circ h)(1/b) \right] \right| \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \\
& \quad \times \left[\int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} [u|f'(a)| + (1-u)|f'(b)|] du \right. \\
& \quad \left. + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} [u|f'(a)| + (1-u)|f'(b)|] du \right] \quad (2.9)
\end{aligned}$$

Calculating following integrals by Lemma 1, we have

$$\begin{aligned}
& \int_0^{\frac{1}{2}} \frac{u^{\alpha+1}}{(ub+(1-u)a)^2} du + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha} u}{(ub+(1-u)a)^2} du \\
& = \int_0^1 \frac{(1-u)^{\alpha} u}{(ub+(1-u)a)^2} du - \int_0^{\frac{1}{2}} \frac{(1-u)^{\alpha} - u^{\alpha}}{(ub+(1-u)a)^2} u du \\
& \leq \int_0^1 \frac{(1-u)^{\alpha} u}{(ub+(1-u)a)^2} du - \int_0^{\frac{1}{2}} \frac{(1-2u)^{\alpha}}{(ub+(1-u)a)^2} u du \\
& = \int_0^1 \frac{(1-u)^{\alpha} u}{(ub+(1-u)a)^2} du - \frac{1}{4} \int_0^1 \frac{(1-u)^{\alpha}}{(\frac{u}{2}b + (1-\frac{u}{2})a)^2} u du \\
& = \int_0^1 (1-u) u^{\alpha} b^{-2} \left(1-u\left(1-\frac{a}{b}\right)\right)^{-2} du \\
& \quad - \frac{1}{4} \int_0^1 (1-v) v^{\alpha} \left(\frac{a+b}{2}\right)^{-2} \left(1-v\left(\frac{b-a}{b+a}\right)\right)^{-2} dv \\
& = \left[\begin{aligned} & \frac{b^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; 1-\frac{a}{b}\right) \\ & - \frac{(a+b)^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; \frac{b-a}{b+a}\right) \end{aligned} \right] \\
& = C_1(\alpha) \quad (2.10)
\end{aligned}$$

and similarly we get

$$\begin{aligned}
& \int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} (1-u) du + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} (1-u) du \\
& = \int_0^1 \frac{(1-u)^{\alpha+1}}{(ub+(1-u)a)^2} du - \int_0^{\frac{1}{2}} \frac{(1-u)^{\alpha} - u^{\alpha}}{(ub+(1-u)a)^2} (1-u) du \\
& \leq \int_0^1 \frac{(1-u)^{\alpha+1}}{(ub+(1-u)a)^2} du - \int_0^{\frac{1}{2}} \frac{(1-2u)^{\alpha}}{(ub+(1-u)a)^2} (1-u) du \\
& = \int_0^1 \frac{(1-u)^{\alpha+1}}{(ub+(1-u)a)^2} du - \int_0^{\frac{1}{2}} \frac{(1-2u)^{\alpha}}{(ub+(1-u)a)^2} du \\
& \quad + \int_0^{\frac{1}{2}} \frac{u(1-2u)^{\alpha}}{(ub+(1-u)a)^2} du
\end{aligned}$$

$$\begin{aligned}
&= \int_0^1 \frac{u^{\alpha+1}}{(ua + (1-u)b)^2} du - \frac{1}{2} \int_0^1 \frac{(1-u)^\alpha}{\left(\frac{u}{2}b + \left(1 - \frac{u}{2}\right)a\right)^2} du \\
&\quad + \frac{1}{4} \int_0^1 \frac{u(1-u)^\alpha}{\left(\frac{u}{2}b + \left(1 - \frac{u}{2}\right)a\right)^2} du \\
&= \int_0^1 \frac{u^{\alpha+1}}{(ua + (1-u)b)^2} du - \frac{1}{2} \int_0^1 v^\alpha \left(\frac{a+b}{2}\right)^{-2} \left(1 - v \left(\frac{b-a}{b+a}\right)\right)^{-2} dv \\
&\quad + \frac{1}{4} \int_0^1 (1-v) v^\alpha \left(\frac{a+b}{2}\right)^{-2} \left(1 - v \left(\frac{b-a}{b+a}\right)\right)^{-2} dv \\
&= \left[\begin{array}{l} \frac{b^{-2}}{\alpha+2} {}_2F_1\left(2, \alpha+2; \alpha+3; 1 - \frac{a}{b}\right) \\ - \frac{2(a+b)^{-2}}{\alpha+1} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{b-a}{b+a}\right) \\ + \frac{(a+b)^{-2}}{(\alpha+1)(\alpha+2)} {}_2F_1\left(2, \alpha+1; \alpha+3; \frac{b-a}{b+a}\right) \end{array} \right] \\
&= C_2(\alpha) \tag{2.11}
\end{aligned}$$

If we use (2.10) and (2.11) in (2.9), we have (2.6). This completes the proof. \square

Corollary 1. *In Theorem 6;*

(1) *If we take $\alpha = 1$ we have the following Hermite-Hadamard-Fejer inequality for harmonically convex functions which is related the left-hand side of (1.5):*

$$\begin{aligned}
&\left| f\left(\frac{2ab}{a+b}\right) \int_a^b \frac{g(x)}{x^2} dx - \int_a^b \frac{f(x)g(x)}{x^2} dx \right| \\
&\leq \|g\|_\infty (b-a)^2 [C_1(1)|f'(a)| + C_2(1)|f'(b)|],
\end{aligned}$$

(2) *If we take $g(x) = 1$ we have following Hermite-Hadamard type inequality for harmonically convex function in fractional integral forms which is related the left-hand side of (2.1):*

$$\begin{aligned}
&\left| f\left(\frac{2ab}{a+b}\right) - \frac{\Gamma(\alpha+1)}{2^{1-\alpha}} \left(\frac{ab}{b-a}\right)^\alpha \left\{ \begin{array}{l} J_{\frac{a+b}{2ab}^+}^\alpha (f \circ h)(1/a) \\ + J_{\frac{a+b}{2ab}^-}^\alpha (f \circ h)(1/b) \end{array} \right\} \right| \\
&\leq \frac{ab(b-a)}{2^{1-\alpha}} [C_1(\alpha)|f'(a)| + C_2(\alpha)|f'(b)|],
\end{aligned}$$

(3) *If we take $\alpha = 1$ and $g(x) = 1$ we have the following Hermite-Hadamard type inequality for harmonically convex function which is related the left-hand side of (1.4):*

$$\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq ab(b-a) [C_1(1)|f'(a)| + C_2(1)|f'(b)|].$$

Theorem 7. *Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ and $a < b$. If $|f'|^q, q \geq 1$, is harmonically convex on $[a, b]$, $g : [a, b] \rightarrow \mathbb{R}$ is continuous and harmonically symmetric with respect to $\frac{2ab}{a+b}$,*

then the following inequality for fractional integrals hold:

$$\begin{aligned}
& \left| \frac{f(a)+f(b)}{2} \left[J_{1/b+}^{\alpha} (g \circ h) (1/a) + J_{1/a-}^{\alpha} (g \circ h) (1/b) \right] \right. \\
& \quad \left. - \left[J_{1/b+}^{\alpha} (fg \circ h) (1/a) + J_{1/a-}^{\alpha} (fg \circ h) (1/b) \right] \right| \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab} \right)^{\alpha} \\
& \quad \times \left[\begin{aligned} & C_3^{1-\frac{1}{q}}(\alpha) \left[\left(\begin{aligned} & C_4(\alpha) |f'(a)|^q \\ & + C_5(\alpha) |f'(b)|^q \end{aligned} \right) \right]^{\frac{1}{q}} \\ & + C_6^{1-\frac{1}{q}}(\alpha) \left[\left(\begin{aligned} & C_7(\alpha) |f'(a)|^q \\ & + C_8(\alpha) |f'(b)|^q \end{aligned} \right) \right]^{\frac{1}{q}} \end{aligned} \right] \quad (2.12)
\end{aligned}$$

where

$$\begin{aligned}
C_3(\alpha) &= \frac{(a+b)^{-2}}{2^{\alpha-1}(\alpha+1)} {}_2F_1\left(2, 1; \alpha+2; \frac{b-a}{b+a}\right), \\
C_4(\alpha) &= \frac{(a+b)^{-2}}{2^{\alpha}(\alpha+2)} {}_2F_1\left(2, 1; \alpha+3; \frac{b-a}{b+a}\right), \\
C_5(\alpha) &= C_3(\alpha) - C_4(\alpha), \\
C_6(\alpha) &= \frac{b^{-2}}{2^{\alpha+1}(\alpha+1)} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{1}{2}\left(1-\frac{a}{b}\right)\right) \\
C_7(\alpha) &= \left[\begin{aligned} & \frac{b^{-2}}{2^{\alpha+1}(\alpha+1)} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{1}{2}\left(1-\frac{a}{b}\right)\right) \\ & - \frac{b^{-2}}{2^{\alpha+2}(\alpha+2)} {}_2F_1\left(2, \alpha+2; \alpha+3; \frac{1}{2}\left(1-\frac{a}{b}\right)\right) \end{aligned} \right] \\
C_8(\alpha) &= C_6(\alpha) - C_7(\alpha),
\end{aligned}$$

with $\alpha > 1$ and $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. Using (2.7), power mean inequality and the harmonically convexity of $|f'|^q$, it follows that

$$\begin{aligned}
& \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{2ab}{2ab}+}^{\alpha} (g \circ h) (1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (g \circ h) (1/b) \right] \right. \\
& \quad \left. - \left[J_{\frac{2ab}{2ab}+}^{\alpha} (fg \circ h) (1/a) + J_{\frac{2ab}{2ab}-}^{\alpha} (fg \circ h) (1/b) \right] \right| \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab} \right)^{\alpha} \left[\begin{aligned} & \int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \\ & + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \end{aligned} \right] \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab} \right)^{\alpha} \\
& \quad \left[\begin{aligned} & \left(\int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} du \right)^{1-\frac{1}{q}} \\ & \times \left(\int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right|^q du \right)^{\frac{1}{q}} \\ & + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} du \right)^{1-\frac{1}{q}} \\ & \times \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^2} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right|^q du \right)^{\frac{1}{q}} \end{aligned} \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{\|g\|_\infty ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^\alpha \\
&\quad \times \left[\begin{aligned} &\left(\int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} du\right)^{1-\frac{1}{q}} \\ &\times \left(\int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} [u|f'(a)|^q + (1-u)|f'(b)|^q] du\right)^{\frac{1}{q}} \\ &\quad + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} du\right)^{1-\frac{1}{q}} \\ &\times \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} [u|f'(a)|^q + (1-u)|f'(b)|^q] du\right)^{\frac{1}{q}} \end{aligned} \right] \\
&= \frac{\|g\|_\infty ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^\alpha \\
&\quad \times \left[\begin{aligned} &\left(\int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} du\right)^{1-\frac{1}{q}} \\ &\times \left(\begin{aligned} &\int_0^{\frac{1}{2}} \frac{u^{\alpha+1}}{(ub+(1-u)a)^2} du |f'(a)|^q \\ &+ \int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} (1-u) du |f'(b)|^q \end{aligned} \right)^{\frac{1}{q}} \\ &\quad + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} du\right)^{1-\frac{1}{q}} \\ &\times \left(\begin{aligned} &\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} u du |f'(a)|^q \\ &+ \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha+1}}{(ub+(1-u)a)^2} du |f'(b)|^q \end{aligned} \right)^{\frac{1}{q}} \end{aligned} \right] \tag{2.13}
\end{aligned}$$

Calculating following integrals, we have

$$\begin{aligned}
\int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} du &= \frac{1}{2^{\alpha+1}} \int_0^1 \frac{u^\alpha}{\left(\frac{u}{2}b + \left(1-\frac{u}{2}\right)a\right)^2} du \\
&= \frac{1}{2^{\alpha+1}} \int_0^1 (1-v)^\alpha \left(\frac{a+b}{2}\right)^{-2} \left(1-v\left(\frac{b-a}{b+a}\right)\right)^{-2} du \\
&= \frac{(a+b)^{-2}}{2^{\alpha-1}(\alpha+1)} {}_2F_1\left(2, 1; \alpha+2; \frac{b-a}{b+a}\right) \\
&= C_3(\alpha) \tag{2.14}
\end{aligned}$$

$$\begin{aligned}
\int_0^{\frac{1}{2}} \frac{u^{\alpha+1}}{(ub+(1-u)a)^2} du &= \frac{1}{2^{\alpha+2}} \int_0^1 \frac{u^{\alpha+1}}{\left(\frac{u}{2}b + \left(1-\frac{u}{2}\right)a\right)^2} du \\
&= \frac{1}{2^{\alpha+2}} \int_0^1 (1-v)^{\alpha+1} \left(\frac{a+b}{2}\right)^{-2} \left(1-v\left(\frac{b-a}{b+a}\right)\right)^{-2} du \\
&= \frac{(a+b)^{-2}}{2^\alpha(\alpha+2)} {}_2F_1\left(2, 1; \alpha+3; \frac{b-a}{b+a}\right) \\
&= C_4(\alpha) \tag{2.15}
\end{aligned}$$

$$\int_0^{\frac{1}{2}} \frac{u^\alpha}{(ub+(1-u)a)^2} (1-u) = C_3(\alpha) - C_4(\alpha) = C_5(\alpha) \tag{2.16}$$

$$\begin{aligned}
\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} du &= \int_0^{\frac{1}{2}} \frac{u^\alpha}{(ua+(1-u)b)^2} du \\
&= \frac{1}{2^{\alpha+1}} \int_0^1 \frac{u^\alpha}{\left(\frac{u}{2}a + \left(1-\frac{u}{2}\right)b\right)^2} du \\
&= \frac{1}{2^{\alpha+1}} \int_0^1 u^\alpha b^{-2} \left(1 - \frac{u}{2} \left(1 - \frac{a}{b}\right)\right)^{-2} du \\
&= \frac{b^{-2}}{2^{\alpha+1}(\alpha+1)} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{1}{2}\left(1 - \frac{a}{b}\right)\right) \\
&= C_6(\alpha) \tag{2.17}
\end{aligned}$$

$$\begin{aligned}
\int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} u du &= \int_0^{\frac{1}{2}} \frac{u^\alpha(1-u)}{(ua+(1-u)b)^2} du \\
&= \int_0^{\frac{1}{2}} \frac{u^\alpha}{(ua+(1-u)b)^2} du - \int_0^{\frac{1}{2}} \frac{u^{\alpha+1}}{(ua+(1-u)b)^2} du \\
&= \frac{1}{2^{\alpha+1}} \int_0^1 \frac{u^\alpha}{\left(\frac{u}{2}a + \left(1-\frac{u}{2}\right)b\right)^2} du \\
&\quad - \frac{1}{2^{\alpha+2}} \int_0^1 \frac{u^{\alpha+1}}{\left(\frac{u}{2}a + \left(1-\frac{u}{2}\right)b\right)^2} du \\
&= \frac{1}{2^{\alpha+1}} \int_0^1 u^\alpha b^{-2} \left(1 - \frac{u}{2} \left(1 - \frac{a}{b}\right)\right)^{-2} du \\
&\quad - \frac{1}{2^{\alpha+2}} \int_0^1 u^{\alpha+1} b^{-2} \left(1 - \frac{u}{2} \left(1 - \frac{a}{b}\right)\right)^{-2} du \\
&\leq \left[\begin{array}{l} \frac{b^{-2}}{2^{\alpha+1}(\alpha+1)} {}_2F_1\left(2, \alpha+1; \alpha+2; \frac{1}{2}\left(1 - \frac{a}{b}\right)\right) \\ -\frac{b^{-2}}{2^{\alpha+2}(\alpha+2)} {}_2F_1\left(2, \alpha+2; \alpha+3; \frac{1}{2}\left(1 - \frac{a}{b}\right)\right) \end{array} \right] \\
&= C_7(\alpha) \tag{2.18}
\end{aligned}$$

$$\begin{aligned}
\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha+1}}{(ub+(1-u)a)^2} du &= \int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} du - \int_{\frac{1}{2}}^1 \frac{(1-u)^\alpha}{(ub+(1-u)a)^2} u du \\
&= C_6(\alpha) - C_7(\alpha) = C_8(\alpha) \tag{2.19}
\end{aligned}$$

If we use (2.14 – 2.19) in (2.13), we have (2.12). This completes the proof. \square

Corollary 2. *In Theorem 7;*

(1) *If we take $\alpha = 1$ we have the following Hermite-Hadamard-Fejer inequality for harmonically convex functions which is related the left-hand side of (1.5):*

$$\begin{aligned}
&\left| f\left(\frac{2ab}{a+b}\right) \int_a^b \frac{g(x)}{x^2} dx - \int_a^b \frac{f(x)g(x)}{x^2} dx \right| \\
&\leq \|g\|_\infty (b-a)^2 \left[\begin{array}{l} C_3^{1-\frac{1}{q}}(1) \left[\left(\begin{array}{l} C_4(1) |f'(a)|^q \\ +C_5(1) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \\ +C_6^{1-\frac{1}{q}}(1) \left[\left(\begin{array}{l} C_7(1) |f'(a)|^q \\ +C_8(1) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \end{array} \right],
\end{aligned}$$

(2) *If we take $g(x) = 1$ we have following Hermite-Hadamard type inequality for harmonically convex function in fractional integral forms which is related the*

left-hand side of (2.1):

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) - \frac{\Gamma(\alpha+1)}{2^{1-\alpha}} \left(\frac{ab}{b-a}\right)^\alpha \left\{ \begin{array}{l} J_{\frac{a+b}{2a^+}}^\alpha (f \circ h)(1/a) \\ + J_{\frac{a+b}{2a^-}}^\alpha (f \circ h)(1/b) \end{array} \right\} \right| \\ & \leq \frac{ab(b-a)}{2^{1-\alpha}} \left[\begin{array}{l} C_3^{1-\frac{1}{q}}(\alpha) \left[\left(\begin{array}{l} C_4(\alpha) |f'(a)|^q \\ + C_5(\alpha) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \\ + C_6^{1-\frac{1}{q}}(\alpha) \left[\left(\begin{array}{l} C_7(\alpha) |f'(a)|^q \\ + C_8(\alpha) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \end{array} \right], \end{aligned}$$

(3) If we take $\alpha = 1$ and $g(x) = 1$ we have the following Hermite-Hadamard type inequality for harmonically convex function which is related the left-hand side of (1.4):>

$$\begin{aligned} & \left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\ & \leq ab(b-a) \left[\begin{array}{l} C_3^{1-\frac{1}{q}}(1) \left[\left(\begin{array}{l} C_4(1) |f'(a)|^q \\ + C_5(1) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \\ + C_6^{1-\frac{1}{q}}(1) \left[\left(\begin{array}{l} C_7(1) |f'(a)|^q \\ + C_8(1) |f'(b)|^q \end{array} \right) \right]^{\frac{1}{q}} \end{array} \right]. \end{aligned}$$

We can state another inequality for $q > 1$ as follows:

Theorem 8. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ and $a < b$. If $|f'|^q, q > 1$, is harmonically convex on $[a, b]$, $g : [a, b] \rightarrow \mathbb{R}$ is continuous and harmonically symmetric with respect to $\frac{2ab}{a+b}$, then the following inequality for fractional integrals hold:

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} \left[J_{1/b^+}^\alpha (g \circ h)(1/a) + J_{1/a^-}^\alpha (g \circ h)(1/b) \right] \right. \\ & \quad \left. - \left[J_{1/b^+}^\alpha (fg \circ h)(1/a) + J_{1/a^-}^\alpha (fg \circ h)(1/b) \right] \right| \\ & \leq \frac{\|g\|_\infty ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^\alpha \\ & \quad \times \left[\begin{array}{l} C_9^{\frac{1}{p}}(\alpha) \left[\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right]^{\frac{1}{q}} \\ + C_{10}^{\frac{1}{p}}(\alpha) \left[\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right]^{\frac{1}{q}} \end{array} \right] \end{aligned} \quad (2.20)$$

where

$$\begin{aligned} C_9(\alpha) &= \frac{(a+b)^{-2p}}{2^{\alpha p - 2p + 1} (\alpha p + 1)} {}_2F_1\left(2p, 1; \alpha p + 2; \frac{b-a}{b+a}\right), \\ C_{10}(\alpha) &= \frac{b^{-2p}}{2^{\alpha p + 1} (\alpha p + 1)} {}_2F_1\left(2, \alpha p + 1; \alpha p + 2; \frac{1}{2}\left(1 - \frac{a}{b}\right)\right), \end{aligned}$$

with $\alpha > 1$, $h(x) = 1/x$, $x \in [\frac{1}{b}, \frac{1}{a}]$ and $1/p + 1/q = 1$.

Proof. Using (2.7), Hölder's inequality and the convexity of $|f'|^q$, it follows that

$$\begin{aligned}
& \left| f\left(\frac{2ab}{a+b}\right) \left[J_{\frac{2ab}{a+b}+}^{\alpha} (g \circ h)(1/a) + J_{\frac{2ab}{a+b}-}^{\alpha} (g \circ h)(1/b) \right] \right. \\
& \quad \left. - \left[J_{\frac{2ab}{a+b}+}^{\alpha} (fg \circ h)(1/a) + J_{\frac{2ab}{a+b}-}^{\alpha} (fg \circ h)(1/b) \right] \right| \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \left[\int_0^{\frac{1}{2}} \frac{u^{\alpha}}{(ub+(1-u)a)^{2p}} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \right. \\
& \quad \left. + \int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha}}{(ub+(1-u)a)^{2p}} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right| du \right] \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \left[\left(\int_0^{\frac{1}{2}} \frac{u^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\
& \quad \left. \times \left(\int_0^{\frac{1}{2}} \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right|^q du \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\
& \quad \left. \times \left(\int_{\frac{1}{2}}^1 \left| f'\left(\frac{ab}{ub+(1-u)a}\right) \right|^q du \right)^{\frac{1}{q}} \right] \\
& \leq \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \left[\left(\int_0^{\frac{1}{2}} \frac{u^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\
& \quad \left. \times \left(\int_0^{\frac{1}{2}} u |f'(a)|^q + (1-u) |f'(b)|^q du \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\
& \quad \left. \times \left(\int_{\frac{1}{2}}^1 u |f'(a)|^q + (1-u) |f'(b)|^q du \right)^{\frac{1}{q}} \right] \\
& = \frac{\|g\|_{\infty} ab(b-a)}{\Gamma(\alpha+1)} \left(\frac{b-a}{ab}\right)^{\alpha} \\
& \quad \times \left[\left(\int_0^{\frac{1}{2}} \frac{u^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \left[\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right]^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha p}}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \left[\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right]^{\frac{1}{q}} \right] \tag{2.21}
\end{aligned}$$

Calculating following integrals, we have

$$\begin{aligned}
\int_0^{\frac{1}{2}} \frac{u^{\alpha p}}{(ub+(1-u)a)^{2p}} du &= \frac{1}{2^{\alpha p+1}} \int_0^1 \frac{u^{\alpha p}}{\left(\frac{u}{2}b + \left(1-\frac{u}{2}\right)a\right)^{2p}} du \\
&= \frac{1}{2^{\alpha p+1}} \int_0^1 (1-v)^{\alpha p} \left(\frac{a+b}{2}\right)^{-2p} \left[1-v\left(\frac{b-a}{b+a}\right)\right]^{-2p} dv \\
&= \frac{(a+b)^{-2p}}{2^{\alpha p-2p+1}(\alpha p+1)} {}_2F_1\left(2p, 1; \alpha p+2; \frac{b-a}{b+a}\right) \\
&= C_9(\alpha) \tag{2.22}
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\int_{\frac{1}{2}}^1 \frac{(1-u)^{\alpha p}}{(ub+(1-u)a)^{2p}} du &= \int_0^{\frac{1}{2}} \frac{u^{\alpha p}}{(ua+(1-u)b)^{2p}} du \\
&= \frac{1}{2^{\alpha p+1}} \int_0^1 \frac{u^{\alpha p}}{\left(\frac{u}{2}a + \left(1-\frac{u}{2}\right)b\right)^{2p}} du \\
&= \frac{1}{2^{\alpha p+1}} \int_0^1 u^{\alpha} b^{-2p} \left(1 - \frac{u}{2} \left(1 - \frac{a}{b}\right)\right)^{-2p} du \\
&= \frac{b^{-2p}}{2^{\alpha p+1}(\alpha p+1)} {}_2F_1\left(2, \alpha p+1; \alpha p+2; \frac{1}{2}\left(1 - \frac{a}{b}\right)\right) \\
&= C_{10}(\alpha) \tag{2.23}
\end{aligned}$$

If we use (2.22) and (2.23) in (2.21), we have (2.20). This completes the proof. \square

Corollary 3. *In Theorem 8;*

(1) *If we take $\alpha = 1$ we have the following Hermite-Hadamard-Fejer inequality for harmonically convex functions which is related the left-hand side of (1.5):*

$$\begin{aligned}
&\left| f\left(\frac{2ab}{a+b}\right) \int_a^b \frac{g(x)}{x^2} dx - \int_a^b \frac{f(x)g(x)}{x^2} dx \right| \\
&\leq \|g\|_{\infty} (b-a)^2 \left[C_9^{\frac{1}{p}}(1) \left[\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right]^{\frac{1}{q}} + C_{10}^{\frac{1}{p}}(1) \left[\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right]^{\frac{1}{q}} \right],
\end{aligned}$$

(2) *If we take $g(x) = 1$ we have following Hermite-Hadamard type inequality for harmonically convex function in fractional integral forms which is related the left-hand side of (2.1):*

$$\begin{aligned}
&\left| f\left(\frac{2ab}{a+b}\right) - \frac{\Gamma(\alpha+1)}{2^{1-\alpha}} \left(\frac{ab}{b-a}\right)^{\alpha} \left\{ \begin{array}{l} J_{\frac{a+b}{2ab}^+}^{\alpha} (f \circ h)(1/a) \\ + J_{\frac{a+b}{2ab}^-}^{\alpha} (f \circ h)(1/b) \end{array} \right\} \right| \\
&\leq \frac{ab(b-a)}{2^{1-\alpha}} \left[C_9^{\frac{1}{p}}(\alpha) \left[\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right]^{\frac{1}{q}} + C_{10}^{\frac{1}{p}}(\alpha) \left[\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right]^{\frac{1}{q}} \right],
\end{aligned}$$

(3) *If we take $\alpha = 1$ and $g(x) = 1$ we have the following Hermite-Hadamard type inequality for harmonically convex function which is related the left-hand side of (1.4):*

$$\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \leq ab(b-a) \left[C_9^{\frac{1}{p}}(1) \left[\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right]^{\frac{1}{q}} + C_{10}^{\frac{1}{p}}(1) \left[\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right]^{\frac{1}{q}} \right].$$

REFERENCES

- [1] M. Bombardelli and S. Varošanec, Properties of h -convex functions related to the Hermite Hadamard Fejer inequalities, *Computers and Mathematics with Applications* 58 (2009), 1869-1877.
- [2] F. Chen and S. Wu, Fejer and Hermite-Hadamard type inequalities for harmonically convex functions, *Journal of applied Mathematics*, volume 2014, article id:386806.
- [3] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.* 1(1) (2010), 51-58.

- [4] L. Fejér, Über die Fourierreihen, II, Math. Naturwiss. Anz Ungar. Akad., Wiss, 24 (1906), 369-390, (in Hungarian).
- [5] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, J. Math. Pures Appl., 58 (1893), 171-215.
- [6] İ. İşcan, New estimates on generalization of some integral inequalities for s -convex functions and their applications, Int. J. Pure Appl. Math., 86(4) (2013), 727-746.
- [7] İ. İşcan, Some new general integral inequalities for h -convex and h -concave functions, Adv. Pure Appl. Math. 5(1) (2014), 21-29 . doi: 10.1515/apam-2013-0029.
- [8] İ. İşcan, Generalization of different type integral inequalities for s -convex functions via fractional integrals, Applicable Analysis, 2013. doi: 10.1080/00036811.2013.851785.
- [9] İ. İşcan, On generalization of different type integral inequalities for s -convex functions via fractional integrals, Mathematical Sciences and Applications E-Notes, 2(1) (2014), 55-67.
- [10] İ. İşcan, S. Wu, Hermite-Hadamard type inequalities for harmonically convex functions via fractional integrals, Appl. Math. Comput., 238 (2014) 237-244.
- [11] İ. İşcan, Hermite-Hadamard type inequalities for harmonically convex functions, Hacet. J. Math. Stat., 43 (6) (2014), 935-942
- [12] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and applications of fractional differential equations. Elsevier, Amsterdam (2006).
- [13] M. A. Latif, S. S. Dragomir and E. Momoniat, Some Fejer type inequalities for harmonically-convex functions with applications to special means, <http://rgmia.org/papers/v18/v18a24.pdf>.
- [14] A. P. Prudnikov, Y. A. Brychkov, O. J. Marichev, Integral and series, Elementary Functions, vol. 1, Nauka, Moscow, 1981.
- [15] M.Z. Sarıkaya, On new Hermite Hadamard Fejér type integral inequalities, Stud. Univ. Babeş-Bolyai Math. 57(3) (2012), 377–386.
- [16] M.Z. Sarıkaya, E. Set, H. Yıldız and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Mathematical and Computer Modelling, 57(9) (2013), 2403-2407.
- [17] K.-L. Tseng, G.-S. Yang and K.-C. Hsu, Some inequalities for differentiable mappings and applications to Fejér inequality and weighted trapezoidal formula, Taiwanese journal of Mathematics, 15(4) (2011), 1737-1747.
- [18] J. Wang, X. Li, M. Fečkan and Y. Zhou, Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity, Appl. Anal., 92(11) (2012), 2241-2253. doi:10.1080/00036811.2012.727986
- [19] J. Wang, C. Zhu and Y. Zhou, New generalized Hermite-Hadamard type inequalities and applications to special means, J. Inequal. Appl., 2013(325) (2013), 15 pages.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, KARADENİZ TECHNICAL UNIVERSITY, TRABZON, TURKEY

E-mail address: mkunt@ktu.edu.tr

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND ARTS, GİRESUN UNIVERSITY, GİRESUN, TURKEY

E-mail address: imdat.iscan@giresun.edu.tr; imdati@yahoo.com