

**SOME HERMITE-HADAMARD TYPE INEQUALITIES FOR
FUNCTIONS WHOSE PRODUCT WITH EXPONENTIALS ARE
CONVEX**

S.S. DRAGOMIR^{1,2} AND I. GOMM¹

ABSTRACT. Some inequalities of Hermite-Hadamard type for functions whose product with exponentials are convex, are obtained.

1. INTRODUCTION

The following integral inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a) + f(b)}{2},$$

which holds for any convex function $f : [a, b] \rightarrow \mathbb{R}$, is well known in the literature as the Hermite-Hadamard inequality.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the Theory of Special Means and in Information Theory for divergence measures, from which we would like to refer the reader to the papers [1] – [59] and the references therein.

Following [20], we denote by $\mathfrak{E}xpconv(I)$ the class of all functions defined on the interval I of real numbers such that $\exp(f)$ is convex on I . If $\mathfrak{C}onv(I)$ is the class of convex functions defined on I then we have the following fact [20]:

Proposition 1. *We have the strict inclusion*

$$\mathfrak{C}onv(I) \subsetneq \mathfrak{E}xpconv(I).$$

Now, if $g \in \mathfrak{E}xpconv(I)$, then by the Hermite-Hadamard inequality for $\exp(g)$ we have for $a, b \in I$ with $a < b$ that

$$(1.2) \quad \exp g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b \exp g(t) dt \leq \frac{1}{2} [\exp g(a) + \exp g(b)].$$

Consider the *identric mean* of two positive numbers

$$I(a, b) := \begin{cases} a & \text{if } a = b, \\ \frac{1}{e} \left(\frac{b^b}{a^a}\right)^{\frac{1}{b-a}} & \text{if } a \neq b, \end{cases} \quad a, b > 0.$$

We observe that

$$\ln I(a, b) = \frac{1}{b-a} \int_a^b \ln u du$$

for $a, b > 0$, $a \neq b$.

1991 *Mathematics Subject Classification.* 26D15; 25D10.

Key words and phrases. Convex functions, Hermite-Hadamard inequality, Special means.

The following result holds [20]:

Theorem 1. *Assume that $f \in \mathfrak{Expconv}(I)$ and $a, b \in I$ with $a < b$. Then we have*

$$(1.3) \quad \exp\left(\frac{1}{b-a} \int_a^b f(t) dt\right) \leq I(\exp f(a), \exp f(b))$$

and

$$(1.4) \quad \begin{aligned} & \exp f\left(\frac{a+b}{2}\right) \\ & \leq \exp\left(\frac{1}{b-a} \int_a^b \ln\left[\frac{\exp f(x) + \exp f(a+b-x)}{2}\right] dx\right) \\ & \leq \frac{1}{b-a} \int_a^b \exp f(x) dx. \end{aligned}$$

Remark 1. *If the function $g : I \rightarrow (0, \infty)$ is convex on I , then $f = \ln g \in \mathfrak{Expconv}(I)$ and for $a, b \in I$ with $a < b$ we have, by (1.3) and (1.4), the following inequalities*

$$(1.5) \quad \exp\left(\frac{1}{b-a} \int_a^b \ln g(t) dt\right) \leq I(g(a), g(b))$$

and

$$(1.6) \quad \begin{aligned} g\left(\frac{a+b}{2}\right) & \leq \exp\left(\frac{1}{b-a} \int_a^b \ln\left[\frac{g(x) + g(a+b-x)}{2}\right] dx\right) \\ & \leq \frac{1}{b-a} \int_a^b g(x) dx. \end{aligned}$$

We denote by $\mathfrak{Expconv}_\alpha(I)$ for $\alpha \neq 0$, the class of all functions f defined on the interval I of real numbers such that the function $g_\alpha(t) := f(t) \exp(\alpha t)$ is convex on I .

Consider the function $\ell(t) = t$, $t \in \mathbb{R}$. Then $g_\alpha(t) = t \exp(\alpha t)$. Taking the second derivative of g_α we have

$$g_\alpha''(t) = \alpha e^{\alpha t} (\alpha t + 2).$$

We observe that for either $\alpha > 0$ or $\alpha < 0$ the function g_α is concave on $(\infty, -\frac{2}{\alpha}]$ and convex on $[-\frac{2}{\alpha}, \infty)$. Therefore for any $\alpha \neq 0$ the function $\ell \in \mathfrak{Expconv}_\alpha\left([-\frac{2}{\alpha}, \infty)\right)$.

The interested reader can give other examples of such functions, for instance the family of functions $f_p(t) = t^p$, $p < 0$ and $t \in (0, \infty)$ are in the class $\mathfrak{Expconv}_\alpha(0, \infty)$ for $\alpha \neq 0$. Indeed, if we take the second derivative of the function $g_{\alpha,p}(t) = t^p \exp(\alpha t)$, then we have

$$g_{\alpha,p}''(t) = e^{\alpha t} t^{p-2} \left((\alpha t + p)^2 - p \right) > 0,$$

for any $\alpha \neq 0$, $t \in (0, \infty)$ and $p < 0$.

Motivated by the above results, we establish in this paper some inequalities of Hermite-Hadamard type for functions from the class $\mathfrak{Expconv}_\alpha(I)$.

2. SOME INEQUALITIES

We have the following result:

Theorem 2. *Assume that the function $f : I \rightarrow \mathbb{R}$ has the property that $f \in \mathfrak{Epproconv}_\alpha(I)$ for some $\alpha \neq 0$. Then f is integrable on any interval $[a, b] \subset I$ and*

$$(2.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) \exp\left[\alpha\left(t - \frac{a+b}{2}\right)\right] dt \\ \leq \frac{1}{2} \left[f(a) \exp\left(-\alpha \frac{b-a}{2}\right) + f(b) \exp\left(\alpha \frac{b-a}{2}\right) \right].$$

Proof. Since $f \in \mathfrak{Epproconv}_\alpha(I)$, $\alpha \neq 0$ then there exists a convex function $g_\alpha : I \rightarrow \mathbb{R}$ such that $f(t) \exp(\alpha t) = g_\alpha(t)$, which gives that $f(t) = g_\alpha(t) \exp(-\alpha t)$, $t \in I$. This shows that f is integrable on $[a, b]$.

Now, using the Hermite-Hadamard inequality for g_α we have

$$f\left(\frac{a+b}{2}\right) \exp\left(\alpha \frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) \exp(\alpha t) dt \\ \leq \frac{1}{2} [f(a) \exp(\alpha a) + f(b) \exp(\alpha b)].$$

Dividing this inequality by $\exp\left(\alpha \frac{a+b}{2}\right)$ we get the desired result (2.1).

An alternative direct proof for the first inequality is as follows.

By the fact that $g_\alpha(t) = f(t) \exp(\alpha t)$ is convex on I , we have

$$f\left(\frac{x+y}{2}\right) \exp\left[\alpha\left(\frac{x+y}{2}\right)\right] \leq \frac{1}{2} [f(x) \exp(\alpha x) + f(y) \exp(\alpha y)],$$

which is equivalent, by division with $\exp\left[\alpha\left(\frac{x+y}{2}\right)\right]$, to

$$(2.2) \quad f\left(\frac{x+y}{2}\right) \leq \frac{1}{2} \left[f(x) \exp\left(-\frac{1}{2}\alpha(y-x)\right) + f(y) \exp\left(\frac{1}{2}\alpha(y-x)\right) \right]$$

for any $x, y \in I$.

If we take

$$y = (1-t)a + tb \text{ and } x = (1-t)b + ta$$

then we have

$$y - x = (2t - 1)(b - a) \text{ and } \frac{x+y}{2} = \frac{a+b}{2},$$

and by (2.2) we get

$$(2.3) \quad f\left(\frac{a+b}{2}\right) \\ \leq \frac{1}{2} f((1-t)b + ta) \exp\left(-\frac{1}{2}\alpha(2t-1)(b-a)\right) \\ + \frac{1}{2} f((1-t)a + tb) \exp\left(\frac{1}{2}\alpha(2t-1)(b-a)\right) \\ = \frac{1}{2} f((1-t)b + ta) \exp\left(-\alpha\left(t - \frac{1}{2}\right)(b-a)\right) \\ + \frac{1}{2} f((1-t)a + tb) \exp\left(\alpha\left(t - \frac{1}{2}\right)(b-a)\right)$$

for any $t \in [0, 1]$.

Integrate over $t \in [0, 1]$ to get

$$(2.4) \quad \begin{aligned} & f\left(\frac{a+b}{2}\right) \\ & \leq \frac{1}{2} \int_0^1 f((1-t)b+ta) \exp\left(-\alpha\left(t-\frac{1}{2}\right)(b-a)\right) dt \\ & \quad + \frac{1}{2} \int_0^1 f((1-t)a+tb) \exp\left(\alpha\left(t-\frac{1}{2}\right)(b-a)\right) dt. \end{aligned}$$

If we change the variable by putting $s = 1 - t$, then we have

$$\begin{aligned} & \int_0^1 f((1-t)a+tb) \exp\left(\alpha\left(t-\frac{1}{2}\right)(b-a)\right) dt \\ & = \int_0^1 f(sa+(1-s)b) \exp\left(-\alpha\left(s-\frac{1}{2}\right)(b-a)\right) ds \end{aligned}$$

and by (2.4) we get

$$(2.5) \quad f\left(\frac{a+b}{2}\right) \leq \int_0^1 f((1-t)b+ta) \exp\left(-\alpha\left(t-\frac{1}{2}\right)(b-a)\right) dt.$$

Now, if we change the variable $u = (1-t)b+ta$, then $t = \frac{b-u}{b-a}$,

$$t - \frac{1}{2} = \frac{b-u}{b-a} - \frac{1}{2} = \frac{2b-2u-b+a}{2(b-a)} = \frac{1}{b-a} \left(\frac{b+a}{2} - u\right),$$

which implies that

$$\begin{aligned} \exp\left(-\alpha\left(t-\frac{1}{2}\right)(b-a)\right) & = \exp\left[\frac{-\alpha}{b-a} \left(\frac{b+a}{2} - u\right)(b-a)\right] \\ & = \exp\left[\alpha\left(u - \frac{b+a}{2}\right)\right]. \end{aligned}$$

Then by (2.5) we get

$$f\left(\frac{a+b}{2}\right) \leq \int_a^b f(u) \exp\left[\alpha\left(u - \frac{b+a}{2}\right)\right] du.$$

□

The following result also holds:

Theorem 3. *Assume that the function $f : I \rightarrow \mathbb{R}$ has the property that $f \in \mathfrak{Epproconv}_\alpha(I)$ for some $\alpha \neq 0$. Then for any $[a, b] \subset I$ with $a < b$ we have*

$$(2.6) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{\alpha(b-a)} \\ & \quad \times \left(\left[1 - \frac{\exp(-\alpha(b-a)) - 1}{\alpha(b-a)} \right] f(a) + \left[\frac{1 - \exp(\alpha(b-a))}{\alpha(b-a)} - 1 \right] f(b) \right). \end{aligned}$$

Proof. Since $f \in \mathfrak{Epproconv}_\alpha(I)$, $\alpha \neq 0$, then

$$\begin{aligned} & f((1-t)a+tb) \exp[\alpha((1-t)a+tb)] \\ & \leq (1-t)f(a) \exp(\alpha a) + tf(b) \exp(\alpha b) \end{aligned}$$

and if we divide by $\exp[\alpha((1-t)a+tb)]$ we get

$$\begin{aligned}
 (2.7) \quad & f((1-t)a+tb) \\
 & \leq (1-t)f(a)\exp(\alpha a - \alpha((1-t)a+tb)) \\
 & \quad + tf(b)\exp(\alpha b - \alpha((1-t)a+tb)) \\
 & = (1-t)f(a)\exp(\alpha t(a-b)) + tf(b)\exp(\alpha(1-t)(b-a))
 \end{aligned}$$

for any $a, b \in I$ and $t \in [0, 1]$.

Integrating (2.7) over t on $[0, 1]$ we get

$$\begin{aligned}
 (2.8) \quad & \int_0^1 f((1-t)a+tb) dt \\
 & \leq f(a) \int_0^1 (1-t)\exp(\alpha t(a-b)) dt + f(b) \int_0^1 t\exp(\alpha(1-t)(b-a)) dt.
 \end{aligned}$$

Observe, for $a \neq b$, that

$$\begin{aligned}
 & \int_0^1 (1-t)\exp(\alpha t(a-b)) dt \\
 & = \frac{1}{\alpha(a-b)} \int_0^1 (1-t) d[\exp(\alpha t(a-b))] \\
 & = \frac{1}{\alpha(a-b)} \left[(1-t)\exp(\alpha t(a-b)) \Big|_0^1 + \int_0^1 \exp(\alpha t(a-b)) dt \right] \\
 & = \frac{1}{\alpha(a-b)} \left[-1 + \frac{1}{\alpha(a-b)} \exp(\alpha t(a-b)) \Big|_0^1 \right] \\
 & = \frac{1}{\alpha(a-b)} \left[\frac{\exp(\alpha(a-b)) - 1}{\alpha(a-b)} - 1 \right]
 \end{aligned}$$

and

$$\begin{aligned}
 & \int_0^1 t\exp(\alpha(1-t)(b-a)) dt \\
 & = \int_0^1 t\exp(\alpha(t-1)(a-b)) dt \\
 & = \frac{1}{\alpha(a-b)} \int_0^1 td[\exp(\alpha(t-1)(a-b))] \\
 & = \frac{1}{\alpha(a-b)} \left[t[\exp(\alpha(t-1)(a-b))] \Big|_0^1 - \int_0^1 \exp(\alpha(t-1)(a-b)) dt \right] \\
 & = \frac{1}{\alpha(a-b)} \left[1 - \frac{1}{\alpha(a-b)} \exp(\alpha(t-1)(a-b)) \Big|_0^1 \right] \\
 & = \frac{1}{\alpha(a-b)} \left[1 - \frac{1 - \exp(-\alpha(a-b))}{\alpha(a-b)} \right]
 \end{aligned}$$

and by (2.8) we get the desired result (2.6). \square

Remark 2. Since the family of functions $f_p(t) = t^p$, $p < 0$ and $t \in (0, \infty)$ are in the class $\mathfrak{Epproconv}_\alpha(0, \infty)$ for $\alpha \neq 0$, then by (2.1) we have for $0 < a < b$ that

$$(2.9) \quad \begin{aligned} \left(\frac{a+b}{2}\right)^p &\leq \frac{1}{b-a} \int_a^b t^p \exp\left[\alpha\left(t - \frac{a+b}{2}\right)\right] dt \\ &\leq \frac{1}{2} \left[a^p \exp\left(-\alpha\frac{b-a}{2}\right) + b^p \exp\left(\alpha\frac{b-a}{2}\right) \right] \end{aligned}$$

and by (2.6) we have

$$(2.10) \quad \begin{aligned} &\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \\ &\leq \frac{1}{\alpha(b-a)} \\ &\times \left(\left[1 - \frac{\exp(-\alpha(b-a)) - 1}{\alpha(b-a)} \right] a^p + \left[\frac{1 - \exp(\alpha(b-a))}{\alpha(b-a)} - 1 \right] b^p \right), \end{aligned}$$

here $p \in (-\infty, 0) \setminus \{-1\}$.

3. FURTHER RESULTS

We also have the following result:

Theorem 4. Assume that the function $f : I \rightarrow \mathbb{R}$ has the property that $f \in \mathfrak{Epproconv}_\alpha(I)$ for some $\alpha \neq 0$. Then for any $[a, b] \subset I$ we have

$$(3.1) \quad \begin{aligned} &\frac{1}{(b-a)^2} \int_a^b \int_a^b f\left(\frac{x+y}{2}\right) dx dy \\ &\leq E\left(-\frac{1}{2}\alpha b, -\frac{1}{2}\alpha a\right) \frac{1}{b-a} \int_a^b f(y) \exp\left(\frac{1}{2}\alpha y\right) dy, \end{aligned}$$

where the exponential mean E is defined as

$$E(c, d) := \frac{\exp d - \exp c}{d - c}, \quad d, c \in \mathbb{R}, d \neq c.$$

Proof. Using the inequality (2.2) and integrating on the square $[a, b] \times [a, b]$ we have

$$(3.2) \quad \begin{aligned} &\frac{1}{(b-a)^2} \int_a^b \int_a^b f\left(\frac{x+y}{2}\right) dx dy \\ &\leq \frac{1}{2} \frac{1}{(b-a)^2} \int_a^b \int_a^b f(x) \exp\left(-\frac{1}{2}\alpha(y-x)\right) dx dy \\ &\quad + \frac{1}{2} \frac{1}{(b-a)^2} \int_a^b \int_a^b f(y) \exp\left(\frac{1}{2}\alpha(y-x)\right) dx dy. \end{aligned}$$

Changing x with y we get

$$\begin{aligned} &\frac{1}{(b-a)^2} \int_a^b \int_a^b f(x) \exp\left(-\frac{1}{2}\alpha(y-x)\right) dx dy \\ &= \frac{1}{(b-a)^2} \int_a^b \int_a^b f(y) \exp\left(\frac{1}{2}\alpha(y-x)\right) dx dy \end{aligned}$$

and by (3.2) we get

$$(3.3) \quad \begin{aligned} & \frac{1}{(b-a)^2} \int_a^b \int_a^b f\left(\frac{x+y}{2}\right) dx dy \\ & \leq \frac{1}{(b-a)^2} \int_a^b \int_a^b f(y) \exp\left(\frac{1}{2}\alpha(y-x)\right) dx dy. \end{aligned}$$

Since

$$\begin{aligned} & \frac{1}{(b-a)^2} \int_a^b \int_a^b f(y) \exp\left(\frac{1}{2}\alpha(y-x)\right) dx dy \\ & = \frac{1}{(b-a)^2} \int_a^b \exp\left(-\frac{1}{2}\alpha x\right) dx \int_a^b f(y) \exp\left(\frac{1}{2}\alpha y\right) dy \\ & = E\left(-\frac{1}{2}\alpha b, -\frac{1}{2}\alpha a\right) \frac{1}{b-a} \int_a^b f(y) \exp\left(\frac{1}{2}\alpha y\right) dy, \end{aligned}$$

then we get from (3.3) the desired result (3.1). \square

Finally we have:

Theorem 5. *Assume that the function $f : I \rightarrow \mathbb{R}$ has the property that $f \in \mathfrak{E}_{\text{pproconv}}_{\alpha}(I)$ for some $\alpha \neq 0$. Then for any $[a, b] \subset I$ with $a < b$ we have*

$$(3.4) \quad \begin{aligned} & f(x) \exp(\alpha x) E(-\alpha a, -\alpha b) - \alpha \frac{1}{b-a} \int_a^b f(y) (x-y) dy \\ & \geq \frac{2}{b-a} \int_a^b f(y) dy - \frac{f(b)(b-x) + f(a)(x-a)}{b-a} \end{aligned}$$

for any $x \in [a, b]$.

In particular, we have

$$(3.5) \quad \begin{aligned} & f\left(\frac{a+b}{2}\right) \exp\left(\alpha \frac{a+b}{2}\right) E(-\alpha a, -\alpha b) \\ & - \alpha \frac{1}{b-a} \int_a^b f(y) \left(\frac{a+b}{2} - y\right) dy \\ & \geq \frac{2}{b-a} \int_a^b f(y) dy - \frac{f(b) + f(a)}{2}. \end{aligned}$$

Proof. Since $f \exp(\alpha \cdot)$ is convex then $f \exp(\alpha \cdot)$ has lateral derivative on the interior \mathring{I} of I and therefore f has lateral derivatives on \mathring{I} .

Using the gradient inequality

$$g(x) - g(y) \geq g'_+(y) (x - y)$$

for any $x, y \in \mathring{I}$, that holds for a convex function g defined on the interval I we have

$$f(x) \exp(\alpha x) - f(y) \exp(\alpha y) \geq [\alpha f(y) \exp(\alpha y) + f'_+(y) \exp(\alpha y)] (x - y)$$

for any $x, y \in \mathring{I}$.

Dividing by $\exp(\alpha y)$ we get

$$f(x) \exp(\alpha x) \exp(-\alpha y) - f(y) \geq [\alpha f(y) + f'_+(y)] (x - y)$$

that is equivalent to

$$(3.6) \quad f(x) \exp(\alpha x) \exp(-\alpha y) - \alpha f(y)(x-y) \geq f(y) + f'_+(y)(x-y)$$

for any $x, y \in \hat{I}$.

Integrating (3.6) over $y \in [a, b]$ and dividing by $b-a$ we get

$$(3.7) \quad f(x) \exp(\alpha x) \frac{1}{b-a} \int_a^b \exp(-\alpha y) dy - \alpha \frac{1}{b-a} \int_a^b f(y)(x-y) dy \\ \geq \frac{1}{b-a} \int_a^b f(y) dy + \frac{1}{b-a} \int_a^b f'_+(y)(x-y) dy.$$

Since

$$\frac{1}{b-a} \int_a^b \exp(-\alpha y) dy = E(-\alpha a, -\alpha b)$$

and

$$\int_a^b f'_+(y)(x-y) dy = f(y)(x-y)|_a^b + \int_a^b f(y) dy \\ = \int_a^b f(y) dy - f(b)(b-x) - f(a)(x-a),$$

then by (3.7) we get the desired result (3.4). \square

Corollary 1. *With the assumptions of Theorem 5 and if*

$$(3.8) \quad \frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} \in [a, b],$$

then

$$(3.9) \quad f \left(\frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} \right) \exp \left(\alpha \frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} \right) E(-\alpha a, -\alpha b) \\ \geq \frac{2}{b-a} \int_a^b f(y) dy \\ - \frac{1}{b-a} \left[f(b) \left(b - \frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} \right) + f(a) \left(\frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} - a \right) \right].$$

Remark 3. *We observe that a sufficient condition for (3.8) to hold is that $f(y) \geq 0$ for any $y \in [a, b]$.*

REFERENCES

- [1] Allasia, G., Giordano, C. & Pečarić, J. (1999). Hadamard-type Inequalities for $(2r)$ -convex Functions with Applications, *Atti Acad. Sci. Torino-Cl. Sc. Fis.*, 133, 1-14.
- [2] Alzer, H. (1989). A Note on Hadamard's Inequalities, *C.R. Math. Rep. Acad. Sci. Canada*, 11, 255-258.
- [3] Alzer, H. (1989). On an Integral Inequality, *Math. Rev. Anal. Numer. Theor. Approx.*, 18, 101-103.
- [4] Azpeitia, A.G. (1994). Convex Functions and the Hadamard Inequality, *Rev.-Colombiana-Mat.*, 28(1), 7-12.
- [5] Barbu, D., Dragomir, S.S. & Buşe, C. (1993). A Probabilistic Argument for the Convergence of Some Sequences Associated to Hadamard's Inequality, *Studia Univ. Babeş-Bolyai, Math.*, 38 (1), 29-33.

- [6] Buge, C., Dragomir, S.S. & Barbu, D. (1996). The Convergence of Some Sequences Connected to Hadamard's Inequality, *Demonstratio Math.*, 29(1) (1996), 53-59.
- [7] Dragomir, S.S. (1991). A Mapping in Connection to Hadamard's Inequalities, *An. Öster. Akad. Wiss. Math.-Natur.*, (Wien), 128, 17-20.
- [8] Dragomir, S.S. (1993). A Refinement of Hadamard's Inequality for Isotonic Linear Functionals, *Tamkang J. of Math.* (Taiwan), 24, 101-106.
- [9] Dragomir, S.S. (1992). On Hadamard's Inequalities for Convex Functions, *Mat. Balkanica*, 6, 215-222.
- [10] Dragomir, S.S. (2000). On Hadamard's Inequality for the Convex Mappings Defined on a Ball in the Space and Applications, *Math. Ineq. & Appl.*, 3(2) , 177-187.
- [11] Dragomir, S.S. (2000). On Hadamard's Inequality on a Disk, *Journal of Ineq. in Pure & Appl. Math.*, 1(1), Article 2, 1-11.
- [12] Dragomir, S.S. (1992). Some Integral Inequalities for Differentiable Convex Functions, *Contributions, Macedonian Acad. of Sci. and Arts*, 13(1) , 13-17.
- [13] Dragomir, S.S. (1992). Two mappings in Connection to Hadamard's Inequalities, *J. Math. Anal. Appl.*, **167**, 49-56.
- [14] Dragomir, S.S. (1994). Some Remarks on Hadamard's Inequalities for Convex Functions, *Extracta Math.*, 9 (2) , 88-94.
- [15] Dragomir, S.S. & Agarwal, R.P. (1998). Two New Mappings Associated with Hadamard's Inequalities for Convex Functions, *Appl. Math. Lett.*, 11 (3), 33-38.
- [16] Dragomir, S.S. & Buşe, C. (1995). Refinements of Hadamard's Inequality for Multiple Integrals, *Utilitas Math.* (Canada), 47, 193-195.
- [17] Dragomir, S.S., Cho, Y.J. & Kim, S.S. (2000). Inequalities of Hadamard's Type for Lipschitzian Mappings and their Applications, *J. of Math. Anal. Appl.*, 245 (2) , 489-501.
- [18] Dragomir, S.S. & Gomm, I. (2011). Bounds for Two Mappings Associated to the Hermite-Hadamard Inequality, *Aust. J. Math. Anal. Appl.*, 8, Art. 5,1-9.
- [19] Dragomir, S.S. & Gomm, I. (2012). Some New Bounds for Two Mappings Related to the Hermite-Hadamard Inequality for Convex Functions, *Num. Alg. Cont. & Opt.* 2(2), 271-278.
- [20] Dragomir, S.S. & Gomm, I. (2015). Some Hermite-Hadamard Type Inequalities for Functions Whose Exponentials are Convex, Preprint, *RGMIA Res. Rep. Coll.* **18** (2015), Art.
- [21] Dragomir, S.S. & Fitzpatrick, S. (1998). The Hadamard's Inequality for s -convex Functions in the First Sense, *Demonstratio Math.*, 31(3) , 633-642.
- [22] Dragomir, S.S. & Fitzpatrick, S. (1999). The Hadamard's Inequality for s -convex Functions in the Second Sense, *Demonstratio Math.*, 32 (4) , 687-696.
- [23] Dragomir, S.S. & Ionescu, N.M. (1990). On Some Inequalities for Convex-dominated Functions, *Anal. Num. Theor. Approx.*, 19, 21-28.
- [24] Dragomir, S.S., Milošević, D.S. & Sándor, J. (1993). On Some Refinements of Hadamard's Inequalities and Applications, *Univ. Belgrad, Publ. Elek. Fak. Sci. Math.*, 4, 21-24.
- [25] Dragomir, S.S. & Mond, B. (1997). On Hadamard's Inequality for a Class of Functions of Godunova and Levin, *Indian J. Math.*, 39(1), 1-9.
- [26] Dragomir, S.S. & Pearce, C.E.M. (1998). Quasi-convex Functions and Hadamard's Inequality, *Bull. Austral. Math. Soc.*, 57, 377-385.
- [27] Dragomir, S.S. & Pearce, C.E.M. (2000). *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, [Online http://rgmia.org/monographs/hermite_hadamard.html].
- [28] Dragomir, S.S., Pearce, C.E.M. & Pečarić, J. (1995). On Jessen's and Related Inequalities for Isotonic Sublinear Functionals, *Acta Math. Sci.* (Szeged), 61, 373-382.
- [29] Dragomir, S.S., Pečarić, J. & Persson, L.E. (1995). Some Inequalities of Hadamard type, *Soochow J. of Math.* (Taiwan), 21, 335-341.
- [30] Dragomir, S.S., Pečarić, J. & Sándor, J. (1990). A Note on the Jensen-Hadamard Inequality, *Anal. Num. Theor. Approx.*, 19, 21-28.
- [31] Dragomir, S.S. & Toader, G.H. (1993). Some Inequalities for m -convex Functions, *Studia Univ. Babeş-Bolyai, Math.*, 38(1) , 21-28.
- [32] Fink, A.M. (1998). A Best Possible Hadamard Inequality, *Math. Ineq. & Appl.*, 2, 223-230.
- [33] Fink, A.M. (1994). Toward a Theory of Best Possible Inequalities, *Nieuw Archief von Wiskunde*, 12, 19-29.
- [34] Fink, A.M. (1995). Two Inequalities, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat.*, 6, 48-49.

- [35] Gavrea, B. (2000). On Hadamard's Inequality for the Convex Mappings Defined on a Convex Domain in the Space, *Journal of Ineq. in Pure & Appl. Math.*, 1(1), Article 9, 1-6.
- [36] Gill, P.M., Pearce, C.E.M. & Pečarić, J. (1997). Hadamard's Inequality for r -convex Functions, *J. of Math. Anal. and Appl.*, 215, 461-470.
- [37] Hardy, G. H., Littlewood, E.J. & Pólya, G. (1952). *Inequalities*, 2nd Ed., Cambridge University Press.
- [38] Lee, K.-C. & Tseng, K.-L. (2000). On a Weighted Generalisation of Hadamard's Inequality for G -convex Functions, *Tamsui Oxford Journal of Math. Sci.*, 16(1) , 91-104.
- [39] Lupaş, A. (1997). The Jensen-Hadamard Inequality for Convex Functions of Higher Order, *Octagon Math. Mag.*, 5(2), 8-9.
- [40] Lupaş, A. (1976). A Generalisation of Hadamard's Inequality for Convex Functions, *Univ. Beograd. Publ. Elek. Fak. Ser. Mat. Fiz.*, No. 544-576, 115-121.
- [41] Maksimović, D. M. (1979). A Short Proof of Generalized Hadamard's Inequalities, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, No. 634-677, 126-128.
- [42] Mitrinović, D.S. & Lacković, I. (1985). Hermite and Convexity, *Aequat. Math.*, 28, 229-232.
- [43] Mitrinović, D.S, Pečarić, J. & Fink, A.M. (1993) *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London.
- [44] Neuman, E. (1986). Inequalities Involving Generalised Symmetric Means, *J. Math. Anal. Appl.*, 120, 315-320.
- [45] Neuman, E. & Pečarić, J. (1989). Inequalities involving multivariate convex functions, *J. Math. Anal. Appl.*, 137, 514-549.
- [46] Neuman, E. (1990). Inequalities Involving Multivariate Convex Functions II, *Proc. Amer. Math. Soc.*, 109, 965-974.
- [47] Niculescu, C.P. (2000). A Note on the Dual Hermite-Hadamard Inequality, *The Math. Gazette*, (Romania), July, 1-3.
- [48] Niculescu, C.P. (2000). Convexity According to the Geometric Mean, *Math. Ineq. & Appl.*, 3(2) , 155-167.
- [49] Pearce, C.E.M. Pečarić, J. & Šimić, V. (1998). Stolarsky Means and Hadamard's Inequality, *J. Math. Anal. Appl.*, 220, 99-109.
- [50] Pearce, C.E.M. & Rubinov, A.M. (1999). P -functions, Quasi-convex Functions and Hadamard-type Inequalities, *J. Math. Anal. Appl.*, 240(1), 92-104.
- [51] Pečarić, J. (1992). Remarks on Two Interpolations of Hadamard's Inequalities, *Contributions, Macedonian Acad. of Sci. and Arts, Sect. of Math. and Technical Sciences*, (Scopje), 13, 9-12.
- [52] Pečarić, J. & Dragomir, S.S.(1991). A Generalization of Hadamard's Integral Inequality for Isotonic Linear Functionals, *Rudovi Mat.* (Sarajevo), 7 , 103-107.
- [53] Pečarić, J., Proschan, F. & Tong, Y. L. (1992) *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, Inc..
- [54] Sándor, J. (1990). A Note on the Jensen-Hadamard Inequality, *Anal. Numer. Theor. Approx.*, 19(1), 29-34.
- [55] Sándor, J. (1990). An Application of the Jensen-Hadamard Inequality, *Nieuw-Arch.-Wisk.*, 8(1), 63-66.
- [56] Sándor, J. (1991). On the Jensen-Hadamard inequality, *Studia Univ. Babeş-Bolyai, Math.*, 36(1), 9-15.
- [57] Vasić, P.M., Lacković, I.B. & Maksimović, D.M. (1980). Note on Convex Functions IV: On Hadamard's Inequality for Weighted Arithmetic Means, *Univ. Beograd Publ. Elek. Fak., Ser. Mat. Fiz.*, No. 678-715, 199-205.
- [58] Yang, G.S. & Hong, M.C. (1997). A Note on Hadamard's Inequality, *Tamkang J. Math.*, 28(1) , 33-37.
- [59] Yang, G.S. & Tseng, K.L. (1999). On Certain Integral Inequalities Related to Hermite-Hadamard Inequalities, *J. Math. Anal. Appl.*, 239, 180-187.

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au; ian.gomm@vu.edu.au

URL: <http://rgmia.org/dragomir>

²SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA.