

SOME FEJÉR TYPE INTEGRAL INEQUALITIES FOR GEOMETRICALLY-ARITHMETICALLY-CONVEX FUNCTIONS WITH APPLICATIONS

M. A. LATIF, S. S. DRAGOMIR^{1,2}, AND E. MOMONIAT

ABSTRACT. In this paper, the notion of geometrically symmetric functions is introduced. A new identity involving geometrically symmetric functions is established, and by using the obtained identity, the Hölder integral inequality and the notion of geometrically-arithmetically convexity, some new Fejér type integral inequalities are presented. Applications of our results to special means of positive real numbers are given as well.

1. INTRODUCTION

The classical or the usual convexity is defined as follows:

A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

holds for all $x, y \in I$ and $t \in [0, 1]$.

A number of papers have been written on inequalities using the classical convexity and one of the most fascinating inequalities in mathematical analysis is stated as follows:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}, \quad (1.1)$$

where $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping and $a, b \in I$ with $a < b$. Both the inequalities hold in reversed direction if f is concave. The inequalities stated in (1.1) are known as Hermite-Hadamard inequalities.

For more results on (1.1) which provide new proofs, noteworthy extensions, generalizations, refinements, counterparts, new Hermite-Hadamard-type inequalities and numerous applications, we refer the interested reader to [2, 3, 6, 9, 10, 20, 21] and the references therein.

The usual notion of convex functions have been generalized in diverse manners. One of them is the so called GA-convex functions and is stated in the definition below.

Date: November 15, 2012.

2000 Mathematics Subject Classification. Primary 26D15, Secondary 26A51, 26E60, 41A55.

Key words and phrases. Hermite-Hadamard's inequality, Fejér's inequality, convex function, GA-convex function, Hölder's inequality, power mean inequality.

Corresponding Author's Email Address: m_amer_latif@hotmail.com.

Corresponding Authors's Cell Number: +966-56-532-9501.

Definition 1. [13, 14] A function $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ is said to be GA-convex function on I if

$$f(x^\lambda y^{1-\lambda}) \leq \lambda f(x) + (1-\lambda)f(y)$$

holds for all $x, y \in I$ and $\lambda \in [0, 1]$, where $x^\lambda y^{1-\lambda}$ and $\lambda f(x) + (1-\lambda)f(y)$ are respectively the weighted geometric mean of two positive numbers x and y and the weighted arithmetic mean of $f(x)$ and $f(y)$.

The definition of GA-convexity is further generalized as GA- s -convexity in the second sense as follows.

Definition 2. [7] A function $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ is said to be s -GA-convex function on I if

$$f(x^\lambda y^{1-\lambda}) \leq \lambda^s f(x) + (1-\lambda)^s f(y)$$

holds for all $x, y \in I$, $\lambda \in [0, 1]$ and for some $s \in (0, 1]$.

For the properties of GA-convex functions and GA- s -convex functions, we refer the reader to [5, 7, 13, 14, 23, 24] and the reference there in.

Most recently, a number of findings have been seen on Hermite-Hadamard type integral inequalities for GA-convex and for GA- s -convex functions.

Zhang et al. in [24] established the following Hermite-Hadamard type integral inequalities for GA-convex function.

Theorem 1. [24] Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q \geq 1$, we have the following inequality:

$$\left| bf(b) - af(a) - \int_a^b f(x) dx \right| \leq \frac{[(b-a)A(a,b)]^{1-\frac{1}{q}}}{2^{\frac{1}{q}}} \times \left\{ [L(a^2, b^2) - a^2] |f'(a)|^q + [b^2 - L(a^2, b^2)] |f'(b)|^q \right\}^{\frac{1}{q}}. \quad (1.2)$$

Theorem 2. [24] Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a function differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q > 1$, we have the following inequality:

$$\left| bf(b) - af(a) - \int_a^b f(x) dx \right| \leq (\ln b - \ln a) \left[L\left(a^{\frac{2q}{q-1}}, b^{\frac{2q}{q-1}}\right) \right]^{1-\frac{1}{q}} \left[A\left(|f'(a)|^q, |f'(b)|^q\right) \right]^{\frac{1}{q}}. \quad (1.3)$$

Theorem 3. [24] Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q \geq 1$, we have the following inequality:

$$\left| bf(b) - af(a) - \int_a^b f(x) dx \right| \leq \frac{(\ln b - \ln a)^{1-\frac{1}{q}}}{(2q)^{\frac{1}{q}}} \left[L\left(a^{\frac{2q}{q-1}}, b^{\frac{2q}{q-1}}\right) \right]^{1-\frac{1}{q}} \times \left\{ [L(a^{2q}, b^{2q}) - a^{2q}] |f'(a)|^q + [b^{2q} - L(a^{2q}, b^{2q})] |f'(b)|^q \right\}^{\frac{1}{q}}. \quad (1.4)$$

Theorem 4. [24] Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q > 1$ and $2q > p > 0$. Then

$$\left| bf(b) - af(a) - \int_a^b f(x) dx \right| \leq \frac{(\ln b - \ln a)^{1-\frac{1}{q}}}{p^{\frac{1}{q}}} \left[L \left(a^{\frac{2q-p}{q-1}}, b^{\frac{2q-p}{q-1}} \right) \right]^{1-\frac{1}{q}} \\ \times \left\{ [L(a^p, b^p) - a^p] |f'(a)|^q + [b^p - L(a^p, b^p)] |f'(b)|^q \right\}^{\frac{1}{q}}. \quad (1.5)$$

Applications of the above results to special means are given in [24] as well.

İşcan [7], proved the following result for GA- s -convex functions in the second sense.

Theorem 5. Suppose that $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ is s -GA-convex in the second sense and $a, b \in I$ with $a < b$. If $f \in L([a, b])$, then one has the inequalities:

$$2^{s-1} f(\sqrt{ab}) \leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \leq \frac{f(a) + f(b)}{s+1}. \quad (1.6)$$

If f in Theorem 5 is GA-convex function, then we get the following inequalities.

$$f(\sqrt{ab}) \leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \leq \frac{f(a) + f(b)}{2}. \quad (1.7)$$

For more results on GA-convex functions and s -GA-convex functions see e.g. [5], [7], [11], [13], [22], [23] and [24].

In Section 2, we will introduce a new notion of geometrically symmetric functions and by using this notion we prove a weighted generalization of (1.7). In Section 2, we will also establish a new weighted identity to provide more general and better estimates for the difference between the right most and the middle terms of the weighted version of (1.7).

2. MAIN RESULTS

Throughout this section we take $U(t) = a^{(1-t)/2} b^{(1+t)/2}$ and $L(t) = a^{(1+t)/2} b^{(1-t)/2}$. The Beta function and the integral from of the hypergeometric function are defined as follows to be used in the sequel of the paper

$$B(\alpha, \beta) = \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt, \alpha > 0, \beta > 0$$

and

$${}_2F_1(\alpha, \beta; \gamma; z) = \frac{1}{B(\beta, \gamma - \beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-zt)^{-\alpha} dt$$

for $|z| < 1, \gamma > \beta > 0$.

The notion of geometrically symmetric functions is given in following definition.

Definition 3. A function $g : [a, b] \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ is said to be geometrically symmetric with respect to \sqrt{ab} if the inequality

$$g\left(\frac{ab}{x}\right) = g(x)$$

holds for all $x \in [a, b]$.

Theorem 6. Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function and $a, b \in I$ with $a < b$. Let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} . Then

$$f(\sqrt{ab}) \int_a^b \frac{g(x)}{x} dx \leq \int_a^b \frac{f(x)g(x)}{x} dx \leq \frac{f(a) + f(b)}{2} \int_a^b \frac{g(x)}{x} dx. \quad (2.1)$$

Proof. By the GA-convexity of f , we have

$$\begin{aligned} f(\sqrt{ab}) \int_0^1 g(a^{1-t}b^t) dt \\ \leq \int_0^1 \left[\frac{1}{2}f(a^tb^{1-t}) + \frac{1}{2}f(a^{1-t}b^t) \right] g(a^{1-t}b^t) dt \\ = \frac{1}{2} \int_0^1 f(a^tb^{1-t}) g(a^{1-t}b^t) dt + \frac{1}{2} \int_0^1 f(a^{1-t}b^t) g(a^{1-t}b^t) dt. \end{aligned} \quad (2.2)$$

By geometrical symmetry of g with respect to \sqrt{ab} , we also have

$$\int_0^1 f(a^tb^{1-t}) g(a^{1-t}b^t) dt = \int_0^1 f(a^{1-t}b^t) g(a^tb^{1-t}) dt. \quad (2.3)$$

Hence by using (2.3) in (2.2) and by the change of variables $x = a^tb^{1-t}$ and $y = a^{1-t}b^t$, we obtain

$$\begin{aligned} \frac{f(\sqrt{ab})}{\ln b - \ln a} \int_a^b \frac{g(x)}{x} dx \\ \leq \frac{1}{2} \int_0^1 f(a^tb^{1-t}) g(a^tb^{1-t}) dt + \frac{1}{2} \int_0^1 f(a^{1-t}b^t) g(a^{1-t}b^t) dt \\ = \frac{1}{2(\ln b - \ln a)} \int_a^b \frac{f(x)g(x)}{x} dx + \frac{1}{2(\ln b - \ln a)} \int_a^b \frac{f(y)g(y)}{y} dy \\ = \frac{1}{2(\ln b - \ln a)} \int_a^b \frac{f(x)g(x)}{x} dx + \frac{1}{2(\ln b - \ln a)} \int_a^b \frac{f(x)g(x)}{x} dx \\ = \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)g(x)}{x} dx. \end{aligned} \quad (2.4)$$

By the GA-convexity on $[a, b]$ and geometrical symmetry of g with respect to \sqrt{ab} , we have

$$f(a^{1-t}b^t) g(a^{1-t}b^t) \leq [(1-t)f(a) + tf(b)] g(a^{1-t}b^t) \quad (2.5)$$

and

$$f(a^tb^{1-t}) g(a^tb^{1-t}) \leq [(1-t)f(b) + tf(a)] g(a^tb^{1-t}). \quad (2.6)$$

Adding (2.5) and (2.6) and integrating with respect to t over $[0, 1]$, we obtain

$$\begin{aligned} \int_0^1 f(a^tb^{1-t}) g(a^tb^{1-t}) dt + \int_0^1 f(a^{1-t}b^t) g(a^{1-t}b^t) dt \\ \leq [f(a) + f(b)] \int_0^1 g(a^{1-t}b^t) dt. \end{aligned} \quad (2.7)$$

By the change of variables $x = a^t b^{1-t}$ and $y = a^{1-t} b^t$ in (2.7), we get

$$\frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)g(x)}{x} dx \leq \frac{[f(a) + f(b)]}{2(\ln b - \ln a)} \int_0^1 \frac{g(x)}{x} dx. \quad (2.8)$$

Combining the inequalities (2.4) and (2.8), we get the required result. \square

Now we prove a weighted integral identity which play a key role in establishing our main results.

Lemma 1. *Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} . If $f' \in L([a, b])$, then the following equality holds*

$$\begin{aligned} & \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \\ &= \frac{\ln b - \ln a}{4} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) [U(t) f'(U(t)) - L(t) f'(L(t))] dt. \end{aligned} \quad (2.9)$$

Proof. Let

$$I_1 = \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) U(t) f'(U(t)) dt$$

and

$$I_2 = \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) L(t) f'(L(t)) dt.$$

Since $g : [a, b] \rightarrow [0, \infty)$ is geometrically symmetric to \sqrt{ab} , hence $g(U(t)) = g(L(t))$ for all $t \in [0, 1]$. By this, we have

$$\begin{aligned} I_1 &= \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) U(t) f'(U(t)) dt \\ &= \frac{2}{\ln b - \ln a} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) d[f(U(t))] \\ &= \frac{2}{\ln b - \ln a} \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) f(U(t)) \Big|_0^1 - \int_0^1 [g(U(t)) + g(L(t))] f(U(t)) dt \\ &= \frac{2f(b)}{\ln b - \ln a} \int_a^b \frac{g(x)}{x} dx - 2 \int_0^1 g(U(t)) f(U(t)) dt \\ &= \frac{2f(b)}{\ln b - \ln a} \int_a^b \frac{g(x)}{x} dx - \frac{4}{\ln b - \ln a} \int_{\sqrt{ab}}^b \frac{g(x) f(x)}{x} dx. \end{aligned} \quad (2.10)$$

Analogously, we have

$$-I_2 = \frac{2f(a)}{\ln b - \ln a} \int_a^b \frac{g(x)}{x} dx - \frac{4}{\ln b - \ln a} \int_a^{\sqrt{ab}} \frac{g(x) f(x)}{x} dx. \quad (2.11)$$

Adding (2.10) and (2.11) and multiplying the result by $\frac{\ln b - \ln a}{4}$, we get the required identity. This completes the proof of the Lemma. \square

Lemma 2. For $u, v > 0$, we have

$$\int_0^1 u^{(1-t)/2} v^{(1+t)/2} dt = \sqrt{v} L(\sqrt{u}, \sqrt{v}),$$

$$\int_0^1 u^{(1+t)/2} v^{(1-t)/2} dt = \sqrt{u} L(\sqrt{u}, \sqrt{v}),$$

$$\Psi(u, v) \triangleq \frac{1}{2} \int_0^1 t u^{(1-t)/2} v^{(1+t)/2} dt = \begin{cases} \frac{v - \sqrt{v} L(\sqrt{u}, \sqrt{v})}{\ln v - \ln u}, & u \neq v \\ \frac{1}{4} u, & u = v, \end{cases}$$

and

$$\Phi(u, v) \triangleq \frac{1}{2} \int_0^1 t^2 u^{(1-t)/2} v^{(1+t)/2} dt = \begin{cases} \frac{4\sqrt{v} L(\sqrt{u}, \sqrt{v}) - 4v + v(\ln v - \ln u)}{(\ln v - \ln u)^2}, & u \neq v \\ \frac{1}{6} u, & u = v. \end{cases}$$

Proof. The proof follows from a straightforward computation. \square

We now establish new Fejér type inequalities for GA-convex functions, which provide weighted generalization of some of the results established in recent literature concerning GA-convex functions.

Theorem 7. Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} such that $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q \geq 1$, then the following inequality holds

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^2}{2^{1+1/q}} \|g\|_\infty \left\{ [\Psi(a, b)]^{1-1/q} \right. \\ & \quad \times \left([\Psi(a, b) - \Phi(a, b)] |f'(a)|^q + [\Psi(a, b) + \Phi(a, b)] |f'(b)|^q \right)^{1/q} \\ & \quad + [\Psi(b, a)]^{1-1/q} \left([\Psi(b, a) + \Phi(b, a)] |f'(a)|^q \right. \\ & \quad \left. \left. + [\Psi(b, a) - \Phi(b, a)] |f'(b)|^q \right)^{1/q} \right\}, \quad (2.12) \end{aligned}$$

where $\|g\|_\infty = \sup_{x \in [a, b]} g(x) < \infty$.

Proof. From Lemma 1 and Hölder's inequality, we have

$$\begin{aligned}
& \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\
& \leq \frac{\ln b - \ln a}{4} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) [U(t) |f'(U(t))| + L(t) |f'(L(t))|] dt \\
& \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \int_0^1 [tU(t) |f'(U(t))| + tL(t) |f'(L(t))|] dt \\
& \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \left\{ \left(\int_0^1 tU(t) \right)^{1-1/q} \left(\int_0^1 tU(t) |f'(U(t))|^q \right)^{1/q} \right. \\
& \quad \left. + \left(\int_0^1 tL(t) \right)^{1-1/q} \left(\int_0^1 tL(t) |f'(L(t))|^q \right)^{1/q} \right\}. \quad (2.13)
\end{aligned}$$

By the GA-convexity of $|f'|^q$ on $[a, b]$ for $q \geq 1$ and by using Lemma 2, we have

$$\begin{aligned}
& \int_0^1 tU(t) |f'(U(t))|^q \\
& \leq |f'(a)|^q \int_0^1 t \left(\frac{1-t}{2} \right) a^{(1-t)/2} b^{(1+t)/2} dt + |f'(b)|^q \int_0^1 t \left(\frac{1+t}{2} \right) a^{(1-t)/2} b^{(1+t)/2} dt \\
& = [\Psi(a, b) - \Phi(a, b)] |f'(a)|^q + [\Psi(a, b) + \Phi(a, b)] |f'(b)|^q \quad (2.14)
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^1 tL(t) |f'(L(t))|^q \\
& \leq |f'(a)|^q \int_0^1 t \left(\frac{1+t}{2} \right) a^{(1+t)/2} b^{(1-t)/2} dt + |f'(b)|^q \int_0^1 t \left(\frac{1-t}{2} \right) a^{(1+t)/2} b^{(1-t)/2} dt \\
& = [\Psi(b, a) + \Phi(b, a)] |f'(a)|^q + [\Psi(b, a) - \Phi(b, a)] |f'(b)|^q. \quad (2.15)
\end{aligned}$$

Using (2.14) and (2.15) in (2.13), we get the required result. This completes the proof of the theorem. \square

Corollary 1. *Suppose the assumptions of Theorem 7 are satisfied. If $q = 1$, then the following inequality holds*

$$\begin{aligned}
& \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\
& \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \left\{ [\Psi(a, b) + \Psi(b, a) - \Phi(a, b) + \Phi(b, a)] |f'(a)| \right. \\
& \quad \left. + [\Psi(a, b) + \Psi(b, a) + \Phi(a, b) - \Phi(b, a)] |f'(b)| \right\}, \quad (2.16)
\end{aligned}$$

where $\|g\|_\infty = \sup_{x \in [a, b]} g(x) < \infty$.

Corollary 2. *If $g(x) = \frac{1}{\ln b - \ln a}$, for all $x \in [a, b]$ in Theorem 7, then*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} - \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)}{2^{1+1/q}} \left\{ [\Psi(a, b)]^{1-1/q} \right. \\ & \quad \times \left([\Psi(a, b) - \Phi(a, b)] \left| f'(a) \right|^q + [\Psi(a, b) + \Phi(a, b)] \left| f'(b) \right|^q \right)^{1/q} \\ & \quad + [\Psi(b, a)]^{1-1/q} \left([\Psi(b, a) + \Phi(b, a)] \left| f'(a) \right|^q \right. \\ & \quad \left. \left. + [\Psi(b, a) - \Phi(b, a)] \left| f'(b) \right|^q \right)^{1/q} \right\}. \quad (2.17) \end{aligned}$$

Corollary 3. *If $q = 1$ in Corollary 2, then we get the following inequality*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} - \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)}{4} \left\{ [\Psi(a, b) + \Psi(b, a) + \Phi(b, a) - \Phi(a, b)] \left| f'(a) \right| \right. \\ & \quad \left. + [\Psi(b, a) + \Psi(a, b) + \Phi(a, b) - \Phi(b, a)] \left| f'(b) \right| \right\}. \quad (2.18) \end{aligned}$$

Theorem 8. *Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} such that $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q > 1$, then the following inequality holds*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^{2-1/q} \|g\|_\infty}{4 \cdot q^{1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ b^{1/2} \left([L(a^{q/2}, b^{q/2}) - a^{q/2}] \left| f'(a) \right|^q \right. \right. \\ & \quad + [2b^{q/2} - a^{q/2} - L(a^{q/2}, b^{q/2})] \left| f'(b) \right|^q \right)^{1/q} \\ & \quad + a^{1/2} \left([L(a^{q/2}, b^{q/2}) + b^{q/2} - 2a^{q/2}] \left| f'(a) \right|^q \right. \\ & \quad \left. \left. + [b^{q/2} - L(a^{q/2}, b^{q/2})] \left| f'(b) \right|^q \right)^{1/q} \right\}. \quad (2.19) \end{aligned}$$

Proof. From Lemma 1 and Hölder's inequality, we have

$$\begin{aligned}
& \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\
& \leq \frac{\ln b - \ln a}{4} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) [U(t) |f'(U(t))| + L(t) |f'(L(t))|] dt \\
& \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \int_0^1 [tU(t) |f'(U(t))| + tL(t) |f'(L(t))|] dt \\
& \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \left(\int_0^1 t^{q/(q-1)} dt \right)^{1-1/q} \\
& \times \left\{ \left(\int_0^1 [U(t)]^q |f'(U(t))|^q dt \right)^{1/q} + \left(\int_0^1 [L(t)]^q |f'(L(t))|^q dt \right)^{1/q} \right\}. \quad (2.20)
\end{aligned}$$

Since

$$\begin{aligned}
\int_0^1 [U(t)]^q |f'(U(t))|^q &= \int_0^1 a^{q(1-t)/2} b^{q(1+t)/2} |f'(a^{(1-t)/2} b^{(1+t)/2})|^q \\
&\leq |f'(a)|^q \int_0^1 \left(\frac{1-t}{2} \right) a^{q(1-t)/2} b^{q(1+t)/2} dt \\
&+ |f'(b)|^q \int_0^1 \left(\frac{1+t}{2} \right) a^{q(1-t)/2} b^{q(1+t)/2} dt \\
&= \frac{b^{q/2} [L(a^{q/2}, b^{q/2}) - a^{q/2}]}{q(\ln b - \ln a)} |f'(a)|^q \\
&+ \frac{b^{q/2} [2b^{q/2} - a^{q/2} - L(a^{q/2}, b^{q/2})]}{q(\ln b - \ln a)} |f'(b)|^q \quad (2.21)
\end{aligned}$$

and

$$\begin{aligned}
\int_0^1 [L(t)]^q |f'(L(t))|^q &= \int_0^1 a^{q(1+t)/2} b^{q(1-t)/2} |f'(a^{(1+t)/2} b^{(1-t)/2})|^q \\
&\leq |f'(a)|^q \int_0^1 \left(\frac{1+t}{2} \right) a^{q(1+t)/2} b^{q(1-t)/2} dt + |f'(b)|^q \int_0^1 \left(\frac{1-t}{2} \right) a^{q(1+t)/2} b^{q(1-t)/2} dt \\
&= \frac{a^{q/2} [L(a^{q/2}, b^{q/2}) + b^{q/2} - 2a^{q/2}]}{q(\ln b - \ln a)} |f'(a)|^q \\
&+ \frac{a^{q/2} [b^{q/2} - L(a^{q/2}, b^{q/2})]}{q(\ln b - \ln a)} |f'(b)|^q. \quad (2.22)
\end{aligned}$$

The inequality (2.19) is proved by applying (2.21) and (2.22) in (2.20). \square

Corollary 4. *If the assumptions of Theorem 8 are satisfied and if $g(x) = \frac{1}{\ln b - \ln a}$ for all $x \in [a, b]$, then the following inequality holds*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} - \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^{1-1/q}}{4 \cdot q^{1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ b^{1/2} \left([L(a^{q/2}, b^{q/2}) - a^{q/2}] |f'(a)|^q \right. \right. \\ & \quad + [2b^{q/2} - a^{q/2} - L(a^{q/2}, b^{q/2})] |f'(b)|^q \Big)^{1/q} \\ & \quad + a^{1/2} \left([L(a^{q/2}, b^{q/2}) + b^{q/2} - 2a^{q/2}] |f'(a)|^q \right. \\ & \quad \left. \left. + [b^{q/2} - L(a^{q/2}, b^{q/2})] |f'(b)|^q \right)^{1/q} \right\}. \quad (2.23) \end{aligned}$$

Theorem 9. *Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} such that $f' \in L([a, b])$. If $|f'|^q$ is GA-convex on $[a, b]$ for $q > 1$, then the following inequality holds*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^{2-1/q} \|g\|_\infty}{2(4q)^{1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ [L(a^q, b^q) - a^q] |f'(a)|^q \right. \\ & \quad \left. + [b^q - L(a^q, b^q)] |f'(b)|^q \right\}^{1/q}. \quad (2.24) \end{aligned}$$

Proof. From Lemma 1 and Hölder's inequality, we have

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{\ln b - \ln a}{4} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) [U(t) |f'(U(t))| + L(t) |f'(L(t))|] dt \\ & \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \int_0^1 [tU(t) |f'(U(t))| + tL(t) |f'(L(t))|] dt \\ & \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \left(\int_0^1 t^{q/(q-1)} dt \right)^{1-1/q} \\ & \times \left\{ \left(\int_0^1 [U(t)]^q |f'(U(t))|^q dt \right)^{1/q} + \left(\int_0^1 [L(t)]^q |f'(L(t))|^q dt \right)^{1/q} \right\}. \quad (2.25) \end{aligned}$$

By the power-mean inequality $(a^r + b^r) \leq 2^{1-r} (a + b)^r$ for $a > 0, b > 0$ and $r < 1$), we have

$$\begin{aligned} & \left(\int_0^1 [U(t)]^q |f'(U(t))|^q dt \right)^{1/q} + \left(\int_0^1 [L(t)]^q |f'(L(t))|^q dt \right)^{1/q} \\ & \leq 2^{1-1/q} \left(\int_0^1 [U(t)]^q |f'(U(t))|^q dt + \int_0^1 [L(t)]^q |f'(L(t))|^q dt \right)^{1/q}. \end{aligned} \quad (2.26)$$

Since $|f'|^q$ is GA-convex on $[a, b]$ for $q > 1$

$$\begin{aligned} & \int_0^1 [U(t)]^q |f'(U(t))|^q dt + \int_0^1 [L(t)]^q |f'(L(t))|^q dt \\ & \leq |f'(a)|^q \left[\int_0^1 \left(\frac{1-t}{2} \right) a^{q(1-t)/2} b^{q(1+t)} dt + \int_0^1 \left(\frac{1+t}{2} \right) a^{q(1+t)/2} b^{q(1-t)} dt \right] \\ & + |f'(b)|^q \left[\int_0^1 \left(\frac{1+t}{2} \right) a^{q(1-t)/2} b^{q(1+t)} dt + \int_0^1 \left(\frac{1-t}{2} \right) a^{q(1+t)/2} b^{q(1-t)} dt \right] \\ & = \left[\frac{2L(a^q, b^q) - 2a^q}{q(\ln b - \ln a)} \right] |f'(a)|^q + \left[\frac{2b^q - 2L(a^q, b^q)}{q(\ln b - \ln a)} \right] |f'(b)|^q. \end{aligned} \quad (2.27)$$

Using (2.26) in (2.27), we get

$$\begin{aligned} & \left(\int_0^1 [U(t)]^q |f'(U(t))|^q dt \right)^{1/q} + \left(\int_0^1 [L(t)]^q |f'(L(t))|^q dt \right)^{1/q} \\ & \leq 2^{1-2/q} \left(\left[\frac{L(a^q, b^q) - a^q}{q(\ln b - \ln a)} \right] |f'(a)|^q + \left[\frac{b^q - L(a^q, b^q)}{q(\ln b - \ln a)} \right] |f'(b)|^q \right)^{1/q} \end{aligned} \quad (2.28)$$

Applying (2.28) in (2.25), we obtain the required inequality (2.24). \square

Corollary 5. *If the assumptions of Theorem 9 are satisfied and if $g(x) = \frac{1}{\ln b - \ln a}$ for all $x \in [a, b]$, then the following inequality holds*

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} - \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^{1-1/q}}{2(4q)^{1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ [L(a^q, b^q) - a^q] |f'(a)|^q \right. \\ & \quad \left. + [b^q - L(a^q, b^q)] |f'(b)|^q \right\}^{1/q}. \end{aligned} \quad (2.29)$$

Theorem 10. *Let $f : I \subseteq \mathbb{R}_+ = (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$ and let $g : [a, b] \rightarrow [0, \infty)$ be continuous positive mapping and geometrically symmetric to \sqrt{ab} such that $f' \in L([a, b])$. If $|f'|$ is GA-convex on*

$[a, b]$, then the following inequality holds for $q > 1$

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{(\ln b - \ln a)^2 [L(a^{q/[2(q-1)]}, b^{q/[2(q-1)]})]^{1-1/q} \|g\|_\infty}{8} \\ & \times \left\{ \left(b^{1/2} [B(q+1, q+1)]^{1/q} + a^{1/2} \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} \right) |f'(a)| \right. \\ & \left. + \left(a^{1/2} [B(q+1, q+1)]^{1/q} + b^{1/2} \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} \right) |f'(b)| \right\}. \end{aligned} \quad (2.30)$$

Proof. From Lemma 1 and the GA-convexity of $|f'|$ on $[a, b]$, we have

$$\begin{aligned} & \left| \frac{f(b) + f(a)}{2} \int_a^b \frac{g(x)}{x} dx - \int_a^b \frac{f(x)g(x)}{x} dx \right| \\ & \leq \frac{\ln b - \ln a}{4} \int_0^1 \left(\int_{L(t)}^{U(t)} \frac{g(x)}{x} dx \right) [U(t) |f'(U(t))| + L(t) |f'(L(t))|] dt \\ & \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \int_0^1 [tU(t) |f'(U(t))| + tL(t) |f'(L(t))|] dt \\ & \leq \frac{(\ln b - \ln a)^2}{4} \|g\|_\infty \left\{ \int_0^1 a^{(1-t)/2} b^{(1+t)/2} \left[t \left(\frac{1-t}{2} \right) |f'(a)| \right. \right. \\ & \left. \left. + t \left(\frac{1+t}{2} \right) |f'(b)| \right] dt + \int_0^1 a^{(1+t)/2} b^{(1-t)/2} \left[t \left(\frac{1+t}{2} \right) |f'(a)| \right. \right. \\ & \left. \left. + t \left(\frac{1-t}{2} \right) |f'(b)| \right] dt \right\}. \end{aligned} \quad (2.31)$$

Using Hölder integral inequality, we have

$$\begin{aligned} & \int_0^1 a^{(1-t)/2} b^{(1+t)/2} \left[t \left(\frac{1-t}{2} \right) |f'(a)| + t \left(\frac{1+t}{2} \right) |f'(b)| \right] dt \\ & \leq \left(\int_0^1 a^{q(1-t)/[2(q-1)]} b^{q(1+t)/[2(q-1)]} dt \right)^{1-1/q} \\ & \times \left\{ \left[\int_0^1 t^q \left(\frac{1-t}{2} \right)^q dt \right]^{1/q} |f'(a)| + \left[\int_0^1 t^q \left(\frac{1+t}{2} \right)^q dt \right]^{1/q} |f'(b)| \right\} \\ & = \frac{1}{2} [b^{q/[2(q-1)]} L(a^{q/[2(q-1)]}, b^{q/[2(q-1)]})]^{1-1/q} \left\{ [B(q+1, q+1)]^{1/q} |f'(a)| \right. \\ & \left. + \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} |f'(b)| \right\}. \end{aligned} \quad (2.32)$$

Similarly, we one have

$$\begin{aligned}
& \int_0^1 a^{(1+t)/2} b^{(1-t)/2} \left[t \left(\frac{1+t}{2} \right) |f'(a)| + t \left(\frac{1-t}{2} \right) |f'(b)| \right] dt \\
& \leq \left(\int_0^1 a^{q(1+t)/[2(q-1)]} b^{q(1-t)/[2(q-1)]} dt \right)^{1-1/q} \\
& \quad \times \left\{ \left[\int_0^1 t^q \left(\frac{1+t}{2} \right)^q dt \right]^{1/q} |f'(a)| + \left[\int_0^1 t^q \left(\frac{1-t}{2} \right)^q dt \right]^{1/q} |f'(b)| \right\} \\
& = \frac{1}{2} \left[a^{q/[2(q-1)]} L \left(a^{q/[2(q-1)]}, b^{q/[2(q-1)]} \right) \right]^{1-1/q} \left\{ [B(q+1, q+1)]^{1/q} |f'(b)| \right. \\
& \quad \left. + \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} |f'(a)| \right\}. \quad (2.33)
\end{aligned}$$

Using (2.32) and (2.33) in (2.31), we obtain the required inequality (2.30). \square

Corollary 6. *Under the assumptions of Theorem 10, if $g(x) = \frac{1}{\ln b - \ln a}$ for all $x \in [a, b]$, then the following inequality holds*

$$\begin{aligned}
& \left| \frac{f(b) + f(a)}{2} - \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \right| \\
& \leq \frac{(\ln b - \ln a) \left[L \left(a^{q/[2(q-1)]}, b^{q/[2(q-1)]} \right) \right]^{1-1/q}}{2} \\
& \times \left\{ \left(b^{1/2} [B(q+1, q+1)]^{1/q} + a^{1/2} \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} \right) |f'(a)| \right. \\
& \quad \left. + \left(a^{1/2} [B(q+1, q+1)]^{1/q} + b^{1/2} \left[{}_2F_1(-q, q+1; q+2; -1) \cdot \frac{1}{q+1} \right]^{1/q} \right) |f'(b)| \right\}. \quad (2.34)
\end{aligned}$$

3. APPLICATIONS TO SPECIAL MEANS

In this section we apply some of the above established inequalities of Hermite-Hadamard type involving the product of a geometrically-arithmetically convex function and a geometrically symmetric function to construct inequalities for special means.

For positive numbers $a > 0$ and $b > 0$ with $a \neq b$

$$A(a, b) = \frac{a+b}{2}, \quad L(a, b) = \frac{b-a}{\ln b - \ln a}, \quad G(a, b) = \sqrt{ab}, \quad H(a, b) = \frac{2ab}{a+b}$$

and

$$L_p(a, b) = \begin{cases} \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}}, & p \neq -1, 0 \\ L(a, b), & p = -1 \\ \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}}, & p = 0 \end{cases}$$

are the arithmetic mean, the logarithmic mean, geometric mean, harmonic mean and the generalized logarithmic mean of order $p \in \mathbb{R}$ respectively. For further information on means, we refer the readers to [17, 18, 19] and the references therein.

Now let $f(x) = x^r$ for $x > 0$, $r \in \mathbb{R}$ with $r \neq 0$. Then

$$\begin{aligned} \left| f'(x^\lambda y^{1-\lambda}) \right|^q &= |r|^q \left[x^{q(r-1)} \right]^\lambda \left[y^{q(r-1)} \right]^{1-\lambda} \\ &\leq |r|^q \left[\lambda x^{q(r-1)} + (1-\lambda) y^{q(r-1)} \right] \end{aligned}$$

for $\lambda \in [0, 1]$, $x, y > 0$ and $q \geq 1$. That is $\left| f'(x) \right|^q = |r|^q x^{q(r-1)}$ is geometrically-arithmetically convex on $[a, b]$ for $q \geq 1$ and $r \neq 1$, where $a, b > 0$.

Let $g : [a, b] \rightarrow \mathbb{R}_0$ be defined as

$$g(x) = \left(\frac{x}{\sqrt{ab}} - \frac{\sqrt{ab}}{x} \right)^2, x \in [a, b].$$

It is obvious that

$$g\left(\frac{ab}{x}\right) = g(x)$$

for all $x \in [a, b]$. Hence $g(x) = \left(\frac{x}{\sqrt{ab}} - \frac{\sqrt{ab}}{x} \right)^2$, $x \in [a, b]$ is geometrically symmetric with respect to $x = \sqrt{ab}$.

Now applications of our results are given in the following theorems to come.

Theorem 11. *Let $0 < a < b$, $r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$ and $q \geq 1$. Then*

$$\begin{aligned} &\left| \frac{2A(a^r, b^r) L(a^2, b^2) - L(a^{r+2}, b^{r+2})}{[G(a, b)]^2} \right. \\ &\quad \left. - [G(a, b)]^2 L(a^{r-2}, b^{r-2}) + 2L(a^r, b^r) - 2A(a^r, b^r) \right| \\ &\leq \frac{|r|(b-a)^2}{2^{1+1/q} G(a, b) L(a, b)} \left\{ [\Psi(a, b)]^{1-1/q} \right. \\ &\times \left[2\Psi(a, b) A(a^{q(r-1)}, b^{q(r-1)}) + \Phi(a, b) (b^{q(r-1)} - a^{q(r-1)}) \right]^{1/q} + [\Psi(b, a)]^{1-1/q} \\ &\quad \left. \times \left[2\Psi(b, a) A(a^{q(r-1)}, b^{q(r-1)}) + \Phi(b, a) (a^{q(r-1)} - b^{q(r-1)}) \right]^{1/q} \right\}, \quad (3.1) \end{aligned}$$

where $\Psi(\cdot, \cdot)$ and $\Phi(\cdot, \cdot)$ are defined as in Lemma 2.

Proof. Applying Theorem 7 to the functions

$$f(x) = x^r \text{ for } x > 0, r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$$

and

$$g(x) = \left(\frac{x}{\sqrt{ab}} - \frac{\sqrt{ab}}{x} \right)^2, x \in [a, b]$$

we get the desired result. \square

Corollary 7. *Suppose the assumptions of Theorem 11 are satisfied and if $r = -1$, the the following inequality holds*

$$\begin{aligned} & \left| \frac{2L(a^2, b^2) [H(a, b)]^{-1} + L(a, b)}{[G(a, b)]^2} - [G(a, b)]^2 L(a^{-3}, b^{-3}) - 2[H(a, b)]^{-1} \right| \\ & \leq \frac{(b-a)^2}{2^{1+1/q} G(a, b) L(a, b)} \left\{ [\Psi(a, b)]^{1-1/q} \right. \\ & \quad \times [2\Psi(a, b) A(a^{-2q}, b^{-2q}) + \Phi(a, b) (b^{-2q} - a^{-2q})]^{1/q} \\ & \quad \left. + [\Psi(b, a)]^{1-1/q} [2\Psi(b, a) A(a^{-2q}, b^{-2q}) + \Phi(b, a) (a^{-2q} - b^{-2q})]^{1/q} \right\}, \quad (3.2) \end{aligned}$$

where $\Psi(\cdot, \cdot)$ and $\Phi(\cdot, \cdot)$ are defined as in Lemma 2.

Corollary 8. *Under the assumptions of Theorem 11, the following inequality holds true for $q = 1$*

$$\begin{aligned} & \left| \frac{2A(a^r, b^r) L(a^2, b^2) - L(a^{r+2}, b^{r+2})}{[G(a, b)]^2} \right. \\ & \quad \left. - [G(a, b)]^2 L(a^{r-2}, b^{r-2}) + 2L(a^r, b^r) - 2A(a^r, b^r) \right| \\ & \leq \frac{|r|(b-a)^2}{4G(a, b) L(a, b)} \left\{ L(\sqrt{a}, \sqrt{b}) A(a^{r-1}, b^{r-1}) \right. \\ & \quad \times [2A(\sqrt{a}, \sqrt{b}) - L(\sqrt{a}, \sqrt{b})] + 2(r-1) (L(\sqrt{a}, \sqrt{b}) \\ & \quad \times [L(\sqrt{a}, \sqrt{b}) - 2A(\sqrt{a}, \sqrt{b})] + A(a, b)) L(a^{r-1}, b^{r-1}) \left. \right\}. \quad (3.3) \end{aligned}$$

Corollary 9. *If we take $r = -1$ in Corollary 8, then the following inequality holds valid*

$$\begin{aligned} & \left| \frac{2L(a^2, b^2) [H(a, b)]^{-1} + L(a, b)}{[G(a, b)]^2} - [G(a, b)]^2 L(a^{-3}, b^{-3}) - 2[H(a, b)]^{-1} \right| \\ & \leq \frac{(b-a)^2}{4G(a, b) L(a, b)} \left\{ L(\sqrt{a}, \sqrt{b}) A(a^{-2}, b^{-2}) \right. \\ & \quad \times [2A(\sqrt{a}, \sqrt{b}) - L(\sqrt{a}, \sqrt{b})] - 4(L(\sqrt{a}, \sqrt{b}) \\ & \quad \times [2A(\sqrt{a}, \sqrt{b}) - L(\sqrt{a}, \sqrt{b})] - A(a, b)) L(a^{-2}, b^{-2}) \left. \right\}. \quad (3.4) \end{aligned}$$

Theorem 12. *Let $0 < a < b$, $r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$ and $q > 1$. Then*

$$\begin{aligned}
& \left| \frac{2A(a^r, b^r) L(a^2, b^2) - L(a^{r+2}, b^{r+2})}{[G(a, b)]^2} \right. \\
& \quad \left. - [G(a, b)]^2 L(a^{r-2}, b^{r-2}) + 2L(a^r, b^r) - 2A(a^r, b^r) \right| \\
& \leq \frac{(b-a)^{2-1/q} |r|}{4 \cdot q^{1/q} G(a, b) [L(a, b)]^{1-1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ b^{1/2} \left([L(a^{q/2}, b^{q/2}) - a^{q/2}] a^{q(r-1)} \right. \right. \\
& \quad \left. \left. + [2b^{q/2} - a^{q/2} - L(a^{q/2}, b^{q/2})] b^{q(r-1)} \right)^{1/q} \right. \\
& \quad \left. + a^{1/2} \left([L(a^{q/2}, b^{q/2}) + b^{q/2} - 2a^{q/2}] a^{q(r-1)} \right. \right. \\
& \quad \left. \left. + [b^{q/2} - L(a^{q/2}, b^{q/2})] b^{q(r-1)} \right)^{1/q} \right\}. \quad (3.5)
\end{aligned}$$

Proof. Applying Theorem 8 to the functions

$$f(x) = x^r \text{ for } x > 0, r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$$

and

$$g(x) = \left(\frac{x}{\sqrt{ab}} - \frac{\sqrt{ab}}{x} \right)^2, x \in [a, b]$$

we get the desired result. \square

Corollary 10. *Suppose the assumptions of Theorem 12 are fulfilled and if $r = -1$, the following inequality holds true*

$$\begin{aligned}
& \left| \frac{2[H(a, b)]^{-1} L(a^2, b^2) + L(a, b)}{[G(a, b)]^2} \right. \\
& \quad \left. - [G(a, b)]^2 L(a^{-3}, b^{-3}) - 2[H(a, b)]^{-1} \right| \\
& \leq \frac{(b-a)^{2-1/q}}{4 \cdot q^{1/q} G(a, b) [L(a, b)]^{1-1/q}} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \left\{ b^{1/2} \left([L(a^{q/2}, b^{q/2}) - a^{q/2}] a^{-2q} \right. \right. \\
& \quad \left. \left. + [2b^{q/2} - a^{q/2} - L(a^{q/2}, b^{q/2})] b^{-2q} \right)^{1/q} \right. \\
& \quad \left. + a^{1/2} \left([L(a^{q/2}, b^{q/2}) + b^{q/2} - 2a^{q/2}] a^{-2q} \right. \right. \\
& \quad \left. \left. + [b^{q/2} - L(a^{q/2}, b^{q/2})] b^{-2q} \right)^{1/q} \right\}. \quad (3.6)
\end{aligned}$$

Theorem 13. Let $0 < a < b$, $r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$ and $q > 1$. Then

$$\begin{aligned} & \left| \frac{2A(a^r, b^r) L(a^2, b^2) - L(a^{r+2}, b^{r+2})}{[G(a, b)]^2} \right. \\ & \quad \left. - [G(a, b)]^2 L(a^{r-2}, b^{r-2}) + 2L(a^r, b^r) - 2A(a^r, b^r) \right| \\ & \leq \frac{|r|(b-a)^2}{2^{2/q+1} G(a, b) L(a, b)} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \\ & \quad \times \left[rL(a^{qr}, b^{qr}) - (r-1)L(a^{q(r-1)}, b^{q(r-1)}) L(a^q, b^q) \right]^{1/q}. \quad (3.7) \end{aligned}$$

Proof. Applying Theorem 9 to the functions

$$f(x) = x^r \text{ for } x > 0, r \in \mathbb{R} \setminus \{-2, 0, 1, 2\}$$

and

$$g(x) = \left(\frac{x}{\sqrt{ab}} - \frac{\sqrt{ab}}{x} \right)^2, x \in [a, b]$$

we get the desired result. \square

Corollary 11. Suppose the assumptions of Theorem 13 are satisfied and if $r = -1$, the the following inequality holds valid

$$\begin{aligned} & \left| \frac{2[H(a, b)]^{-1} L(a^2, b^2) + L(a, b)}{[G(a, b)]^2} \right. \\ & \quad \left. - [G(a, b)]^2 L(a^{-3}, b^{-3}) - 2[H(a, b)]^{-1} \right| \\ & \leq \frac{(b-a)^2}{2^{2/q+1} G(a, b) L(a, b)} \left(\frac{2q-1}{q-1} \right)^{1-\frac{1}{q}} \\ & \quad \times \left[2L(a^{-2q}, b^{-2q}) L(a^{-1}, b^{-1}) - L(a^{-q}, b^{-q}) \right]^{1/q}. \quad (3.8) \end{aligned}$$

REFERENCES

- [1] P. S. Bullen, *Handbook of Means and Their Inequalities*, Mathematics and its Applications, Volume 560, Kluwer Academic Publishers, Dordrecht/Boston/London, 2003.
- [2] S. S. Dragomir, Hermite-Hadamard's type inequalities for convex functions of selfadjoint operators in Hilbert spaces, *Linear Algebra Appl.* 436 (2012), no. 5, 1503–1515.
- [3] S. S. Dragomir, Superadditivity and monotonicity of some functionals associated with the Hermite-Hadamard inequality for convex functions in linear spaces, *Rocky Mountain J. Math.* 42 (2012), no. 5, 1447–1459.
- [4] H. Hudzik and L. Maligranda, Some remarks on s -convex functions, *Aequationes Math.* 48 (1994), no. 1, 100–111. V. N. Huy and N. T. Chung, Some generalizations of the Fejér and Hermite-Hadamard inequalities in Hölder spaces, *J. Appl. Math. Inform.* 29 (2011), no. 3-4, 859–868.
- [5] J. Hua, B.-Y. Xi, and F. Qi, Hermite-Hadamard type inequalities for geometrically-arithmetically s -convex functions, *Commun. Korean Math. Soc.* 29 (2014), No. 1, pp. 51–63.
- [6] D. A. Ion, Some estimates on the Hermite-Hadamard inequality through quasi-convex functions, *An. Univ. Craiova Ser. Mat. Inform.* 34 (2007), 83–88.
- [7] İ. İşcan, Hermite-Hadamard type inequalities for GA- s -convex functions, *Le Matematiche*, LXIX (2014)-Fasc. II, pp. 129–146.
- [8] A. P. Ji, T. Y. Zhang, F. Qi, Integral Inequalities of Hermite-Hadamard Type for (α, m) -GA-Convex Functions, *Journal of Function Spaces and Applications*, 2013 (2013), Article ID 823856, 8 pages.

- [9] M. Klaričić Bakula and J. Pečarić, Note on some Hadamard-type inequalities, JIPAM. J. Inequal. Pure Appl. Math. 5(2004), no. 3, Article 74, 9 pp.
- [10] M. Klaričić Bakula, M. E. Özdemir and J. Pečarić, Hadamard type inequalities for m -convex and (α, m) -convex functions, JIPAM. J. Inequal. Pure Appl. Math. 9 (2008), no. 4, Article 96, 12 pp.
- [11] M. A. Latif, New Hermite-Hadamard type integral inequalities for GA-convex functions with applications, Volume 34, Issue 4 (Nov 2014).
- [12] V. G. Miheșan, A generalization of the convexity, Seminar on Functional Equations, Approx. and Convex., Cluj-Napoca, Romania, 1993.
- [13] C. P. Niculescu, Convexity according to the geometric mean, Math. Inequal. Appl. 3 (2000), no. 2, 155–167.
- [14] C. P. Niculescu, Convexity according to means, Math. Inequal. Appl. 6 (2003), no. 4, 571–579.
- [15] J. Park, Hermite-Hadamard-type inequalities for real α -star s -convex mappings, J. Appl. Math. Inform. 28 (2010), no. 5-6, 1507–1518.
- [16] J. Park, Generalizations of Hermite-Hadamard-like type inequalities for co-ordinated convex and concave mappings, Far East J. Math. Sci. (FJMS) 63 (2012), no. 2, 235–251.
- [17] F. Qi, Generalized abstracted mean values, J. Inequal. Pure Appl. Math. 1 (1) (2000), Art. 4.
- [18] F. Qi, Generalized weighted mean values with two parameters, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. 454 (1978) (1998), 2723–2732.
- [19] F. Qi and B. -Y. Xi, Some Hermite-Hadamard type inequalities for geometrically quasi-convex functions, Proc. Indian Acad. Sci. (Math. Sci.) Vol. 124, No. 3, August 2014, pp. 333–342.
- [20] M. Z. Sarikaya, On new Hermite Hadamard Fejér type integral inequalities, Stud. Univ. Babeș-Bolyai Math. 57 (2012), no. 3, 377–386.
- [21] M. Z. Sarikaya, A. Saglam and H. Yildirim, On some Hadamard-type inequalities for h -convex functions, J. Math. Inequal. 2 (2008), no. 3, 335–341.
- [22] Y. Shuang, H. P. Yin, F. Qi, Hermite-Hadamard type integral inequalities for geometric-arithmetically s -convex functions, Analysis 33, (2013), 1001–1010.
- [23] B. -Y. Xi and F. Qi, Hermite-Hadamard type inequalities for geometrically r -convex functions, Studia Scientiarum Mathematicarum Hungarica 51 (4), 530–546 (2014).
- [24] T. Y. Zhang, A. P. Ji, F. Qi, Some inequalities of Hermite-Hadamard type for GA-Convex functions with applications to means. Le Matematiche, 48 (2013), no. 1, 229–239.

SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: m.amer.latif@hotmail.com

¹SCHOOL OF ENGINEERING AND SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428 MELBOURNE CITY, MC 8001, AUSTRALIA

²SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: sever.dragomir@vu.edu.au

CENTER FOR DIFFERENTIAL EQUATIONS, CONTINUUM MECHANICS AND APPLICATIONS, SCHOOL OF COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, WITS 2050, JOHANNESBURG, SOUTH AFRICA

E-mail address: ebrahim.momoniat@wits.ac.za