

**NEW WEIGHTED INTEGRAL INEQUALITIES FOR TWICE  
DIFFERENTIABLE CONVEX FUNCTIONS**

MEHMET ZEKI SARIKAYA AND SAMET ERDEN

ABSTRACT. In this paper, we establish several new weighted inequalities for some twice differentiable mappings that are connected with the celebrated Hermite-Hadamard type and Ostrowski type integral inequalities. Some of the new inequalities are Hermite-Hadamard-type inequalities involving fractional. The results presented here would provide extensions of those given in earlier works.

1. INTRODUCTION

**Definition 1.** *The function  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ , is said to be convex if the following inequality holds*

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$

for all  $x, y \in [a, b]$  and  $\lambda \in [0, 1]$ . We say that  $f$  is concave if  $(-f)$  is convex.

The following inequality is well known in the literature as the Hermite-Hadamard integral inequality (see, [6]):

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}$$

where  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  is a convex function on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ .

A largely applied inequality for convex functions, due to its geometrical significance, is Hadamard's inequality, (see [4], [5], [18]-[23]) which has generated a wide range of directions for extension and a rich mathematical literature.

In 1938, the classical integral inequality established by Ostrowski [8] as follows:

**Theorem 1.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  whose derivative  $f' : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ , i.e.,  $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$ . Then, the inequality holds:*

$$(1.2) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for all  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is the best possible.

---

2000 *Mathematics Subject Classification.* 26D07, 26D15.

*Key words and phrases.* Hermite-Hadamard inequality, Ostrowski inequality, Trapezoid inequality, convex function, Hölder inequality, Riemann-Liouville integrals.

Inequality (1.2) has wide applications in numerical analysis and in the theory of some special means; estimating error bounds for some special means, some midpoint, trapezoid and Simpson rules and quadrature rules, etc. Hence inequality (1.2) has attracted considerable attention and interest from mathematicians and researchers. Due to this, over the years, the interested reader is also referred to ([1]-[3], [7],[9]-[16]) for integral inequalities in several independent variables. In addition, the current approach of obtaining the bounds, for a particular quadrature rule, have depended on the use of Peano kernel. The general approach in the past has involved the assumption of bounded derivatives of degree greater than one.

In this article, using functions whose twice derivatives absolute values are convex, we obtaine new weighted inequalities that are connected with the celebrated Hermite-Hadamard type and Ostrowski type integral inequalities. In addition, we obtained new inequalities of Hermite-Hadamard type and Ostrowski type involving fractional integrals. The results presented here would provide extensions of those given in earlier works.

## 2. MAIN RESULTS

Throughout this section, let us define the  $S(\alpha; w, f)$  operator as follow:

$$\begin{aligned}
S(\alpha; w, f) = & \left[ \left( \int_a^x (u-x)w(u)du \right)^\alpha - \left( \int_b^x (u-x)w(u)du \right)^\alpha \right] f'(x) \\
& + \alpha \left[ \left( \int_a^x (u-x)w(u)du \right)^{\alpha-1} \left( \int_a^x w(u)du \right) \right. \\
& \left. - \left( \int_b^x (u-x)w(u)du \right)^{\alpha-1} \left( \int_b^x w(u)du \right) \right] f(x) \\
& + \alpha(\alpha-1) \left\{ \int_a^x \left( \int_a^t (u-t)w(u)du \right)^{\alpha-2} \left( \int_a^t w(u)du \right)^2 f(t)dt \right. \\
& \left. + \int_x^b \left( \int_b^t (u-t)w(u)du \right)^{\alpha-2} \left( \int_b^t w(u)du \right)^2 f(t)dt \right\} \\
& - \alpha \left[ \int_a^x \left( \int_a^t (u-t)w(u)du \right)^{\alpha-1} w(t)f(t)dt + \int_x^b \left( \int_b^t (u-t)w(u)du \right)^{\alpha-1} w(t)f(t)dt \right].
\end{aligned}$$

In order to prove our main results we need the following lemma:

**Lemma 1.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be twice differentiable function on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ ,  $f''$  is absolutely continuous on  $[a, b]$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be nonnegative and continuous on  $[a, b]$ . Then the following identity holds:*

$$(2.1) \quad S(\alpha; w, f) = \int_a^b P_w(x, t) f''(t) dt$$

where

$$P_w(x, t) := \begin{cases} \left( \int_a^t (u-t) w(u) du \right)^\alpha, & a \leq t < x \\ \left( \int_b^t (u-t) w(u) du \right)^\alpha, & x \leq t \leq b. \end{cases}$$

for  $\alpha \geq 1$ .

*Proof.* By integration by parts, we have the following identity:

$$\begin{aligned} & \int_a^b P_w(x, t) f''(t) dt \\ &= \int_a^x \left( \int_a^t (u-t) w(u) du \right)^\alpha f''(t) dt + \int_x^b \left( \int_b^t (u-t) w(u) du \right)^\alpha f''(t) dt \\ &= \left( \int_a^t (u-t) w(u) du \right)^\alpha f'(t) \Big|_a^x + \alpha \int_a^x \left( \int_a^t (u-t) w(u) du \right)^{\alpha-1} \left( \int_a^t w(u) du \right) f'(t) dt \\ &\quad + \left( \int_b^t (u-t) w(u) du \right)^\alpha f'(t) \Big|_x^b + \alpha \int_x^b \left( \int_b^t (u-t) w(u) du \right)^{\alpha-1} \left( \int_b^t w(u) du \right) f'(t) dt \\ &= \left[ \left( \int_a^x (u-t) w(u) du \right)^\alpha - \left( \int_b^x (u-t) w(u) du \right)^\alpha \right] f'(x) \\ &\quad + \alpha \left\{ \left( \int_a^t (u-t) w(u) du \right)^{\alpha-1} \left( \int_a^t w(u) du \right) f(t) \Big|_a^x \right. \\ &\quad + \int_a^x \left[ (\alpha-1) \left( \int_a^t (u-t) w(u) du \right)^{\alpha-2} \left( \int_a^t w(u) du \right)^2 \right. \\ &\quad \left. \left. - \left( \int_a^t (u-t) w(u) du \right)^{\alpha-1} w(t) \right] f(t) dt \right. \\ &\quad \left. + \left( \int_b^t (u-t) w(u) du \right)^{\alpha-1} \left( \int_b^t w(u) du \right) f(t) \Big|_x^b \right\} \end{aligned}$$

$$\begin{aligned}
& + \int_x^b \left[ (\alpha - 1) \left( \int_b^t (u - t) w(u) du \right)^{\alpha-2} \left( \int_b^t w(u) du \right)^2 \right. \\
& \left. - \left( \int_b^t (u - t) w(u) du \right)^{\alpha-1} w(t) \right] f(t) dt \Big\} \\
= & \left[ \left( \int_a^x (u - x) w(u) du \right)^\alpha - \left( \int_b^x (u - x) w(u) du \right)^\alpha \right] f'(x) \\
& + \alpha \left[ \left( \int_a^x (u - x) w(u) du \right)^{\alpha-1} \left( \int_a^x w(u) du \right) \right. \\
& \left. - \left( \int_b^x (u - x) w(u) du \right)^{\alpha-1} \left( \int_b^x w(u) du \right) \right] f(x) \\
& + \alpha(\alpha - 1) \left\{ \int_a^x \left( \int_a^t (u - t) w(u) du \right)^{\alpha-2} \left( \int_a^t w(u) du \right)^2 f(t) dt \right. \\
& \left. + \int_x^b \left( \int_b^t (u - t) w(u) du \right)^{\alpha-2} \left( \int_b^t w(u) du \right)^2 f(t) dt \right\} \\
& - \alpha \left[ \int_a^x \left( \int_a^t (u - t) w(u) du \right)^{\alpha-1} w(t) f(t) dt + \int_x^b \left( \int_b^t (u - t) w(u) du \right)^{\alpha-1} w(t) f(t) dt \right]
\end{aligned}$$

which is the required identity in (2.1) and the proof is completed.  $\square$

**Remark 1.** Under the same assumptions of Lemma 1 with  $\alpha = 1$ , then the following identity holds:

$$\begin{aligned}
& S(1; w, f) \\
= & \left( \int_a^b (u - x) w(u) du \right) f'(x) + \left( \int_a^b w(u) du \right) f(x) - \int_a^b w(t) f(t) dt
\end{aligned}$$

which was proved by Sarikaya and Yaldiz in [16].

**Definition 2.** Let  $f \in L_1[a, b]$ . The Riemann-Liouville integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x - t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t - x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function and  $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$ .

**Corollary 1.** *Under the same assumptions of Lemma 1 with  $w(s) = 1$ , then the following identity holds:*

$$\begin{aligned}
& S(\alpha; 1, f) \\
&= \left(-\frac{1}{2}\right)^\alpha \left[ (a-x)^{2\alpha} - (b-x)^{2\alpha} \right] f'(x) \\
(2.2) \quad & + \alpha \left(-\frac{1}{2}\right)^{\alpha-1} \left[ (a-x)^{2\alpha-1} - (b-x)^{2\alpha-1} \right] f(x) \\
& + \frac{\alpha\Gamma(2\alpha)}{2} \left(-\frac{1}{2}\right)^{\alpha-2} \left[ J_{x-}^{2\alpha-1} f(a) + J_{x+}^{2\alpha-1} f(b) \right].
\end{aligned}$$

**Remark 2.** *If we take  $x = \frac{a+b}{2}$  in (2.2), we get*

$$\begin{aligned}
& \int_a^{\frac{a+b}{2}} \left( \int_a^t (u-t) du \right)^\alpha f''(t) dt + \int_{\frac{a+b}{2}}^b \left( \int_b^t (u-t) du \right)^\alpha f''(t) dt \\
&= \alpha \left(-\frac{1}{2}\right)^{\alpha-1} \frac{(b-a)^{2\alpha-1}}{2^{2\alpha-2}} f\left(\frac{a+b}{2}\right) \\
& + \frac{\alpha\Gamma(2\alpha)}{2} \left(-\frac{1}{2}\right)^{\alpha-2} \left[ J_{\left(\frac{a+b}{2}\right)-}^{2\alpha-1} f(a) + J_{\left(\frac{a+b}{2}\right)+}^{2\alpha-1} f(b) \right].
\end{aligned}$$

**Theorem 2.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be twice differentiable function on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ ,  $f''$  is absolutely continuous on  $[a, b]$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be nonnegative and continuous on  $[a, b]$ . If  $|f''|$  is convex on  $[a, b]$  then for all  $x \in [a, b]$ , the following inequalities hold:*

$$\begin{aligned}
(2.3) \quad & |S(\alpha; w, f)| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha(b-a)} \\
& \times \left[ \left( \frac{(b-a)(x-a)^{2\alpha+1}}{2\alpha+1} - \frac{(x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)| + \frac{(x-a)^{2\alpha+2}}{2\alpha+2} |f''(b)| \right] \\
& + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha(b-a)} \\
& \times \left[ \frac{(b-x)^{2\alpha+2}}{2\alpha+2} |f''(a)| + \left( \frac{(b-a)(b-x)^{2\alpha+1}}{2\alpha+1} - \frac{(b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)| \right] \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{2^\alpha(b-a)} \left[ \left( \frac{(b-a)(x-a)^{2\alpha+1}}{2\alpha+1} + \frac{(b-x)^{2\alpha+2} - (x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)| \right. \\
& \left. + \left( \frac{(b-a)(b-x)^{2\alpha+1}}{2\alpha+1} + \frac{(x-a)^{2\alpha+2} - (b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)| \right]
\end{aligned}$$

where  $\alpha \geq 1$  and  $\|w\|_\infty = \sup_{t \in [a,b]} |w(t)|$ .

*Proof.* We take absolute value of (2.1). Using bounded of the mapping  $w$ , we find that

$$\begin{aligned} |S(\alpha; w, f)| &\leq \int_a^b |P(x, t)| |f''(t)| dt \\ &= \int_a^x \left( \int_a^t (t-u) w(u) du \right)^\alpha |f''(t)| dt + \int_x^b \left( \int_t^b (u-t) w(u) du \right)^\alpha |f''(t)| dt \\ &\leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha} \int_a^x (t-a)^{2\alpha} |f''(t)| dt + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha} \int_x^b (b-t)^{2\alpha} |f''(t)| dt \end{aligned}$$

Since  $|f''(t)|$  is convex on  $[a, b] = [a, x] \cup [x, b]$

$$(2.4) \quad \left| f'' \left( \frac{b-t}{b-a}a + \frac{t-a}{b-a}b \right) \right| \leq \frac{b-t}{b-a} |f''(a)| + \frac{t-a}{b-a} |f''(b)|.$$

From (2.4), it follows that

$$\begin{aligned} &|S(\alpha; w, f)| \\ &\leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha(b-a)} \int_a^x (t-a)^{2\alpha} \left[ (b-t) |f''(a)| + (t-a) |f''(b)| \right] dt \\ &\quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha(b-a)} \int_x^b (b-t)^{2\alpha} \left[ (b-t) |f''(a)| + (t-a) |f''(b)| \right] dt \\ &= \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha(b-a)} \left[ \left( \frac{(b-a)(x-a)^{2\alpha+1}}{2\alpha+1} - \frac{(x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)| + \frac{(x-a)^{2\alpha+2}}{2\alpha+2} |f''(b)| \right] \\ &\quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha(b-a)} \left[ \frac{(b-x)^{2\alpha+2}}{2\alpha+2} |f''(a)| + \left( \frac{(b-a)(b-x)^{2\alpha+1}}{2\alpha+1} - \frac{(b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)| \right] \end{aligned}$$

and because of  $\|w\|_{[a,x],\infty} \leq \|w\|_{[a,b],\infty}$  and  $\|w\|_{[x,b],\infty} \leq \|w\|_{[a,b],\infty}$ , we obtain

$$\begin{aligned} &\leq \frac{\|w\|_{[a,b],\infty}^\alpha}{2^\alpha(b-a)} \left[ \left( \frac{(b-a)(x-a)^{2\alpha+1}}{2\alpha+1} + \frac{(b-x)^{2\alpha+2} - (x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)| \right. \\ &\quad \left. + \left( \frac{(b-a)(b-x)^{2\alpha+1}}{2\alpha+1} + \frac{(x-a)^{2\alpha+2} - (b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)| \right] \end{aligned}$$

which this completes the proof.  $\square$

**Remark 3.** Under the same assumptions of Theorem 2 with  $\alpha = 1$ , then the following inequality holds:

$$\begin{aligned} &|S(1; w, f)| \\ &\leq \frac{\|w\|_{[a,b],\infty}}{2(b-a)} \left[ \left( \frac{(b-a)(x-a)^3}{3} + \frac{(b-x)^4 - (x-a)^4}{4} \right) |f''(a)| \right. \\ &\quad \left. + \left( \frac{(b-a)(b-x)^3}{3} + \frac{(x-a)^4 - (b-x)^4}{4} \right) |f''(b)| \right] \end{aligned}$$

which was proved by Sarikaya and Yaldiz in [16].

**Corollary 2.** Under the same assumptions of Theorem 2 with  $w(s) = 1$ , then the following inequality holds:

$$(2.5) \quad |S(\alpha; 1, f)| \leq \frac{1}{2^\alpha(b-a)} \left[ \left( \frac{(b-a)(x-a)^{2\alpha+1}}{2\alpha+1} + \frac{(b-x)^{2\alpha+2} - (x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)| + \left( \frac{(b-a)(b-x)^{2\alpha+1}}{2\alpha+1} + \frac{(x-a)^{2\alpha+2} - (b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)| \right].$$

**Remark 4.** If we take  $\alpha = 1$  in (2.5), we get

$$\begin{aligned} & \left| \left( \frac{a+b}{2} - x \right) f'(x) + f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2(b-a)^2} \left[ \left( (b-a) \frac{(x-a)^3}{3} + \frac{(b-x)^4 - (x-a)^4}{4} \right) |f''(a)| + \left( \frac{(x-a)^4 - (b-x)^4}{4} + (b-a) \frac{(b-x)^3}{3} \right) |f''(b)| \right]. \end{aligned}$$

which is proved by Sarikaya and Yaldiz in [16].

**Corollary 3.** Under the same assumptions of Theorem 2 with  $w(s) = 1$  and  $x = \frac{a+b}{2}$ , then, we have

$$(2.6) \quad \begin{aligned} & \left| \alpha \left( -\frac{1}{2} \right)^{\alpha-1} \frac{(b-a)^{2\alpha-1}}{2^{2\alpha-2}} f\left( \frac{a+b}{2} \right) + \frac{\alpha\Gamma(2\alpha)}{2} \left( -\frac{1}{2} \right)^{\alpha-2} \left[ J_{\frac{a+b}{2}-}^{2\alpha-1} f(a) + J_{\frac{a+b}{2}+}^{2\alpha-1} f(b) \right] \right| \\ & \leq \frac{(b-a)^{2\alpha+1}}{(2\alpha+1)2^{3\alpha+1}} \left[ |f''(a)| + |f''(b)| \right]. \end{aligned}$$

**Remark 5.** If we take  $\alpha = 1$  in (2.6), we obtain

$$\left| f\left( \frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^2}{24} \left( \frac{|f''(a)| + |f''(b)|}{2} \right)$$

which is proved by Sarikaya and Yildirim in [17].

**Corollary 4.** Under the same assumptions of Theorem 2 with  $|f''(a)| = |f''(b)|$  in (2.3), then the following inequalities hold:

$$\begin{aligned} & |S(\alpha; w, f)| \\ & \leq \frac{1}{(2\alpha+1)2^\alpha} \left[ \|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+1} + \|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+1} \right] |f''(a)| \\ & \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{(2\alpha+1)2^\alpha} \left[ (x-a)^{2\alpha+1} + (b-x)^{2\alpha+1} \right] |f''(a)|. \end{aligned}$$

**Theorem 3.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be twice differentiable function on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ ,  $f''$  is absolutely continuous on  $[a, b]$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be nonnegative and continuous on  $[a, b]$ . If  $|f''|^q$  is convex on  $[a, b]$ ,  $q > 1$ , then for all  $x \in [a, b]$ , the following inequalities hold:

$$\begin{aligned}
& |S(\alpha; w, f)| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+\frac{1}{p}}}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+\frac{1}{p}}}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left\{ (x-a)^{2\alpha+\frac{1}{p}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\
& \quad \left. + (b-x)^{2\alpha+\frac{1}{p}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right\}
\end{aligned}$$

where  $\alpha \geq 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $\|w\|_\infty = \sup_{t \in [a,b]} |w(t)|$ .

*Proof.* We take absolute value of (2.1). Using Holder's inequality, we find that

$$\begin{aligned}
& |S(\alpha; w, f)| \\
& \leq \int_a^x \left( \int_a^t (t-u) w(u) du \right)^\alpha |f''(t)| dt + \int_x^b \left( \int_t^b (u-t) w(u) du \right)^\alpha |f''(t)| dt \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha} \int_a^x (t-a)^{2\alpha} |f''(t)| dt + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha} \int_x^b (b-t)^{2\alpha} |f''(t)| dt \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^\alpha} \left( \int_a^x (t-a)^{2\alpha p} dt \right)^{\frac{1}{p}} \left( \int_a^x |f''(t)|^q dt \right)^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^\alpha} \left( \int_x^b (b-t)^{2\alpha p} dt \right)^{\frac{1}{p}} \left( \int_x^b |f''(t)|^q dt \right)^{\frac{1}{q}}.
\end{aligned}$$

Since  $|f''(t)|^q$  is convex on  $[a, b] = [a, x] \cup [x, b]$

$$(2.7) \quad \left| f'' \left( \frac{b-t}{b-a} a + \frac{t-a}{b-a} b \right) \right|^q \leq \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q.$$



From (2.7), it follows that

$$\begin{aligned}
& |S(\alpha; w, f)| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+\frac{1}{p}}}{2^\alpha (2\alpha p+1)^{\frac{1}{p}}} \left( \int_a^x \left[ \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q \right] dt \right)^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+\frac{1}{p}}}{2^\alpha (2\alpha p+1)^{\frac{1}{p}}} \left( \int_x^b \left[ \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q \right] dt \right)^{\frac{1}{q}} \\
& = \frac{\|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+\frac{1}{p}}}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+\frac{1}{p}}}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}}
\end{aligned}$$

and because of  $\|w\|_{[a,x],\infty} \leq \|w\|_{[a,b],\infty}$  and  $\|w\|_{[x,b],\infty} \leq \|w\|_{[a,b],\infty}$ . Hence, the proof is completed.  $\square$

**Remark 6.** Under the same assumptions of Theorem 3 with  $\alpha = 1$ , then the following inequality holds:

$$\begin{aligned}
& |S(1; w, f)| \\
& \leq \frac{\|w\|_{[a,b],\infty}}{2(b-a)^{\frac{1}{q}}(2p+1)^{\frac{1}{p}}} \left\{ (x-a)^{2+\frac{1}{p}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\
& \quad \left. + (b-x)^{2+\frac{1}{p}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right\}
\end{aligned}$$

which is given by Sarikaya and Yaldiz in [16].

**Corollary 5.** Under the same assumptions of Theorem 3 with  $w(s) = 1$ , then the following inequality holds:

$$\begin{aligned}
& (2.8) \\
& |S(\alpha; 1, f)| \\
& \leq \frac{1}{2^\alpha (b-a)^{\frac{1}{q}} (2\alpha p+1)^{\frac{1}{p}}} \left\{ (x-a)^{2\alpha+\frac{1}{p}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\
& \quad \left. + (b-x)^{2\alpha+\frac{1}{p}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right\}.
\end{aligned}$$

**Remark 7.** If we take  $\alpha = 1$  in (2.8), we get

$$\begin{aligned} & \left| \left( \frac{a+b}{2} - x \right) f'(x) + f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2(b-a)^{1+\frac{1}{q}} (2p+1)^{\frac{1}{p}}} \left\{ (x-a)^{2+\frac{1}{p}} \left[ \frac{(b-a)^2 - (b-x)^2}{2} |f''(a)|^q + \frac{(x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\ & \quad \left. + (b-x)^{2+\frac{1}{p}} \left[ \frac{(b-x)^2}{2} |f''(a)|^q + \frac{(b-a)^2 - (x-a)^2}{2} |f''(b)|^q \right]^{\frac{1}{q}} \right\} \end{aligned}$$

which is given by Sarikaya and Yaldiz in [16].

**Corollary 6.** Under the same assumptions of Theorem 3 with  $w(s) = 1$  and  $x = \frac{a+b}{2}$ , then, we have

$$\begin{aligned} (2.9) \quad & \left| \alpha \left( -\frac{1}{2} \right)^{\alpha-1} \frac{(b-a)^{2\alpha-1}}{2^{2\alpha-2}} f\left( \frac{a+b}{2} \right) \right. \\ & \left. + \frac{\alpha \Gamma(2\alpha)}{2} \left( -\frac{1}{2} \right)^{\alpha-2} \left[ J_{\left( \frac{a+b}{2} \right)^-}^{2\alpha-1} f(a) + J_{\left( \frac{a+b}{2} \right)^+}^{2\alpha-1} f(b) \right] \right| \\ & \leq \frac{(b-a)^{2\alpha+1}}{(2\alpha p + 1)^{\frac{1}{p}} 2^{3\alpha+1}} \left\{ \left[ \frac{3|f''(a)|^q + |f''(b)|^q}{4} \right]^{\frac{1}{q}} + \left[ \frac{|f''(a)|^q + 3|f''(b)|^q}{4} \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 8.** If we take  $\alpha = 1$  in (2.9), we obtain

$$\begin{aligned} & \left| f\left( \frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{(b-a)^2}{16(2p+1)^{\frac{1}{p}}} \left\{ \left[ \frac{3|f''(a)|^q + |f''(b)|^q}{4} \right]^{\frac{1}{q}} \right. \\ & \quad \left. + \left[ \frac{|f''(a)|^q + 3|f''(b)|^q}{4} \right]^{\frac{1}{q}} \right\} \end{aligned}$$

which is proved by Sarikaya and Yildirim in [17].

**Theorem 4.** Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be twice differentiable function on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ ,  $f''$  is absolutely continuous on  $[a, b]$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be nonnegative and continuous on  $[a, b]$ . If  $|f''|^q$  is convex on  $[a, b]$ ,  $q > 1$ , then for all  $x \in [a, b]$ ,

the following inequalities hold:

$$\begin{aligned}
(2.10) \quad & |S(\alpha; w, f)| \\
& \leq \frac{(b-a)^{\frac{1}{q}}}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} \left[ \|w\|_{[a,x],\infty}^{\alpha p} (x-a)^{2\alpha p+1} + \|w\|_{[x,b],\infty}^{\alpha p} (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}} \\
& \quad \times \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}} \\
& \leq \frac{\|w\|_{[a,b],\infty}^\alpha (b-a)^{\frac{1}{q}}}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} \left[ (x-a)^{2\alpha p+1} + (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}
\end{aligned}$$

where  $\alpha \geq 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $\|w\|_\infty = \sup_{t \in [a,b]} |w(t)|$ .

*Proof.* We take absolute value of (2.1). Using Holder's inequality, we find that

$$\begin{aligned}
|S(\alpha; w, f)| & \leq \int_a^b |P(x, t)| |f''(t)| dt \\
& \leq \left[ \int_a^x \left( \int_a^t (t-u) w(u) du \right)^{\alpha p} dt + \int_x^b \left( \int_t^b (u-t) w(u) du \right)^{\alpha p} dt \right]^{\frac{1}{p}} \left( \int_a^b |f''(t)|^q dt \right)^{\frac{1}{q}} \\
& \leq \left[ \frac{\|w\|_{[a,x],\infty}^{\alpha p} (x-a)^{2\alpha p+1}}{2^{\alpha p}} + \frac{\|w\|_{[x,b],\infty}^{\alpha p} (b-x)^{2\alpha p+1}}{2^{\alpha p}} \right]^{\frac{1}{p}} \left( \int_a^b |f''(t)|^q dt \right)^{\frac{1}{q}}.
\end{aligned}$$

From (2.7), it follows that

$$\begin{aligned}
& |S(\alpha; w, f)| \\
& \leq \left[ \frac{\|w\|_{[a,x],\infty}^{\alpha p} (x-a)^{2\alpha p+1}}{2^{\alpha p}} + \frac{\|w\|_{[x,b],\infty}^{\alpha p} (b-x)^{2\alpha p+1}}{2^{\alpha p}} \right]^{\frac{1}{p}} \\
& \quad \times \left[ \int_a^b \left[ \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q \right] dt \right]^{\frac{1}{q}} \\
& = \frac{(b-a)^{\frac{1}{q}}}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} \left[ \|w\|_{[a,x],\infty}^{\alpha p} (x-a)^{2\alpha p+1} + \|w\|_{[x,b],\infty}^{\alpha p} (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}} \\
& \quad \times \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}
\end{aligned}$$

and because of  $\|w\|_{[a,x],\infty} \leq \|w\|_{[a,b],\infty}$  and  $\|w\|_{[x,b],\infty} \leq \|w\|_{[a,b],\infty}$ . Hence, the proof is completed.  $\square$

**Remark 9.** Under the same assumptions of Theorem 4 with  $\alpha = 1$ , then the following inequality holds:

$$\begin{aligned} & |S(1; w, f)| \\ & \leq \frac{\|w\|_{[a,b],\infty} (b-a)^{\frac{1}{q}}}{2(2p+1)^{\frac{1}{p}}} \left[ (x-a)^{2p+1} + (b-x)^{2p+1} \right]^{\frac{1}{p}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}} \end{aligned}$$

which is proved by Sarikaya and Yaldiz in [16].

**Corollary 7.** Under the same assumptions of Theorem 4 with  $w(s) = 1$ , then the following inequality holds:

$$(2.11) \quad \begin{aligned} & |S(\alpha; 1, f)| \\ & \leq \frac{(b-a)^{\frac{1}{q}}}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} \left[ (x-a)^{2\alpha p+1} + (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}. \end{aligned}$$

**Remark 10.** If we take  $\alpha = 1$  in (2.11), we get

$$\begin{aligned} & \left| \left( \frac{a+b}{2} - x \right) f'(x) + f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2(b-a)^{\frac{1}{p}} (2p+1)^{\frac{1}{p}}} \left[ (x-a)^{2p+1} + (b-x)^{2p+1} \right]^{\frac{1}{p}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}. \end{aligned}$$

**Corollary 8.** Under the same assumptions of Theorem 4 with  $w(s) = 1$  and  $x = \frac{a+b}{2}$ , then, we have

$$(2.12) \quad \begin{aligned} & \left| \alpha \left( -\frac{1}{2} \right)^{\alpha-1} \frac{(b-a)^{2\alpha-1}}{2^{2\alpha-2}} f \left( \frac{a+b}{2} \right) \right. \\ & \quad \left. + \frac{\alpha \Gamma(2\alpha)}{2} \left( -\frac{1}{2} \right)^{\alpha-2} \left[ J_{\left(\frac{a+b}{2}\right)-}^{2\alpha-1} f(a) + J_{\left(\frac{a+b}{2}\right)+}^{2\alpha-1} f(b) \right] \right| \\ & \leq \frac{(b-a)^{2\alpha+1}}{(2\alpha p + 1)^{\frac{1}{p}} 2^{3\alpha}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]. \end{aligned}$$

**Remark 11.** If we take  $\alpha = 1$  in (2.12), we obtain

$$\left| f \left( \frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^2}{8(2p+1)^{\frac{1}{p}}} \left[ \frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}$$

which is proved by Sarikaya and Yildirim in [17].

**Corollary 9.** *Under the same assumptions of Theorem 4 with  $|f''(a)|^q = |f''(b)|^q$  in (2.10), then the following inequality holds:*

$$\begin{aligned} & |S(\alpha; w, f)| \\ & \leq \frac{(b-a)^{\frac{1}{q}} |f''(a)|}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} \left[ \|w\|_{[a,x],\infty}^{\alpha p} (x-a)^{2\alpha p+1} + \|w\|_{[x,b],\infty}^{\alpha p} (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}} \\ & \leq \frac{\|w\|_{[a,b],\infty}^\alpha (b-a)^{\frac{1}{q}}}{2^\alpha (2\alpha p + 1)^{\frac{1}{p}}} |f''(a)| \left[ (x-a)^{2\alpha p+1} + (b-x)^{2\alpha p+1} \right]^{\frac{1}{p}}. \end{aligned}$$

**Theorem 5.** *Let  $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$  be twice differentiable function on  $I^\circ$ ,  $a, b \in I^\circ$  with  $a < b$ ,  $f''$  is absolutely continuous on  $[a, b]$  and let  $w : [a, b] \rightarrow \mathbb{R}$  be nonnegative and continuous on  $[a, b]$ . If  $|f''|^q$  is convex on  $[a, b]$ ,  $q \geq 1$ , then for all  $x \in [a, b]$ , the following inequalities hold:*

$$\begin{aligned} & |(2.13); w, f)| \\ & \leq \frac{\|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+1}}{2^{2\alpha} (2\alpha+1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \left[ \frac{(b-a) + (2\alpha+1)(b-x)}{(2\alpha+1)(2\alpha+2)} |f''(a)|^q + \frac{(x-a)}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} \\ & \quad + \frac{\|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+1}}{2^{2\alpha} (2\alpha+1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \left[ \frac{(b-a) + (2\alpha+1)(x-a)}{(2\alpha+1)(2\alpha+2)} |f''(b)|^q + \frac{(b-x)}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}} \\ & \leq \frac{\|w\|_{[a,b],\infty}^\alpha}{2^{2\alpha} (2\alpha+1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \\ & \quad \times \left\{ (x-a)^{2\alpha+1} \left[ \frac{(b-a) + (2\alpha+1)(b-x)}{(2\alpha+1)(2\alpha+2)} |f''(a)|^q + \frac{(x-a)}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\ & \quad \left. + (b-x)^{2\alpha+1} \left[ \frac{(b-a) + (2\alpha+1)(x-a)}{(2\alpha+1)(2\alpha+2)} |f''(b)|^q + \frac{(b-x)}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}} \right\} \end{aligned}$$

where  $\alpha \geq 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $\|w\|_\infty = \sup_{t \in [a,b]} |w(t)|$ .

*Proof.* We take absolute value of (2.1). Because of  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\alpha \left( \frac{1}{p} + \frac{1}{q} \right)$  can be written instead of  $\alpha$ . Using Holder's inequality and the convexity of  $|f''|^q$ , we find that

$$\begin{aligned} & |S(\alpha; w, f)| \\ & \leq \left( \int_a^x \left( \int_a^t (t-u) w(u) du \right)^\alpha dt \right)^{\frac{1}{p}} \left( \int_a^x \left( \int_a^t (t-u) w(u) du \right)^\alpha |f''(t)|^q dt \right)^{\frac{1}{q}} \\ & \quad + \left( \int_x^b \left( \int_t^b (u-t) w(u) du \right)^\alpha dt \right)^{\frac{1}{p}} \left( \int_x^b \left( \int_t^b (u-t) w(u) du \right)^\alpha |f''(t)|^q dt \right)^{\frac{1}{q}} \end{aligned}$$

From (2.7), it follows that

$$\begin{aligned}
& |S(\alpha; w, f)| \\
& \leq \frac{\|w\|_{[a,x],\infty}^\alpha}{2^{2\alpha}} \left( \frac{(x-a)^{2\alpha+1}}{2\alpha+1} \right)^{\frac{1}{p}} \left( \int_a^x (t-a)^{2\alpha} \left[ \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q \right] dt \right)^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^{2\alpha}} \left( \frac{(b-x)^{2\alpha+1}}{2\alpha+1} \right)^{\frac{1}{p}} \left( \int_x^b (b-t)^{2\alpha} \left[ \frac{b-t}{b-a} |f''(a)|^q + \frac{t-a}{b-a} |f''(b)|^q \right] dt \right)^{\frac{1}{q}} \\
& = \frac{\|w\|_{[a,x],\infty}^\alpha}{2^{2\alpha} (b-a)^{\frac{1}{q}}} \left( \frac{(x-a)^{2\alpha+1}}{2\alpha+1} \right)^{\frac{1}{p}} \\
& \quad \times \left[ \left( (b-a) \frac{(x-a)^{2\alpha+1}}{2\alpha+1} - \frac{(x-a)^{2\alpha+2}}{2\alpha+2} \right) |f''(a)|^q + \frac{(x-a)^{2\alpha+2}}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha}{2^{2\alpha} (b-a)^{\frac{1}{q}}} \left( \frac{(b-x)^{2\alpha+1}}{2\alpha+1} \right)^{\frac{1}{p}} \\
& \quad \times \left[ \left( (b-a) \frac{(b-x)^{2\alpha+1}}{2\alpha+1} - \frac{(b-x)^{2\alpha+2}}{2\alpha+2} \right) |f''(b)|^q + \frac{(b-x)^{2\alpha+2}}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}} \\
& = \frac{\|w\|_{[a,x],\infty}^\alpha (x-a)^{2\alpha+1}}{2^{2\alpha} (2\alpha+1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \left[ \left( \frac{(b-a)}{2\alpha+1} - \frac{(x-a)}{2\alpha+2} \right) |f''(a)|^q + \frac{(x-a)}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} \\
& \quad + \frac{\|w\|_{[x,b],\infty}^\alpha (b-x)^{2\alpha+1}}{2^{2\alpha} (2\alpha+1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \left[ \left( \frac{(b-a)}{2\alpha+1} - \frac{(b-x)}{2\alpha+2} \right) |f''(b)|^q + \frac{(b-x)}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}}
\end{aligned}$$

and because of  $\|w\|_{[a,x],\infty} \leq \|w\|_{[a,b],\infty}$  and  $\|w\|_{[x,b],\infty} \leq \|w\|_{[a,b],\infty}$ , therefore, the proof is completed.  $\square$

**Corollary 10.** *Under the same assumptions of Theorem 5 with  $\alpha = 1$ , then the following inequality holds:*

$$\begin{aligned}
& |S(1; w, f)| \\
& \leq \frac{\|w\|_{[a,b],\infty}}{4 \cdot 3^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \left\{ (x-a)^3 \left[ \frac{4b-a-3x}{12} |f''(a)|^q + \frac{(x-a)}{4} |f''(b)|^q \right]^{\frac{1}{q}} \right. \\
& \quad \left. + (b-x)^3 \left[ \frac{3x-4a+b}{12} |f''(b)|^q + \frac{(b-x)}{4} |f''(a)|^q \right]^{\frac{1}{q}} \right\}.
\end{aligned}$$

**Corollary 11.** *Under the same assumptions of Theorem 5 with  $w(s) = 1$ , then the following inequality holds:*

$$(2.14) \quad |S(\alpha; 1, f)| \leq \frac{1}{2^{2\alpha} (2\alpha + 1)^{\frac{1}{p}} (b-a)^{\frac{1}{q}}} \times \left\{ (x-a)^{2\alpha+1} \left[ \frac{(b-a) + (2\alpha+1)(b-x)}{(2\alpha+1)(2\alpha+2)} |f''(a)|^q + \frac{(x-a)}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} + (b-x)^{2\alpha+1} \left[ \frac{(b-a) + (2\alpha+1)(x-a)}{(2\alpha+1)(2\alpha+2)} |f''(b)|^q + \frac{(b-x)}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}} \right\}.$$

**Corollary 12.** *If we take  $\alpha = 1$  in (2.14), we get*

$$\left| \left( \frac{a+b}{2} - x \right) f'(x) + f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{4 \cdot 3^{\frac{1}{p}} (b-a)^{\frac{1}{q}+1}} \left\{ (x-a)^3 \left[ \frac{4b-a-3x}{12} |f''(a)|^q + \frac{(x-a)}{4} |f''(b)|^q \right]^{\frac{1}{q}} + (b-x)^3 \left[ \frac{3x-4a+b}{12} |f''(b)|^q + \frac{(b-x)}{4} |f''(a)|^q \right]^{\frac{1}{q}} \right\}.$$

**Corollary 13.** *Under the same assumptions of Theorem 5 with  $w(s) = 1$  and  $x = \frac{a+b}{2}$ , we obtain*

$$(2.15) \quad \left| \alpha \left( -\frac{1}{2} \right)^{\alpha-1} \frac{(b-a)^{2\alpha-1}}{2^{2\alpha-2}} f \left( \frac{a+b}{2} \right) + \frac{\alpha \Gamma(2\alpha)}{2} \left( -\frac{1}{2} \right)^{\alpha-2} \left[ J_{\left( \frac{a+b}{2} \right)_-}^{2\alpha-1} f(a) + J_{\left( \frac{a+b}{2} \right)_+}^{2\alpha-1} f(b) \right] \right| \leq \frac{(b-a)^{2\alpha+1}}{(2\alpha+1)^{\frac{1}{p}} 2^{4\alpha+1+\frac{1}{q}}} \left\{ \left[ \frac{(2\alpha+3)}{(2\alpha+1)(2\alpha+2)} |f''(a)|^q + \frac{1}{2\alpha+2} |f''(b)|^q \right]^{\frac{1}{q}} + \left[ \frac{(2\alpha+3)}{(2\alpha+1)(2\alpha+2)} |f''(b)|^q + \frac{1}{2\alpha+2} |f''(a)|^q \right]^{\frac{1}{q}} \right\}.$$

**Corollary 14.** *Under the same assumptions of Corollary 13 with  $\alpha = 1$ , then, we have*

$$\left| f \left( \frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{(b-a)^2}{2^{\frac{1}{q}+5} \times 3^{\frac{1}{p}}} \left\{ \left[ \frac{5 |f''(a)|^q + 3 |f''(b)|^q}{12} \right]^{\frac{1}{q}} + \left[ \frac{3 |f''(a)|^q + 5 |f''(b)|^q}{12} \right]^{\frac{1}{q}} \right\}.$$

## REFERENCES

- [1] F. Ahmad, N. S. Barnett and S. S. Dragomir, *New Weighted Ostrowski and Cebysev Type Inequalities*, *Nonlinear Analysis: Theory, Methods & Appl.*, 71 (12), (2009), 1408-1412.
- [2] F. Ahmad, A. Rafiq, N. A. Mir, *Weighted Ostrowski type inequality for twice differentiable mappings*, *Global Journal of Research in Pure and Applied Math.*, 2 (2) (2006), 147-154.
- [3] N. S. Barnett and S. S. Dragomir, *An Ostrowski type inequality for double integrals and applications for cubature formulae*, *Soochow J. Math.*, 27(1), (2001), 109-114.
- [4] S. S. Dragomir and R.P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, *Appl. Math. lett.*, 11(5) (1998), 91-95.
- [5] J. Deng and J. Wang, *Fractional Hermite-Hadamard inequalities for  $(\alpha, m)$ -logarithmically convex functions*.
- [6] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.
- [7] S. Hussain, M.A.Latif and M. Alomari, *Generalized duple-integral Ostrowski type inequalities on time scales*, *Appl. Math. Letters*, 24(2011), 1461-1467.
- [8] A. M. Ostrowski, *Über die absolutabweichung einer differentiebaren funktion von ihrem integralmittelwert*, *Comment. Math. Helv.* 10(1938), 226-227.
- [9] A. Qayyum, *A weighted Ostrowski-Grüss type inequality and applications*, *Proceeding of the World Cong. on Engineering*, Vol:2, 2009, 1-9.
- [10] A. Rafiq and F. Ahmad, *Another weighted Ostrowski-Grüss type inequality for twice differentiable mappings*, *Kragujevac Journal of Mathematics*, 31 (2008), 43-51.
- [11] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, *Acta Math. Univ. Comenianae*, Vol. LXXIX, 1(2010), pp. 129-134.
- [12] M. Z. Sarikaya *On the Ostrowski type integral inequality for double integrals*, *Demonstratio Mathematica*, Vol. XLV, No 3, 2012, pp. 533-540.
- [13] M. Z. Sarikaya and H. Ogunmez, *On the weighted Ostrowski type integral inequality for double integrals*, *The Arabian Journal for Science and Engineering (AJSE)-Mathematics*, (2011) 36:1153-1160.
- [14] M.Z.Sarikaya and S. Erden, *On the weighted integral inequalities for convex function*, *Acta Universitatis Sapientiae Mathematica*, in press.
- [15] M. Z. Sarikaya and S. Erden, *On The Hermite- Hadamard-Fejer Type Integral Inequality for Convex Function*, *Turkish Journal of Analysis and Number Theory*, 2014, Vol. 2, No. 3, 85-89.
- [16] M. Z. Sarikaya and H. Yaldiz, *Some inequalities associated with the for the probability density function*, Submitted, 2014.
- [17] M. Z. Sarikaya and H. Yildirim, *Some new integral inequalities for twice differentiable convex mappings*, *Nonlinear Analysis Forum* 17, pp. 1-14, 2012.
- [18] C.-L. Wang, X.-H. Wang, *On an extension of Hadamard inequality for convex functions*, *Chin. Ann. Math.* 3 (1982) 567-570.
- [19] S. Wasowicz and A. Witkowski, *On some inequality of Hermite-Hadamard type*, *Opuscula Math.* 32(2), (2012), pp:591-600.
- [20] S.-H. Wu, *On the weighted generalization of the Hermite-Hadamard inequality and its applications*, *The Rocky Mountain J. of Math.*, vol. 39, no. 5, pp. 1741-1749, 2009.
- [21] B-Y, Xi and F. Qi, *Some Hermite-Hadamard type inequalities for differentiable convex functions and applications*, *Hacet. J. Math. Stat.* 42(3), 243-257 (2013).
- [22] B-Y, Xi and F. Qi, *Hermite-Hadamard type inequalities for functions whose derivatives are of convexities*, *Nonlinear Funct. Anal. Appl.* 18(2), 163-176 (2013).
- [23] J. Wang, X. Li, M. Feckan and Y. Zhou, *Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity*, *Appl. Anal.* (2012). doi:10.1080/00036811.2012.727986.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, KUNURALP CAMPUS, DÜZCE-TURKEY  
*E-mail address:* sarikayamz@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BARTIN UNIVERSITY, BARTIN-TURKEY  
*E-mail address:* erdem1627@gmail.com