

Received 06/05/15

**ON EXPONENTIAL POMPEIU'S TYPE INEQUALITIES FOR
DOUBLE INTEGRALS WITH APPLICATIONS TO
OSTROWSKI'S INEQUALITY**

SAMET ERDEN AND MEHMET ZEKI SARIKAYA

ABSTRACT. The main of this paper is to derive some new inequalities of Ostrowski type using Pompeiu's mean value theorem for double integrals involving functions of two independent variables via being used exponential function.

1. INTRODUCTION

In 1938, the classical integral inequality established by Ostrowski [10] as follows:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative $f' : (a, b) \rightarrow \mathbb{R}$ is bounded on (a, b) , i.e., $\|f'\|_{\infty} = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then, the inequality holds:*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_{\infty}$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ is the best possible.

Inequality (1.1) has wide applications in numerical analysis and in the theory of some special means; estimating error bounds for some special means, some midpoint, trapezoid and Simpson rules and quadrature rules, etc. Hence inequality (1.1) has attracted considerable attention and interest from mathematicians and researchers. Due to this, over the years, the interested reader is also referred to ([2]-[5], [8],[14]-[18]) for integral inequalities in several independent variables.. In addition, the current approach of obtaining the bounds, for a particular quadrature rule, have depended on the use of Peano kernel. The general approach in the past has involved the assumption of bounded derivatives of degree greater than one.

In 1946, Pompeiu [12] derived a variant of Lagrange's mean value theorem, now known as *Pompeiu's mean value theorem*.

Theorem 2. *For every real valued function f differentiable on an interval $[a, b]$ not containing 0 and for all pairs $x_1 \neq x_2$ in $[a, b]$, there exist a point ξ between x_1 and x_2 such that*

$$\frac{x_1 f(x_2) - x_2 f(x_1)}{x_1 - x_2} = f(\xi) - \xi f'(\xi).$$

In [13], E.C. Popa using a mean value theorem obtained following theorem.

2000 Mathematics Subject Classification. 26D10, 26D15, 41A55.

Key words and phrases. Ostrowski's inequality, Pompeiu's inequalities, absolutely continuous function, double integrals.

Theorem 3. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) . Assume that $\alpha \notin [a, b]$. Then for any $x \in [a, b]$, we have the inequality

$$\begin{aligned} & \left| \left(\frac{a+b}{2} - \alpha \right) f(x) + \frac{\alpha-x}{b-a} \int_a^b f(t) dt \right| \\ & \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f - l_\alpha f'\|_\infty. \end{aligned}$$

In [11], the authors have proved the following Ostrowski type inequality:

Theorem 4. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) with $0 < a < b$. Then for $1 \leq p, q \leq \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$ we have the inequality

$$\left| \frac{a+b}{2} \cdot \frac{f(x)}{x} + \frac{1}{b-a} \int_a^b f(t) dt \right| \leq PU(x, p) \|f - lf'\|_p,$$

for $x \in [a, b]$, where

$$\begin{aligned} PU(x, p) & := (b-a)^{\frac{1}{p}-1} \left[\left(\frac{a^{2-q} - x^{2-q}}{(1-2q)(2-q)} + \frac{x^{2-q} - a^{1+q}x^{1-2q}}{(1-2q)(1+q)} \right)^{\frac{1}{p}} \right. \\ & \quad \left. + \left(\frac{b^{2-q} - x^{2-q}}{(1-2q)(2-q)} + \frac{x^{2-q} - b^{1+q}x^{1-2q}}{(1-2q)(1+q)} \right)^{\frac{1}{q}} \right]. \end{aligned}$$

In the cases $(p, q) = (1, \infty), (\infty, 1)$ and $(2, 2)$ the quantity $PU(x, p)$ has to be taken as the limit as $p \rightarrow 1, \infty$ and 2, respectively.

In [7], Dragomir has proved the Ostrowski type inequalities for complex valued absolutely continuous functions as follows:

Theorem 5. Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous function on the interval $[a, b]$ and $\alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$. Then for any $x \in [a, b]$ we have

$$\begin{aligned} & \left| \frac{\exp(\alpha b) - \exp(\alpha a)}{\alpha} f(x) - \exp(\alpha x) \int_a^b f(t) dt \right| \\ & \leq \begin{cases} |\operatorname{Re}(\alpha)| \|f' - \alpha f\|_\infty B_1(a, b, x, \alpha) & \text{if } f' - \alpha f \in L_\infty[a, b] \\ q^{\frac{1}{q}} |\operatorname{Re}(\alpha)|^{\frac{1}{q}} (b-a)^{\frac{1}{p}} \\ \times \|f' - \alpha f\|_p |B_q(a, b, x, \alpha)|^{\frac{1}{q}} & \text{if } f' - \alpha f \in L_p[a, b], \\ \|f' - \alpha f\|_1 B_\infty(a, b, x, \alpha) & \text{if } f' - \alpha f \in L_1[a, b], \end{cases} \end{aligned}$$

where

$$\begin{aligned} B_q(a, b, x, \alpha) & := \left[2 \exp(xq \operatorname{Re}(\alpha)) \left(x - \frac{a+b}{2} \right) \right. \\ & \quad \left. + \frac{1}{q \operatorname{Re}(\alpha)} \left(\frac{\exp(bq \operatorname{Re}(\alpha)) + \exp(aq \operatorname{Re}(\alpha))}{2} - \exp(xq \operatorname{Re}(\alpha)) \right) \right] \end{aligned}$$

for $q \geq 1$ and

$$B_\infty(a, b, x, \alpha) := \exp(x \operatorname{Re}(\alpha))(x - a) + \frac{\exp(b \operatorname{Re}(\alpha)) - \exp(x \operatorname{Re}(\alpha))}{\operatorname{Re}(\alpha)}.$$

The interested reader is also referred to ([1]-[9], [11], [13], [7], [4]-[22]) for integral inequalities by using Pompeiu's mean value theorem. The main aim of this paper is to establish some Pompeiu's type inequality for complex valued absolutely continuous functions with double integrals involving functions of two independent variables.

2. MAIN RESULT

To prove our theorems, we need the following lemma:

Lemma 1. *$f : \Delta = [a, b] \times [c, d] \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ and $\alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$. Then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have*

$$(2.1) \quad \begin{aligned} & \frac{f(x, y)}{\exp(\alpha x) \exp(\alpha y)} - \frac{f(x, s)}{\exp(\alpha x) \exp(\alpha s)} - \frac{f(t, y)}{\exp(\alpha t) \exp(\alpha y)} + \frac{f(t, s)}{\exp(\alpha t) \exp(\alpha s)} \\ &= \int_t^x \int_s^y \left[\frac{f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] dv du \end{aligned}$$

or, equivalently

$$(2.2) \quad \begin{aligned} & \exp(\alpha t) \exp(\alpha s) f(x, y) - \exp(\alpha y) \exp(\alpha t) f(x, s) \\ & - \exp(\alpha x) \exp(\alpha s) f(t, y) + \exp(\alpha x) \exp(\alpha y) f(t, s) \\ &= \exp(\alpha x) \exp(\alpha y) \exp(\alpha t) \exp(\alpha s) \\ & \times \int_t^x \int_s^y \left[\frac{f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] dv du. \end{aligned}$$

Proof. Since f is continuously differentiable function, $\frac{f(u, v)}{\exp(\alpha u) \exp(\alpha v)}$ is an absolutely continuous function on Δ . Then we get

$$(2.3) \quad \begin{aligned} & \int_t^x \int_s^y \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] dv du \\ &= \int_t^x \frac{\partial}{\partial u} \left(\frac{f(u, y)}{\exp(\alpha u) \exp(\alpha y)} - \frac{f(u, s)}{\exp(\alpha u) \exp(\alpha s)} \right) du \\ &= \frac{f(x, y)}{\exp(\alpha x) \exp(\alpha y)} - \frac{f(x, s)}{\exp(\alpha x) \exp(\alpha s)} \\ & - \frac{f(t, y)}{\exp(\alpha t) \exp(\alpha y)} + \frac{f(t, s)}{\exp(\alpha t) \exp(\alpha s)}, \end{aligned}$$

for all $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$. On the other hand,

$$(2.4) \quad \begin{aligned} & \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] \\ &= \frac{f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \end{aligned}$$

for all $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$. Thus, from (2.3) and (2.4), it follows that

$$\begin{aligned} & \int_t^x \int_s^y \frac{\partial^2}{\partial u \partial v} \left[\frac{f(u, v)}{u^r v^r} \right] dv du \\ &= \int_t^x \int_s^y \left[\frac{f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] dv du \\ &= \frac{f(x, y)}{\exp(\alpha x) \exp(\alpha y)} - \frac{f(x, s)}{\exp(\alpha x) \exp(\alpha s)} - \frac{f(t, y)}{\exp(\alpha t) \exp(\alpha y)} + \frac{f(t, s)}{\exp(\alpha t) \exp(\alpha s)} \end{aligned}$$

which this completes the proof. \square

Theorem 6. *$f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ and $\alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$. Then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have*

$$(2.5) \quad \begin{aligned} & |\exp(\alpha t) \exp(\alpha s) f(x, y) - \exp(\alpha t) \exp(\alpha y) f(x, s) \\ & - \exp(\alpha x) \exp(\alpha s) f(t, y) + \exp(\alpha x) \exp(\alpha y) f(t, s)| \\ & \leq \begin{cases} \operatorname{Re}^2(\alpha) \|F\|_\infty |\exp(x \operatorname{Re}(\alpha)) - \exp(t \operatorname{Re}(\alpha))| \\ \times |\exp(y \operatorname{Re}(\alpha)) - \exp(s \operatorname{Re}(\alpha))| & \text{if } F \in L_\infty(\Delta), \\ q^{\frac{2}{q}} \operatorname{Re}^{\frac{2}{q}}(\alpha) \|F\|_p \\ \times |\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))|^{\frac{1}{q}} \\ \times |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))|^{\frac{1}{q}} & \text{if } F \in L_p(\Delta), \\ \|F\|_1 \max \{|\exp(t \operatorname{Re}(\alpha))|, |\exp(x \operatorname{Re}(\alpha))|\} \\ \times \max \{|\exp(s \operatorname{Re}(\alpha))|, |\exp(y \operatorname{Re}(\alpha))|\} & \text{if } F \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F = f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f$.

Proof. From Lemma 1 and by using modulus, we have

$$\begin{aligned}
(2.6) \quad & \left| \frac{f(x, y)}{\exp(\alpha x) \exp(\alpha y)} - \frac{f(x, s)}{\exp(\alpha x) \exp(\alpha s)} - \frac{f(t, y)}{\exp(\alpha t) \exp(\alpha y)} + \frac{f(t, s)}{\exp(\alpha t) \exp(\alpha s)} \right| \\
= & \left| \int_t^x \int_s^y \left[\frac{f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)}{\exp(\alpha u) \exp(\alpha v)} \right] dv du \right| \\
\leq & \left| \int_t^x \int_s^y \frac{|f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)|}{|\exp(\alpha u) \exp(\alpha v)|} dv du \right| := |I|
\end{aligned}$$

Firstly, we will consider the case $p = \infty$ and $q = 1$. Then, we have

$$\begin{aligned}
(2.7) \quad |I| & \leq \sup_{(u, v) \in [t, x] \times [s, y]} |f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)| \\
& \times \left| \int_t^x \int_s^y \frac{1}{|\exp(\alpha u) \exp(\alpha v)|} dv du \right|.
\end{aligned}$$

Since $\alpha = \operatorname{Re}(\alpha) + i \operatorname{Im}(\alpha)$ and $(u, v) \in \Delta$, we have

$$(2.8) \quad |\exp(\alpha u) \exp(\alpha v)| = \exp(u \operatorname{Re}(\alpha)) \exp(v \operatorname{Re}(\alpha)).$$

From (2.8)

$$\begin{aligned}
(2.9) \quad & \int_t^x \int_s^y \frac{1}{|\exp(\alpha u) \exp(\alpha v)|} dv du \\
= & \int_t^x \int_s^y \frac{1}{\exp(u \operatorname{Re}(\alpha)) \exp(v \operatorname{Re}(\alpha))} dv du \\
= & \left(\int_t^x \frac{1}{\exp(u \operatorname{Re}(\alpha))} du \right) \left(\int_s^y \frac{1}{\exp(v \operatorname{Re}(\alpha))} dv \right) \\
= & \operatorname{Re}^2(\alpha) \left[\frac{1}{\exp(t \operatorname{Re}(\alpha))} - \frac{1}{\exp(x \operatorname{Re}(\alpha))} \right] \left[\frac{1}{\exp(s \operatorname{Re}(\alpha))} - \frac{1}{\exp(y \operatorname{Re}(\alpha))} \right]
\end{aligned}$$

and by (2.7) and (2.9) we obtain

$$\begin{aligned}
|I| & \leq \operatorname{Re}^2(\alpha) \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_\infty \\
& \times \left| \frac{1}{\exp(t \operatorname{Re}(\alpha))} - \frac{1}{\exp(x \operatorname{Re}(\alpha))} \right| \left| \frac{1}{\exp(s \operatorname{Re}(\alpha))} - \frac{1}{\exp(y \operatorname{Re}(\alpha))} \right|.
\end{aligned}$$

Now, consider the case $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, by using Hölder's inequality we have

$$\begin{aligned} |I| &\leq \left| \int_t^x \int_s^y |f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)|^p dv du \right|^{\frac{1}{p}} \\ &\quad \times \left| \int_t^x \int_s^y \frac{1}{|\exp(\alpha u) \exp(\alpha v)|^q} dv du \right|^{\frac{1}{q}} \\ &= q^{\frac{2}{q}} \operatorname{Re}^{\frac{2}{q}}(\alpha) \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_p \\ &\quad \times \left| \frac{1}{\exp(tq \operatorname{Re}(\alpha))} - \frac{1}{\exp(xq \operatorname{Re}(\alpha))} \right|^{\frac{1}{q}} \left| \frac{1}{\exp(sq \operatorname{Re}(\alpha))} - \frac{1}{\exp(yq \operatorname{Re}(\alpha))} \right|^{\frac{1}{q}}. \end{aligned}$$

Finally, we consider the case $p = 1$ and $q = \infty$. Then, we get

$$\begin{aligned} |I| &\leq \left| \int_t^x \int_s^y |f_{uv}(u, v) - \alpha f_u(u, v) - \alpha f_v(u, v) + \alpha^2 f(u, v)| dv du \right| \\ &\quad \times \sup_{(u,v) \in [t,x] \times [s,y]} \left(\frac{1}{\exp(\alpha u) \exp(\alpha v)} \right) \\ &= \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_1 \\ &\quad \times \frac{1}{\min \{\exp(t \operatorname{Re}(\alpha)), \exp(x \operatorname{Re}(\alpha))\}} \frac{1}{\min \{\exp(s \operatorname{Re}(\alpha)), \exp(y \operatorname{Re}(\alpha))\}}. \end{aligned}$$

This completes the proof. \square

Now, we examine some particular case of Theorem 6.

Corollary 1. *$f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$. Then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have*

$$\begin{aligned} &|\exp(t) \exp(s) f(x, y) - \exp(t) \exp(y) f(x, s) \\ &\quad - \exp(x) \exp(s) f(t, y) + \exp(x) \exp(y) f(t, s)| \\ &\leq \begin{cases} \|F_1\|_\infty |\exp(x) - \exp(t)| |\exp(y) - \exp(s)| & \text{if } F_1 \in L_\infty(\Delta), \\ q^{\frac{2}{q}} \|F_1\|_p |\exp(xq) - \exp(tq)|^{\frac{1}{q}} |\exp(yq) - \exp(sq)|^{\frac{1}{q}} & \text{if } F_1 \in L_p(\Delta), \\ \|F_1\|_1 \max \{\exp(t), \exp(x)\} \max \{\exp(s), \exp(y)\} & \text{if } F_1 \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F_1 = f_{uv} - f_u - f_v + f$.

Remark 1. If we take $\operatorname{Re}(\alpha) = 0$, then the inequality (2.5) becomes for any $(t, s), (x, y) \in \Delta$

$$(2.10) \quad \begin{aligned} & \left| \frac{f(x, y)}{\exp(i \operatorname{Im}(\alpha)x) \exp(i \operatorname{Im}(\alpha)y)} - \frac{f(x, s)}{\exp(i \operatorname{Im}(\alpha)x) \exp(i \operatorname{Im}(\alpha)s)} \right. \\ & \left. - \frac{f(t, y)}{\exp(i \operatorname{Im}(\alpha)t) \exp(i \operatorname{Im}(\alpha)y)} + \frac{f(t, s)}{\exp(i \operatorname{Im}(\alpha)t) \exp(i \operatorname{Im}(\alpha)s)} \right| \\ & \leq \begin{cases} \|F_2\|_\infty |x - t| |y - s| & \text{if } F_2 \in L_\infty(\Delta), \\ \|F_2\|_p |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} & \text{if } F_2 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \|F_2\|_1 & \text{if } F_2 \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F_2 = f_{uv} - i \operatorname{Im}(\alpha)f_u - i \operatorname{Im}(\alpha)f_v - \operatorname{Im}^2(\alpha)f$.
In particular, we have for any $(t, s), (x, y) \in \Delta$

$$\begin{aligned} & \left| \frac{f(x, y)}{\exp(ix) \exp(iy)} - \frac{f(x, s)}{\exp(ix) \exp(is)} - \frac{f(t, y)}{\exp(it) \exp(iy)} + \frac{f(t, s)}{\exp(it) \exp(is)} \right| \\ & \leq \begin{cases} \|F_3\|_\infty |x - t| |y - s| & \text{if } F_3 \in L_\infty(\Delta), \\ \|F_3\|_p |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} & \text{if } F_3 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \|F_3\|_1 & \text{if } F_3 \in L_1(\Delta) \end{cases} \end{aligned}$$

or, equivalently

$$\begin{aligned} & |\exp(it) \exp(is)f(x, y) - \exp(it) \exp(iy)f(x, s) \\ & \quad - \exp(ix) \exp(is)f(t, y) + \exp(ix) \exp(iy)f(t, s)| \\ & \leq \begin{cases} \|F_3\|_\infty |x - t| |y - s| & \text{if } F_3 \in L_\infty(\Delta), \\ \|F_3\|_p |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} & \text{if } F_3 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \|F_3\|_1 & \text{if } F_3 \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F_3 = f_{uv} - if_u - if_v - f$.

Theorem 7. $f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ and $\alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$. Then for

any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have

$$\begin{aligned}
(2.11) \quad & \left| \frac{(\exp(\alpha b) - \exp(\alpha a)) (\exp(\alpha d) - \exp(\alpha c))}{\alpha^2} f(x, y) \right. \\
& - \frac{(\exp(\alpha b) - \exp(\alpha a))}{\alpha} \exp(\alpha y) \int_c^d f(x, s) ds \\
& - \frac{(\exp(\alpha d) - \exp(\alpha c))}{\alpha} \exp(\alpha x) \int_a^b f(t, y) dt \\
& \left. + \exp(\alpha x) \exp(\alpha y) \int_a^b \int_c^d f(t, s) ds dt \right| \\
& \leq \begin{cases} \operatorname{Re}^2(\alpha) \|F\|_\infty B_1(a, b, x, \alpha) B_1(c, d, y, \alpha) & \text{if } F \in L_\infty(\Delta), \\ q^{\frac{2}{q}} \operatorname{Re}^{\frac{2}{q}}(\alpha) (b-a)^{\frac{1}{p}} (d-c)^{\frac{1}{p}} \\ \times \|F\|_p |B_q(a, b, x, \alpha)|^{\frac{1}{q}} |B_q(c, d, y, \alpha)|^{\frac{1}{q}} & \text{if } F \in L_p(\Delta), \\ \|F\|_1 B_\infty(a, b, x, \alpha) B_\infty(c, d, y, \alpha) & \text{if } F \in L_1(\Delta) \end{cases}
\end{aligned}$$

where $F = f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f$,

$$\begin{aligned}
B_q(a, b, x, \alpha) & : = 2 \left[\exp(xq \operatorname{Re}(\alpha)) \left(x - \frac{a+b}{2} \right) \right. \\
& \left. + \frac{1}{q \operatorname{Re}(\alpha)} \left(\frac{\exp(bq \operatorname{Re}(\alpha)) + \exp(aq \operatorname{Re}(\alpha))}{2} - \exp(xq \operatorname{Re}(\alpha)) \right) \right]
\end{aligned}$$

for $q \geq 1$ and

$$B_\infty(a, b, x, \alpha) := \exp(x \operatorname{Re}(\alpha))(x-a) + \frac{\exp(b \operatorname{Re}(\alpha)) - \exp(x \operatorname{Re}(\alpha))}{\operatorname{Re}(\alpha)}.$$

Proof. By using the first inequality in (2.5), it follows that

$$\begin{aligned}
|A| &:= \left| \frac{(\exp(\alpha b) - \exp(\alpha a))(\exp(\alpha d) - \exp(\alpha c))}{\alpha^2} f(x, y) \right. \\
&\quad - \frac{(\exp(\alpha b) - \exp(\alpha a))}{\alpha} \exp(\alpha y) \int_c^d f(x, s) ds \\
&\quad \left. - \frac{(\exp(\alpha d) - \exp(\alpha c))}{\alpha} \exp(\alpha x) \int_a^b f(t, y) dt + \exp(\alpha x) \exp(\alpha y) \int_a^b \int_c^d f(t, s) ds dt \right| \\
&\leq \int_a^b \int_c^d |\exp(\alpha t) \exp(\alpha s) f(x, y) - \exp(\alpha t) \exp(\alpha y) f(x, s) \\
&\quad - \exp(\alpha x) \exp(\alpha s) f(t, y) + \exp(\alpha x) \exp(\alpha y) f(t, s)| ds dt \\
&\leq \operatorname{Re}^2(\alpha) \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_\infty \\
&\quad \times \int_a^b \int_c^d |\exp(x \operatorname{Re}(\alpha)) - \exp(t \operatorname{Re}(\alpha))| |\exp(y \operatorname{Re}(\alpha)) - \exp(s \operatorname{Re}(\alpha))| ds dt.
\end{aligned}$$

Then for $\operatorname{Re}(\alpha) > 0$, we have

$$\begin{aligned}
&\int_a^b |\exp(x \operatorname{Re}(\alpha)) - \exp(t \operatorname{Re}(\alpha))| dt \\
&= \int_a^x (\exp(x \operatorname{Re}(\alpha)) - \exp(t \operatorname{Re}(\alpha))) dt + \int_x^b (\exp(t \operatorname{Re}(\alpha)) - \exp(x \operatorname{Re}(\alpha))) dt \\
&= 2 \left[\exp(x \operatorname{Re}(\alpha)) \left(x - \frac{a+b}{2} \right) \right. \\
&\quad \left. + \frac{1}{\operatorname{Re}(\alpha)} \left(\frac{\exp(b \operatorname{Re}(\alpha)) + \exp(a \operatorname{Re}(\alpha))}{2} - \exp(x \operatorname{Re}(\alpha)) \right) \right] \\
&= B_1(a, b, x, \alpha)
\end{aligned}$$

and similarly,

$$\begin{aligned}
&\int_c^d |\exp(y \operatorname{Re}(\alpha)) - \exp(s \operatorname{Re}(\alpha))| ds \\
&= \int_c^y (\exp(y \operatorname{Re}(\alpha)) - \exp(s \operatorname{Re}(\alpha))) ds + \int_y^d (\exp(s \operatorname{Re}(\alpha)) - \exp(y \operatorname{Re}(\alpha))) ds \\
&= B_1(c, d, y, \alpha)
\end{aligned}$$

thus, we obtain the first inequality in (2.11).

By using the second inequality in (2.5), it follows that

$$\begin{aligned}
|A| &\leq \int_a^b \int_c^d |\exp(\alpha t) \exp(\alpha s) f(x, y) - \exp(\alpha t) \exp(\alpha y) f(x, s) \\
&\quad - \exp(\alpha x) \exp(\alpha s) f(t, y) + \exp(\alpha x) \exp(\alpha y) f(t, s)| ds dt \\
&\leq q^{\frac{2}{q}} \operatorname{Re}^{\frac{2}{q}}(\alpha) \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_p \\
&\quad \times \int_a^b \int_c^d |\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))|^{\frac{1}{q}} |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))|^{\frac{1}{q}} ds dt.
\end{aligned}$$

By Hölder's integral inequality we also have

$$\begin{aligned}
&\int_a^b \int_c^d |\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))|^{\frac{1}{q}} |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))|^{\frac{1}{q}} ds dt \\
&= \left(\int_a^b \int_c^d ds dt \right)^{\frac{1}{p}} \\
&\quad \times \left[\int_a^b \int_c^d \left(|\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))|^{\frac{1}{q}} |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))|^{\frac{1}{q}} \right)^q ds dt \right]^{\frac{1}{q}} \\
&= (b-a)^{\frac{1}{p}} (d-c)^{\frac{1}{p}} \\
&\quad \times \left[\int_a^b \int_c^d |\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))| |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))| ds dt \right]^{\frac{1}{q}}
\end{aligned}$$

Then for $\operatorname{Re}(\alpha) > 0$, we have

$$\int_a^b |\exp(xq \operatorname{Re}(\alpha)) - \exp(tq \operatorname{Re}(\alpha))| dt = B_q(a, b, x, \alpha)$$

and similarly,

$$\int_c^d |\exp(yq \operatorname{Re}(\alpha)) - \exp(sq \operatorname{Re}(\alpha))| ds = B_q(c, d, y, \alpha)$$

thus, we obtain the second inequality in (2.11).

Finally, using the third (2.5) we have

$$\begin{aligned}
|A| &\leq \int_a^b \int_c^d |\exp(\alpha t) \exp(\alpha s) f(x, y) - \exp(\alpha t) \exp(\alpha y) f(x, s) \\
&\quad - \exp(\alpha x) \exp(\alpha s) f(t, y) + \exp(\alpha x) \exp(\alpha y) f(t, s)| ds dt \\
&\leq \|f_{uv} - \alpha f_u - \alpha f_v + \alpha^2 f\|_1 \\
&\quad \times \int_a^b \int_c^d \max \{\exp(t \operatorname{Re}(\alpha)), \exp(x \operatorname{Re}(\alpha))\} \max \{\exp(s \operatorname{Re}(\alpha)), \exp(y \operatorname{Re}(\alpha))\} ds dt.
\end{aligned}$$

Then for $\operatorname{Re}(\alpha) > 0$, we have

$$\begin{aligned}
&\int_a^b \max \{\exp(t \operatorname{Re}(\alpha)), \exp(x \operatorname{Re}(\alpha))\} dt \\
&= \int_a^x \max \{\exp(t \operatorname{Re}(\alpha)), \exp(x \operatorname{Re}(\alpha))\} dt + \int_x^b \max \{\exp(t \operatorname{Re}(\alpha)), \exp(x \operatorname{Re}(\alpha))\} dt \\
&= \int_a^x \exp(x \operatorname{Re}(\alpha)) dt + \int_x^b \exp(t \operatorname{Re}(\alpha)) dt \\
&= \exp(x \operatorname{Re}(\alpha))(x - a) + \frac{\exp(b \operatorname{Re}(\alpha)) - \exp(x \operatorname{Re}(\alpha))}{\operatorname{Re}(\alpha)} \\
&= B_\infty(a, b, x, \alpha)
\end{aligned}$$

and similarly,

$$\begin{aligned}
&\int_c^d \max \{\exp(s \operatorname{Re}(\alpha)), \exp(y \operatorname{Re}(\alpha))\} ds \\
&= B_\infty(c, d, y, \alpha)
\end{aligned}$$

and we get the third part of (2.11). Thus, This proof is completed. \square

Remark 2. If $\operatorname{Re}(\alpha) < 0$, then a similar result may be stated. Furthermore, if taken the equality (2.1) instead of the equality (2.2) in Theorem 6, then a similar inequality may be obtained. However the details are left to the interested reader.

Corollary 2. $f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$. Then for any $(t, s), (x, y) \in \Delta$ with

$x \neq y \neq t \neq s$, we have

$$\begin{aligned}
(2.12) \quad & |(\exp(b) - \exp(a)) (\exp(d) - \exp(c)) f(x, y) \\
& - (\exp(b) - \exp(a)) \exp(y) \int_c^d f(x, s) ds \\
& - (\exp(d) - \exp(c)) \exp(x) \int_a^b f(t, y) dt \\
& + \exp(x) \exp(y) \int_a^b \int_c^d f(t, s) ds dt| \\
\leq & \begin{cases} \|F_1\|_\infty B_1(a, b, x) B_1(c, d, y) & \text{if } F_1 \in L_\infty(\Delta), \\ q^{\frac{2}{q}} (b-a)^{\frac{1}{p}} (d-c)^{\frac{1}{p}} \|F_1\|_p \\ \times |B_q(a, b, x)|^{\frac{1}{q}} |B_q(c, d, y)|^{\frac{1}{q}} & \text{if } F_1 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \|F_1\|_1 B_\infty(a, b, x) B_\infty(c, d, y) & \text{if } F_1 \in L_1(\Delta) \end{cases}
\end{aligned}$$

where $F_1 = f_{uv} - f_u - f_v + f$,

$$B_q(a, b, x) := 2 \left[\exp(xq) \left(x - \frac{a+b}{2} \right) + \frac{1}{q} \left(\frac{\exp(bq) + \exp(aq)}{2} - \exp(xq) \right) \right]$$

for $q \geq 1$ and

$$B_\infty(a, b, x) := \exp(x)(x-a) + \exp(b) - \exp(x).$$

Remark 3. If we take $(x, y) = \left(\frac{a+b}{2}, \frac{c+d}{2}\right)$ in (2.12), we get

$$\begin{aligned}
& \left| (\exp(b) - \exp(a)) (\exp(d) - \exp(c)) f \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \right. \\
& - (\exp(b) - \exp(a)) \exp \left(\frac{c+d}{2} \right) \int_c^d f \left(\frac{a+b}{2}, s \right) ds \\
& - (\exp(d) - \exp(c)) \exp \left(\frac{a+b}{2} \right) \int_a^b f \left(t, \frac{c+d}{2} \right) dt \\
& \left. + \exp \left(\frac{a+b}{2} \right) \exp \left(\frac{c+d}{2} \right) \int_a^b \int_c^d f(t, s) ds dt \right| \\
\leq & \begin{cases} \|F_1\|_\infty B_1(a, b) B_1(c, d) & \text{if } F_1 \in L_\infty(\Delta) \\ q^{\frac{2}{q}} \|F_1\|_p (b-a)^{\frac{1}{p}} (d-c)^{\frac{1}{p}} \\ \times |B_q(a, b)|^{\frac{1}{q}} |B_q(c, d)|^{\frac{1}{q}} & \text{if } F_1 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \|F_1\|_1 B_\infty(a, b) B_\infty(c, d) & \text{if } F_1 \in L_1(\Delta) \end{cases}
\end{aligned}$$

where $F_1 = f_{uv} - f_u - f_v + f$,

$$B_q(a, b) := \frac{2}{q} \left(\frac{\exp(bq) + \exp(aq)}{2} - \exp\left(\frac{a+b}{2}q\right) \right)$$

for $q \geq 1$ and

$$B_\infty(a, b) := \frac{b-a}{2} \exp\left(\frac{a+b}{2}\right) + \exp(b) - \exp\left(\frac{a+b}{2}\right).$$

Theorem 8. *$f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$ and $\alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) = 0$ and $\operatorname{Im}(\alpha) \neq 0$. Then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have*

$$\begin{aligned} & (2.13) \quad \left| -\frac{(\exp(i \operatorname{Im}(\alpha) b) - \exp(i \operatorname{Im}(\alpha) a)) (\exp(i \operatorname{Im}(\alpha) d) - \exp(i \operatorname{Im}(\alpha) c))}{\operatorname{Im}(\alpha)} f(x, y) \right. \\ & \quad - \frac{(\exp(i \operatorname{Im}(\alpha) b) - \exp(i \operatorname{Im}(\alpha) a))}{i \operatorname{Im}(\alpha)} \exp(i \operatorname{Im}(\alpha) y) \int_c^d f(x, s) ds \\ & \quad - \frac{(\exp(i \operatorname{Im}(\alpha) d) - \exp(i \operatorname{Im}(\alpha) c))}{i \operatorname{Im}(\alpha)} \exp(i \operatorname{Im}(\alpha) x) \int_a^b f(t, y) dt \\ & \quad \left. + \exp(i \operatorname{Im}(\alpha) x) \exp(i \operatorname{Im}(\alpha) y) \int_a^b \int_c^d f(t, s) ds dt \right| \\ & \leq \begin{cases} (b-a)^2(d-c)^2 \|F_2\|_\infty & \text{if } F_2 \in L_\infty(\Delta), \\ \times \left[\frac{1}{4} + \left(\frac{x-\frac{a+b}{2}}{b-a} \right)^2 \right] \left[\frac{1}{4} + \left(\frac{y-\frac{c+d}{2}}{d-c} \right)^2 \right] \\ \frac{q^2(b-a)^{\frac{q+1}{q}}(d-c)^{\frac{q+1}{q}}}{(q+1)^2} \|F_2\|_p & \text{if } F_2 \in L_p(\Delta), \\ \left[\left(\frac{b-x}{b-a} \right)^{\frac{q+1}{q}} + \left(\frac{x-a}{b-a} \right)^{\frac{q+1}{q}} \right] \left[\left(\frac{d-x}{d-c} \right)^{\frac{q+1}{q}} + \left(\frac{x-c}{d-c} \right)^{\frac{q+1}{q}} \right] \\ (b-a)(d-c) \|F_2\|_1 & \text{if } F_2 \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F_2 = f_{uv} - i \operatorname{Im}(\alpha) f_u - i \operatorname{Im}(\alpha) f_v - \operatorname{Im}^2(\alpha) f$.

Proof. Utilizing the inequality (2.10) we have

$$\begin{aligned}
& \left| -\frac{(\exp(i \operatorname{Im}(\alpha) b) - \exp(i \operatorname{Im}(\alpha) a)) (\exp(i \operatorname{Im}(\alpha) d) - \exp(i \operatorname{Im}(\alpha) c))}{\operatorname{Im}(\alpha)} f(x, y) \right. \\
& \quad - \frac{(\exp(i \operatorname{Im}(\alpha) b) - \exp(i \operatorname{Im}(\alpha) a))}{i \operatorname{Im}(\alpha)} \exp(i \operatorname{Im}(\alpha) y) \int_c^d f(x, s) ds \\
& \quad - \frac{(\exp(i \operatorname{Im}(\alpha) d) - \exp(i \operatorname{Im}(\alpha) c))}{i \operatorname{Im}(\alpha)} \exp(i \operatorname{Im}(\alpha) x) \int_a^b f(t, y) dt \\
& \quad \left. + \exp(i \operatorname{Im}(\alpha) x) \exp(i \operatorname{Im}(\alpha) y) \int_a^b \int_c^d f(t, s) ds dt \right| \\
& \leq \int_a^b \int_c^d |\exp(i \operatorname{Im}(\alpha) t) \exp(i \operatorname{Im}(\alpha) s) f(x, y) - \exp(i \operatorname{Im}(\alpha) t) \exp(i \operatorname{Im}(\alpha) y) f(x, s) \\
& \quad - \exp(i \operatorname{Im}(\alpha) x) \exp(i \operatorname{Im}(\alpha) s) f(t, y) + \exp(i \operatorname{Im}(\alpha) x) \exp(i \operatorname{Im}(\alpha) y) f(t, s)| ds dt \\
& \leq \begin{cases} \|f_{uv} - i \operatorname{Im}(\alpha) f_u - i \operatorname{Im}(\alpha) f_v - \operatorname{Im}^2(\alpha) f\|_\infty \int_a^b \int_c^d |x - t| |y - s| ds dt \\ \|f_{uv} - i \operatorname{Im}(\alpha) f_u - i \operatorname{Im}(\alpha) f_v - \operatorname{Im}^2(\alpha) f\|_p \int_a^b \int_c^d |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} ds dt \\ \|f_{uv} - i \operatorname{Im}(\alpha) f_u - i \operatorname{Im}(\alpha) f_v - \operatorname{Im}^2(\alpha) f\|_1 \int_a^b \int_c^d ds dt. \end{cases}
\end{aligned}$$

Since

$$\int_a^b |x - t| dt = \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a)^2$$

and

$$\int_c^d |y - s| ds = \left[\frac{1}{4} + \left(\frac{y - \frac{c+d}{2}}{d-c} \right)^2 \right] (d-c)^2$$

we obtain the first inequality in (2.13).

Since

$$\int_a^b \int_c^d |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} ds dt = \frac{q(b-a)^{\frac{q+1}{q}}}{q+1} \left[\left(\frac{b-x}{b-a} \right)^{\frac{q+1}{q}} + \left(\frac{x-a}{b-a} \right)^{\frac{q+1}{q}} \right]$$

and

$$\int_a^b \int_c^d |x - t|^{\frac{1}{q}} |y - s|^{\frac{1}{q}} ds dt = \frac{q(d-c)^{\frac{q+1}{q}}}{q+1} \left[\left(\frac{d-x}{d-c} \right)^{\frac{q+1}{q}} + \left(\frac{x-c}{d-c} \right)^{\frac{q+1}{q}} \right]$$

we obtain the second inequality in (2.13). Hence, the proof of theorem is completed. \square

Corollary 3. *$f : \Delta \rightarrow \mathbb{C}$ be an absolutely continuous function such that the partial derivative of order 2 exists for all $(t, s) \in \Delta$. Then for any $(t, s), (x, y) \in \Delta$ with $x \neq y \neq t \neq s$, we have*

$$(2.14) \quad \begin{aligned} & | -(\exp(ib) - \exp(ia)) (\exp(id) - \exp(ic)) f(x, y) \\ & \quad - \frac{(\exp(ib) - \exp(ia))}{i} \exp(iy) \int_c^d f(x, s) ds \\ & \quad - \frac{(\exp(id) - \exp(ic))}{i} \exp(ix) \int_a^b f(t, y) dt + \exp(ix) \exp(iy) \int_a^b \int_c^d f(t, s) ds dt | \\ & \leq \begin{cases} \|F_3\|_\infty (b-a)^2 (d-c)^2 \\ \times \left[\frac{1}{4} + \left(\frac{x-\frac{a+b}{2}}{b-a} \right)^2 \right] \left[\frac{1}{4} + \left(\frac{y-\frac{c+d}{2}}{d-c} \right)^2 \right] & \text{if } F_3 \in L_\infty(\Delta), \\ \frac{q^2(b-a)^{\frac{q+1}{q}}(d-c)^{\frac{q+1}{q}}}{(q+1)^2} \|F_3\|_p & \text{if } F_3 \in L_p(\Delta), \\ \left[\left(\frac{b-x}{b-a} \right)^{\frac{q+1}{q}} + \left(\frac{x-a}{b-a} \right)^{\frac{q+1}{q}} \right] \left[\left(\frac{d-x}{d-c} \right)^{\frac{q+1}{q}} + \left(\frac{x-c}{d-c} \right)^{\frac{q+1}{q}} \right] & p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ (b-a)(d-c) \|F_3\|_1 & \text{if } F_3 \in L_1(\Delta) \end{cases} \end{aligned}$$

where $F_3 = f_{uv} - if_u - if_v - f$.

Remark 4. If we take $(x, y) = (\frac{a+b}{2}, \frac{c+d}{2})$ in (2.14), we get

$$\begin{aligned} & \left| -(\exp(ib) - \exp(ia)) (\exp(id) - \exp(ic)) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & \quad \left. - \frac{(\exp(ib) - \exp(ia))}{i} \exp\left(i\frac{c+d}{2}\right) \int_c^d f\left(\frac{a+b}{2}, s\right) ds \right. \\ & \quad \left. - \frac{(\exp(id) - \exp(ic))}{i} \exp\left(i\frac{a+b}{2}\right) \int_a^b f\left(t, \frac{c+d}{2}\right) dt \right. \\ & \quad \left. + \exp\left(i\frac{a+b}{2}\right) \exp\left(i\frac{a+b}{2}\right) \int_a^b \int_c^d f(t, s) ds dt \right| \\ & \leq \begin{cases} \frac{1}{16} \|F_3\|_\infty (b-a)^2 (d-c)^2 & \text{if } F_3 \in L_\infty(\Delta), \\ \frac{q^2(b-a)^{\frac{q+1}{q}}(d-c)^{\frac{q+1}{q}}}{(q+1)^2 2^{\frac{2}{q}}} \|F_3\|_p & \text{if } F_3 \in L_p(\Delta), \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1. & \end{cases} \end{aligned}$$

REFERENCES

- [1] A. M. Acu, A. Babos and F. D. Sofonea, *The mean value theorems and inequalities of Ostrowski type*. Sci. Stud. Res. Ser. Math. Inform. 21 (2011), no. 1, 5–16.

- [2] F. Ahmad, N. S. Barnett and S. S. Dragomir, *New Weighted Ostrowski and Cebyshev Type Inequalities*, Nonlinear Analysis: Theory, Methods & Appl., 71 (12), (2009), 1408-1412.
- [3] F. Ahmad, A. Rafiq, N. A. Mir, *Weighted Ostrowski type inequality for twice differentiable mappings*, Global Journal of Research in Pure and Applied Math., 2 (2) (2006), 147-154.
- [4] F. Ahmad, N. A. Mir and M.Z. Sarikaya, *An inequality of Ostrowski type via variant of Pompeiu's mean value theorem*, J. Basic. Appl. Sci. Res., 4(4)204-211, 2014.
- [5] N. S. Barnett and S. S. Dragomir, *An Ostrowski type inequality for double integrals and applications for cubature formulae*, Soochow J. Math., 27(1), (2001), 109-114.
- [6] S.S. Dragomir, *An inequality of Ostrowski type via Pompeiu's mean value theorem*, J. of Inequal. in Pure and Appl. Math., 6(3) (2005), Art. 83.
- [7] S. S. Dragomir, *Exponential Pompeiu's type inequalities with applications to Ostrowski's inequality*, Acta Math. Univ. Comenianae, Vol. LXXXIV, 1 (2015), pp. 39-50.
- [8] S. Hussain, M.A.Latif and M. Alomari, *Generalized double-integral Ostrowski type inequalities on time scales*, Appl. Math. Letters, 24(2011), 1461-1467.
- [9] I. Muntean, *Extensions of some mean value theorems*, Babes-Bolyai University, Faculty of Mathematics, Research Seminars on Mathematical Analysis, Preprint Nr. 7, 1991, 7-24.
- [10] A. M. Ostrowski, *Über die absolutabweichung einer differentiablen funktion von ihrem integralmittelwert*, Comment. Math. Helv. 10(1938), 226-227.
- [11] J. Pečarić and Š. Ungar, *On an inequality of Ostrowski type*, J. Ineq. Pure & Appl. Math. 7 (2006). No. 4. Art. 151.
- [12] D. Pompeiu, *Sur une proposition analogue au théorème des accroissements finis*, Mathematica (Cluj, Romania), 22(1946), 143-146.
- [13] E. C. Popa, *An inequality of Ostrowski type via a mean value theorem*, General Mathematics Vol. 15, No. 1, 2007, 93-100.
- [14] A. Qayyum, *A weighted Ostrowski-Grüss type inequality and applications*, Proceeding of the World Cong. on Engineering, Vol:2, 2009, 1-9.
- [15] A. Rafiq and F. Ahmad, *Another weighted Ostrowski-Grüss type inequality for twice differentiable mappings*, Kragujevac Journal of Mathematics, 31 (2008), 43-51.
- [16] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [17] M. Z. Sarikaya *On the Ostrowski type integral inequality for double integrals*, Demonstratio Mathematica, Vol. XLV, No 3 2012.
- [18] M. Z. Sarikaya and H. Ogummez, *On the weighted Ostrowski type integral inequality for double integrals*, The Arabian Journal for Science and Engineering (AJSE)-Mathematics, (2011) 36:1153-1160.
- [19] M. Z. Sarikaya and H. Budak, *On an inequality of Ostrowski type via variant of Pompeiu's mean value theorem*, Turkish Journal of Analysis and Number Theory, 2014, Vol. 2, No. 3, 80-84.
- [20] M. Z. Sarikaya, *On an inequality of Grüss type via variant of Pompeiu's mean value theorem*, Pure and Applied Mathematics Letters, 2(2014)26-30.
- [21] M. Z. Sarikaya, *Some new integral inequalities via variant of Pompeiu's mean value theorem*, RGMIA Research Report Collection, 17(2014), Article 76, 7 pp.
- [22] M. Z. Sarikaya, S. Erden and H. Yaldiz, *On power Pompeiu's type inequalities for double integrals*, Application and Applied Mathematics, in press.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BARTIN UNIVERSITY, BARTIN-TURKEY
E-mail address: erdensmt@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, KONURALP CAMPUS, DÜZCE-TURKEY
E-mail address: sarikayamz@gmail.com