

**SOME INEQUALITIES FOR WEIGHTED HARMONIC AND
ARITHMETIC OPERATOR MEANS**

S. S. DRAGOMIR^{1,2}

ABSTRACT. In this paper we establish some upper and lower bounds for the difference between the weighted arithmetic and harmonic operator means under various assumption for the positive invertible operators A, B . Some applications when A, B are bounded above and below by positive constants are given as well.

1. INTRODUCTION

Throughout this paper A, B are positive invertible operators on a complex Hilbert space $(H, \langle \cdot, \cdot \rangle)$. We use the following notations for operators

$$A\nabla_{\nu}B := (1 - \nu)A + \nu B,$$

the *weighted operator arithmetic mean*,

$$A\sharp_{\nu}B := A^{1/2} \left(A^{-1/2} B A^{-1/2} \right)^{\nu} A^{1/2},$$

the *weighted operator geometric mean* and

$$A!_{\nu}B := \left((1 - \nu)A^{-1} + \nu B^{-1} \right)^{-1}$$

the *weighted operator harmonic mean*, where $\nu \in [0, 1]$.

When $\nu = \frac{1}{2}$, we write $A\nabla B$, $A\sharp B$ and $A!B$ for brevity, respectively.

The following fundamental inequality between the weighted arithmetic, geometric and harmonic operator means holds

$$(1.1) \quad A!_{\nu}B \leq A\sharp_{\nu}B \leq A\nabla_{\nu}B$$

for any $\nu \in [0, 1]$.

For various recent inequalities between these means we recommend the recent papers [2]-[5], [7]-[10] and the references therein.

In this paper we establish some upper and lower bounds for the difference $A\nabla_{\nu}B - A!_{\nu}B$ for $\nu \in [0, 1]$ under various assumption for the positive invertible operators A, B . Some applications when A, B are bounded above and below by positive constants are given as well.

1991 *Mathematics Subject Classification.* 47A63, 47A30, 15A60, 26D15; 26D10.

Key words and phrases. Young's Inequality, Convex functions, Arithmetic mean-Harmonic mean inequality, Operator means, Operator inequalities.

2. MAIN RESULTS

We have the following result:

Theorem 1. *Let A, B be positive invertible operators. Then for any $\nu \in [0, 1]$ we have*

$$(2.1) \quad \begin{aligned} rA(B-A)A^{-1}(B-A)(B+A)^{-1}A \\ \leq A\nabla_{\nu}B - A!_{\nu}B \\ \leq RA(B-A)A^{-1}(B-A)(B+A)^{-1}A, \end{aligned}$$

where $r = \min\{\nu, 1 - \nu\}$ and $R = \max\{\nu, 1 - \nu\}$.

Proof. Recall the following result obtained by Dragomir in 2006 [6] that provides a refinement and a reverse for the weighted Jensen's discrete inequality:

$$(2.2) \quad \begin{aligned} n \min_{j \in \{1, 2, \dots, n\}} \{p_j\} \left[\frac{1}{n} \sum_{j=1}^n \Phi(x_j) - \Phi\left(\frac{1}{n} \sum_{j=1}^n x_j\right) \right] \\ \leq \frac{1}{P_n} \sum_{j=1}^n p_j \Phi(x_j) - \Phi\left(\frac{1}{P_n} \sum_{j=1}^n p_j x_j\right) \\ \leq n \max_{j \in \{1, 2, \dots, n\}} \{p_j\} \left[\frac{1}{n} \sum_{j=1}^n \Phi(x_j) - \Phi\left(\frac{1}{n} \sum_{j=1}^n x_j\right) \right], \end{aligned}$$

where $\Phi : C \rightarrow \mathbb{R}$ is a convex function defined on convex subset C of the linear space X , $\{x_j\}_{j \in \{1, 2, \dots, n\}}$ are vectors in C and $\{p_j\}_{j \in \{1, 2, \dots, n\}}$ are nonnegative numbers with $P_n = \sum_{j=1}^n p_j > 0$.

For $n = 2$, we deduce from (2.2) that

$$(2.3) \quad \begin{aligned} 2 \min\{\nu, 1 - \nu\} \left[\frac{\Phi(x) + \Phi(y)}{2} - \Phi\left(\frac{x+y}{2}\right) \right] \\ \leq \nu\Phi(x) + (1 - \nu)\Phi(y) - \Phi[\nu x + (1 - \nu)y] \\ \leq 2 \max\{\nu, 1 - \nu\} \left[\frac{\Phi(x) + \Phi(y)}{2} - \Phi\left(\frac{x+y}{2}\right) \right] \end{aligned}$$

for any $x, y \in C$ and $\nu \in [0, 1]$.

If we write the inequality (2.3) for the convex function $\Phi(x) = \frac{1}{x}$, $x > 0$, then we have

$$(2.4) \quad \begin{aligned} 2r \left(\frac{\frac{1}{x} + \frac{1}{y}}{2} - \frac{2}{x+y} \right) \leq \frac{\nu}{x} + \frac{1 - \nu}{y} - \frac{1}{\nu x + (1 - \nu)y} \\ \leq 2R \left(\frac{\frac{1}{x} + \frac{1}{y}}{2} - \frac{2}{x+y} \right) \end{aligned}$$

for any $x, y > 0$ where $r = \min\{\nu, 1 - \nu\}$ and $R = \max\{\nu, 1 - \nu\}$.

If we take $y = \frac{1}{a}$, $x = \frac{1}{b}$ in (2.4), then we have

$$(2.5) \quad 2r \left(\frac{b+a}{2} - \frac{2}{\frac{1}{b} + \frac{1}{a}} \right) \leq \nu b + (1-\nu)a - (\nu b^{-1} + (1-\nu)a^{-1})^{-1} \\ \leq 2R \left(\frac{b+a}{2} - \frac{2}{\frac{1}{b} + \frac{1}{a}} \right)$$

for any $a, b > 0$ and $\nu \in [0, 1]$ where $r = \min\{\nu, 1-\nu\}$ and $R = \max\{\nu, 1-\nu\}$.
Since

$$\frac{b+a}{2} - \frac{2}{\frac{1}{b} + \frac{1}{a}} = \frac{b+a}{2} - \frac{2ab}{b+a} = \frac{1}{2} \frac{(b-a)^2}{a+b}$$

hence by (2.5) we have

$$(2.6) \quad r \frac{(b-a)^2}{a+b} \leq \nu b + (1-\nu)a - (\nu b^{-1} + (1-\nu)a^{-1})^{-1} \leq R \frac{(b-a)^2}{a+b}$$

for any $a, b > 0$ and $\nu \in [0, 1]$.

This is an inequality of interest in itself.

If we take $a = 1$ and $b = t$ in (2.6), then we get

$$(2.7) \quad r(t-1)^2(t+1)^{-1} \leq \nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq R(t-1)^2(t+1)^{-1}$$

for any $t > 0$.

If we use the continuous functional calculus for the positive invertible operator X we get

$$(2.8) \quad r(X-I)^2(X+I)^{-1} \leq \nu X + (1-\nu)I - (\nu X^{-1} + (1-\nu)I)^{-1} \\ \leq R(X-I)^2(X+I)^{-1}.$$

If we write the inequality (2.8) for $X = A^{-1/2}BA^{-1/2}$, then we get

$$(2.9) \quad r \left(A^{-1/2}BA^{-1/2} - I \right)^2 \left(A^{-1/2}BA^{-1/2} + I \right)^{-1} \\ \leq \nu A^{-1/2}BA^{-1/2} + (1-\nu)I - \left(\nu \left(A^{-1/2}BA^{-1/2} \right)^{-1} + (1-\nu)I \right)^{-1} \\ \leq R \left(A^{-1/2}BA^{-1/2} - I \right)^2 \left(A^{-1/2}BA^{-1/2} + I \right)^{-1}.$$

If we multiply the inequality (2.9) both sides with $A^{1/2}$, then we get

$$(2.10) \quad r A^{1/2} \left(A^{-1/2}BA^{-1/2} - I \right)^2 \left(A^{-1/2}BA^{-1/2} + I \right)^{-1} A^{1/2} \\ \leq \nu B + (1-\nu)A - A^{1/2} \left(\nu \left(A^{-1/2}BA^{-1/2} \right)^{-1} + (1-\nu)I \right)^{-1} A^{1/2} \\ \leq R A^{1/2} \left(A^{-1/2}BA^{-1/2} - I \right)^2 \left(A^{-1/2}BA^{-1/2} + I \right)^{-1} A^{1/2}.$$

Since

$$\begin{aligned}
& A^{1/2} \left(\nu \left(A^{-1/2} B A^{-1/2} \right)^{-1} + (1 - \nu) I \right)^{-1} A^{1/2} \\
&= A^{1/2} \left(\nu A^{1/2} B^{-1} A^{1/2} + (1 - \nu) I \right)^{-1} A^{1/2} \\
&= A^{1/2} \left(A^{1/2} \left(\nu B^{-1} + (1 - \nu) A^{-1} \right) A^{1/2} \right)^{-1} A^{1/2} \\
&= A^{1/2} \left(A^{1/2} \left(\nu B^{-1} + (1 - \nu) A^{-1} \right) A^{1/2} \right)^{-1} A^{1/2} \\
&= A^{1/2} A^{-1/2} \left(\nu B^{-1} + (1 - \nu) A^{-1} \right)^{-1} A^{-1/2} A^{1/2} = A!_{\nu} B
\end{aligned}$$

and

$$\begin{aligned}
& A^{1/2} \left(A^{-1/2} B A^{-1/2} - I \right)^2 \left(A^{-1/2} B A^{-1/2} + I \right)^{-1} A^{1/2} \\
&= A^{1/2} \left(A^{-1/2} (B - A) A^{-1/2} \right)^2 \left(A^{-1/2} (B + A) A^{-1/2} \right)^{-1} A^{1/2} \\
&= A^{1/2} A^{-1/2} (B - A) A^{-1/2} A^{-1/2} (B - A) A^{-1/2} A^{1/2} (B + A)^{-1} A^{1/2} A^{1/2} \\
&= A (B - A) A^{-1} (B - A) (B + A)^{-1} A,
\end{aligned}$$

then by (2.10) we get the desired result (2.1). \square

Remark 1. *Since, as above,*

$$2(A\nabla B - A!B) = A(B - A)A^{-1}(B - A)(B + A)^{-1}A$$

then (2.1) can be written as

$$(2.11) \quad 2r(A\nabla B - A!B) \leq A\nabla_{\nu} B - A!_{\nu} B \leq 2R(A\nabla B - A!B)$$

The first inequality in (2.11) was obtained in [10].

We observe that, if $\nu = \frac{1}{2}$, (2.1) becomes equality.

When some boundedness conditions are known, then we have the following result as well.

Theorem 2. *Let A, B be positive invertible operators and $M > m > 0$ such that*

$$(2.12) \quad MA \geq B \geq mA.$$

Then for any $\nu \in [0, 1]$ we have

$$(2.13) \quad rk(m, M)A \leq A\nabla_{\nu} B - A!_{\nu} B \leq RK(m, M)A$$

where $r = \min\{\nu, 1 - \nu\}$, $R = \max\{\nu, 1 - \nu\}$ and the bounds $K(m, M)$ and $k(m, M)$ are given by

$$(2.14) \quad K(m, M) := \begin{cases} (m - 1)^2 (m + 1)^{-1} & \text{if } M < 1, \\ \max\left\{(m - 1)^2 (m + 1)^{-1}, (M - 1)^2 (M + 1)^{-1}\right\} & \text{if } m \leq 1 \leq M, \\ (M - 1)^2 (M + 1)^{-1} & \text{if } 1 < m \end{cases}$$

and

$$(2.15) \quad k(m, M) := \begin{cases} (M-1)^2(M+1)^{-1} & \text{if } M < 1, \\ 0 & \text{if } m \leq 1 \leq M, \\ (m-1)^2(m+1)^{-1} & \text{if } 1 < m. \end{cases}$$

In particular,

$$(2.16) \quad \frac{1}{2}k(m, M)A \leq A\nabla B - A!B \leq \frac{1}{2}K(m, M)A.$$

Proof. As in the proof of Theorem 1 we have

$$(2.17) \quad r\varphi(t) \leq \nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq R\varphi(t)$$

for any $t > 0$, where $\varphi(t) = (t-1)^2(t+1)^{-1}$.

If we take the derivative of φ , we have

$$\begin{aligned} \varphi'(t) &= 2(t-1)(t+1)^{-1} - (t+1)^{-2}(t-1)^2 \\ &= (t-1)(t+1)^{-2}[2(t+1) - (t-1)] \\ &= (t-1)(t+1)^{-2}(2t+3) \end{aligned}$$

for any $t > 0$.

We observe that the function φ is decreasing on $(0, 1)$ and increasing on $(1, \infty)$. We have $\varphi(0) = 1$, $\varphi(1) = 0$ and $\lim_{t \rightarrow \infty} \varphi(t) = \infty$.

Using the properties of the function φ we have

$$\max_{t \in [m, M]} \varphi(t) = \begin{cases} \varphi(m) & \text{if } M < 1, \\ \max\{\varphi(m), \varphi(M)\} & \text{if } m \leq 1 \leq M, \\ \varphi(M) & \text{if } 1 < m, \end{cases} = K(m, M)$$

and

$$\min_{t \in [m, M]} \varphi(t) = \begin{cases} \varphi(M) & \text{if } M < 1, \\ 0 & \text{if } m \leq 1 \leq M, \\ \varphi(m) & \text{if } 1 < m, \end{cases} = k(m, M).$$

From (2.17) we have

$$(2.18) \quad rk(m, M) \leq \nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq RK(m, M)$$

for all $t \in [m, M]$.

If we use the continuous functional calculus for the positive invertible operator X with $mI \leq X \leq MI$, then we have

$$(2.19) \quad rk(m, M)I \leq \nu X + (1 - \nu)I - (\nu X^{-1} + (1 - \nu)I)^{-1} \leq RK(m, M)I.$$

If we multiply (2.12) both sides by $A^{-1/2}$ we get $MI \geq A^{-1/2}BA^{-1/2} \geq mI$.

By writing the inequality (2.19) for $X = A^{-1/2}BA^{-1/2}$ we obtain

$$(2.20) \quad rk(m, M)I \\ \leq \nu A^{-1/2}BA^{-1/2} + (1-\nu)I - \left(\nu \left(A^{-1/2}BA^{-1/2} \right)^{-1} + (1-\nu)I \right)^{-1} \\ \leq RK(m, M)I.$$

Finally, if we multiply both sides of (2.20) by $A^{1/2}$ we get the desired result (2.13). \square

Remark 2. Since $\varphi(t) \in [0, 1]$ for $t \in [0, 1]$, then $B \leq A$ implies that

$$(0 \leq) A\nabla_{\nu}B - A!_{\nu}B \leq RA$$

for any $\nu \in [0, 1]$. In particular,

$$(0 \leq) A\nabla B - A!B \leq \frac{1}{2}A.$$

We also have:

Theorem 3. Let A, B be positive invertible operators. Then for any $\nu \in [0, 1]$ we have

$$(2.21) \quad (0 \leq) A\nabla_{\nu}B - A!_{\nu}B \leq \nu(1-\nu)(B-A)A^{-1}(B-A)(B^{-1}+A^{-1})A \\ \leq \frac{1}{4}(B-A)A^{-1}(B-A)(B^{-1}+A^{-1})A.$$

Proof. In [1] we obtained the following reverse of Jensen's inequality:

$$(2.22) \quad 0 \leq (1-\nu)f(x) + \nu f(y) - f((1-\nu)x + \nu y) \\ \leq \nu(1-\nu)(y-x)[f'(y) - f'(x)].$$

for any $x, y \in \mathring{I}$ and $\nu \in [0, 1]$, provided the function $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable convex function on \mathring{I} , the interior of the interval I .

If we write the inequality (2.22) for the convex function $\Phi(x) = \frac{1}{x}$, $x > 0$, then we have

$$(2.23) \quad \frac{\nu}{y} + \frac{1-\nu}{x} - \frac{1}{\nu y + (1-\nu)x} \leq \nu(1-\nu)(y-x) \left(\frac{1}{x^2} - \frac{1}{y^2} \right)$$

for any $x, y > 0$.

If we take $y = \frac{1}{b}$ and $x = \frac{1}{a}$ with $a, b > 0$ in (2.23), then we get

$$\nu b + (1-\nu)a - (\nu b^{-1} + (1-\nu)a^{-1})^{-1} \leq \nu(1-\nu) \left(\frac{1}{b} - \frac{1}{a} \right) (a^2 - b^2)$$

namely,

$$(2.24) \quad \nu b + (1-\nu)a - (\nu b^{-1} + (1-\nu)a^{-1})^{-1} \leq \nu(1-\nu) \frac{a+b}{ab} (b-a)^2$$

for any $a, b > 0$ and $\nu \in [0, 1]$.

This is an inequality of interest in itself.

If we take $a = 1$ and $b = t$ in (2.24), then we get

$$\nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq \nu(1-\nu)(t-1)^2(1+t^{-1})$$

for any $t > 0$.

If we use the continuous functional calculus for the positive invertible operator X we get

$$(2.25) \quad \nu X + (1 - \nu) I - (\nu X^{-1} + (1 - \nu) I)^{-1} \leq \nu(1 - \nu) (X - I)^2 (X^{-1} + I).$$

If we write the inequality (2.8) for $X = A^{-1/2} B A^{-1/2}$, then we get

$$(2.26) \quad \nu A^{-1/2} B A^{-1/2} + (1 - \nu) I - \left(\nu \left(A^{-1/2} B A^{-1/2} \right)^{-1} + (1 - \nu) I \right)^{-1} \\ \leq \nu(1 - \nu) \left(A^{-1/2} B A^{-1/2} - I \right)^2 \left(\left(A^{-1/2} B A^{-1/2} \right)^{-1} + I \right),$$

and $\nu \in [0, 1]$.

If we multiply the inequality (2.9) both sides with $A^{1/2}$, then we get

$$(2.27) \quad A \nabla_{\nu} B - A!_{\nu} B \\ \leq \nu(1 - \nu) A^{1/2} \left(A^{-1/2} B A^{-1/2} - I \right)^2 \left(\left(A^{-1/2} B A^{-1/2} \right)^{-1} + I \right) A^{1/2},$$

and since

$$A^{1/2} \left(A^{-1/2} B A^{-1/2} - I \right)^2 \left(\left(A^{-1/2} B A^{-1/2} \right)^{-1} + I \right) A^{1/2} \\ = A^{1/2} A^{-1/2} (B - A) A^{-1/2} A^{-1/2} (B - A) A^{-1/2} A^{1/2} (B^{-1} + A^{-1}) A^{1/2} A^{1/2} \\ = (B - A) A^{-1} (B - A) (B^{-1} + A^{-1}) A$$

hence from (2.27) we get the desired result (2.21).

The last part is obvious from the fact that $\nu(1 - \nu) \leq \frac{1}{4}$, $\nu \in [0, 1]$. \square

We also have:

Theorem 4. *Let A, B be positive invertible operators and $M > m > 0$ such that the condition (2.12) is valid. Then for any $\nu \in [0, 1]$ we have*

$$(2.28) \quad (0 \leq) A \nabla_{\nu} B - A!_{\nu} B \leq \nu(1 - \nu) L(m, M) A$$

where

$$(2.29) \quad L(m, M) \\ := \begin{cases} (m - 1)^2 (1 + m^{-1}) & \text{if } M < 1, \\ \max \left\{ (m - 1)^2 (1 + m^{-1}), (M - 1)^2 (1 + M^{-1}) \right\} & \text{if } m \leq 1 \leq M, \\ (M - 1)^2 (1 + M^{-1}) & \text{if } 1 < m. \end{cases}$$

In particular,

$$(2.30) \quad (0 \leq) A \nabla B - A! B \leq \frac{1}{4} L(m, M) A.$$

Proof. As in the proof of Theorem 3 we have

$$(2.31) \quad \nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq \nu(1 - \nu) \psi(t)$$

for any $t > 0$ and $\nu \in [0, 1]$, where $\psi(t) = (t - 1)^2 (1 + t^{-1})$.

If we take the derivative of ψ , we have

$$\begin{aligned}\psi'(t) &= 2(t-1)(1+t^{-1}) - (t-1)^2 t^{-2} \\ &= (t-1)(2+2t^{-1}-t^{-1}+t^{-2}) = (t-1)(2+t^{-1}+t^{-2})\end{aligned}$$

for any $t > 0$.

We observe that the function ψ is decreasing on $(0, 1)$ and increasing on $(1, \infty)$. We have $\lim_{t \rightarrow 0^+} \psi(t) = \infty$, $\varphi(1) = 0$ and $\lim_{t \rightarrow \infty} \varphi(t) = \infty$.

Using the properties of the function ψ we have

$$\max_{t \in [m, M]} \psi(t) = \begin{cases} \psi(m) & \text{if } M < 1, \\ \max\{\psi(m), \psi(M)\} & \text{if } m \leq 1 \leq M, \\ \psi(M) & \text{if } 1 < m, \end{cases} = L(m, M).$$

Therefore, by (2.31) we have

$$\nu t + 1 - \nu - (\nu t^{-1} + 1 - \nu)^{-1} \leq \nu(1 - \nu)L(m, M)$$

for all $t \in [m, M]$ and $\nu \in [0, 1]$.

By utilizing a similar argument to the one in the proof of Theorem 2 we deduce the desired result (2.29). \square

3. APPLICATIONS

For two positive invertible operators A, B and positive real numbers m, m', M, M' assume that one of the following conditions (i) $0 < mI \leq A \leq m'I < M'I \leq B \leq MI$ and (ii) $0 < mI \leq B \leq m'I < M'I \leq A \leq MI$, holds. Put $h := \frac{M}{m}$ and $h' := \frac{M'}{m'}$. We observe that $h, h' > 1$ and if either of the condition (i) or (ii) holds, then $h \geq h'$.

If (i) is valid, then we have

$$(3.1) \quad A < h'A = \frac{M'}{m'}A \leq B \leq \frac{M}{m}A = hA,$$

while, if (ii) is valid, then we have

$$(3.2) \quad \frac{1}{h}A \leq B \leq \frac{1}{h'}A < A.$$

Proposition 1. *Let A, B positive invertible operators and positive real numbers m, m', M, M' such that the condition (i) holds. Then for any $\nu \in [0, 1]$ we have*

$$(3.3) \quad r(h'-1)^2(h'+1)^{-1}A \leq A\nabla_\nu B - A!_\nu B \leq R(h-1)^2(h+1)^{-1}A,$$

where $r = \min\{\nu, 1-\nu\}$, $R = \max\{\nu, 1-\nu\}$ and

$$(3.4) \quad A\nabla_\nu B - A!_\nu B \leq \nu(1-\nu)(h-1)^2(1+h^{-1})A.$$

In particular, we have

$$(3.5) \quad \frac{1}{2}(h'-1)^2(h'+1)^{-1}A \leq A\nabla B - A!B \leq \frac{1}{2}(h-1)^2(h+1)^{-1}A,$$

and

$$(3.6) \quad A\nabla B - A!B \leq \frac{1}{4}(h-1)^2(1+h^{-1})A.$$

The proof follows by utilizing the inequality (3.1), Theorem 2 and Theorem 4.

Proposition 2. Let A, B positive invertible operators and positive real numbers m, m', M, M' such that the condition (ii) holds. Then for any $\nu \in [0, 1]$ we have

$$(3.7) \quad r(h' - 1)^2 (h' + 1)^{-1} (h')^{-1} A \leq A\nabla_\nu B - A!_\nu B \leq R(h - 1)^2 (h + 1)^{-1} h^{-1} A,$$

and

$$(3.8) \quad A\nabla_\nu B - A!_\nu B \leq \nu(1 - \nu)(h - 1)^2 (1 + h^{-1}) h^{-1} A.$$

In particular, we have

$$(3.9) \quad \frac{1}{2} (h' - 1)^2 (h' + 1)^{-1} (h')^{-1} A \leq A\nabla B - A!B \leq \frac{1}{2} (h - 1)^2 (h + 1)^{-1} h^{-1} A,$$

and

$$(3.10) \quad A\nabla B - A!B \leq \frac{1}{4} (h - 1)^2 (1 + h^{-1}) h^{-1} A.$$

The proof follows by utilizing the inequality (3.2), Theorem 2 and Theorem 4.

If we consider the function $D(x, y) : [1, 10] \times [0, 1] \rightarrow \mathbb{R}$,

$$D(x, y) = y(1 - y)(1 + x^{-1}) - \max\{y, 1 - y\}(x + 1)^{-1}$$

then the plot of this function in Figure 1 shows that it take both positive and negative values, meaning that some time the upper bound for the quantity $A\nabla_\nu B - A!_\nu B$ provided by (3.3) is better and other time worse than the one from (3.7).




FIGURE 1. Plot of the difference function $D(x, y)$

REFERENCES

- [1] S. S. Dragomir, A note on Young's inequality, Preprint *RGMA Res. Rep. Coll.* **18** (2015), Art. 126. [Online <http://rgmia.org/papers/v18/v18a126.pdf>].
- [2] S. S. Dragomir, Some new reverses of Young's operator inequality, Preprint *RGMA Res. Rep. Coll.* **18** (2015), Art. 130. [<http://rgmia.org/papers/v18/v18a130.pdf>].
- [3] S. S. Dragomir, On new refinements and reverses of Young's operator inequality, Preprint *RGMA Res. Rep. Coll.* **18** (2015), Art. 135. [<http://rgmia.org/papers/v18/v18a135.pdf>].

- [4] S. S. Dragomir, Some inequalities for operator weighted geometric mean, Preprint *RGMA Res. Rep. Coll.* **18** (2015), Art. 139. [<http://rgmia.org/papers/v18/v18a139.pdf>].
- [5] S. S. Dragomir, Some reverses and a refinement of Hölder operator inequality, Preprint *RGMA Res. Rep. Coll.* **18** (2015), Art. 147. [<http://rgmia.org/papers/v18/v18a147.pdf>].
- [6] S. S. Dragomir, Bounds for the normalized Jensen functional, *Bull. Austral. Math. Soc.* **74** (3) (2006), 417-478.
- [7] S. Furuichi, Refined Young inequalities with Specht's ratio, *J. Egyptian Math. Soc.* **20** (2012), 46-49.
- [8] S. Furuichi, On refined Young inequalities and reverse inequalities, *J. Math. Inequal.* **5** (2011), 21-31.
- [9] W. Liao, J. Wu and J. Zhao, New versions of reverse Young and Heinz mean inequalities with the Kantorovich constant, *Taiwanese J. Math.* **19** (2015), No. 2, pp. 467-479.
- [10] G. Zuo, G. Shi and M. Fujii, Refined Young inequality with Kantorovich constant, *J. Math. Inequal.*, **5** (2011), 551-556.

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: `sever.dragomir@vu.edu.au`

URL: <http://rgmia.org/dragomir>

²SCHOOL OF COMPUTER SCIENCE & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA