

# Self Adjoint Operator Korovkin type and polynomial direct Approximations with rates

George A. Anastassiou  
Department of Mathematical Sciences  
University of Memphis  
Memphis, TN 38152, U.S.A.  
ganastss@memphis.edu

## Abstract

Here we present self adjoint operator Korovkin type theorems, via self adjoint operator Shisha-Mond type inequalities, also we give self adjoint operator polynomial approximations. This is a quantitative treatment to determine the degree of self adjoint operator uniform approximation with rates, of sequences of self adjoint operator positive linear operators. The same kind of work is performed over important operator polynomial sequences. Our approach is direct based on Gelfand isometry.

**2010 AMS Subject Classification:** 41A17, 41A25, 41A36, 41A80, 47A58, 47A60, 47A67.

**Key Words and Phrases:** Self adjoint operator, Hilbert space, Korovkin theory, Shisha-Mond inequality, positive linear operator, Bernstein type polynomials.

## 1 Background

Let  $A$  be a selfadjoint linear operator on a complex Hilbert space  $(H; \langle \cdot, \cdot \rangle)$ . The Gelfand map establishes a  $*$ -isometrically isomorphism  $\Phi$  between the set  $C(Sp(A))$  of all continuous functions defined on the spectrum of  $A$ , denoted  $Sp(A)$ , and the  $C^*$ -algebra  $C^*(A)$  generated by  $A$  and the identity operator  $1_H$  on  $H$  as follows (see e.g. [6, p. 3]):

For any  $f, g \in C(Sp(A))$  and any  $\alpha, \beta \in \mathbb{C}$  we have

- (i)  $\Phi(\alpha f + \beta g) = \alpha \Phi(f) + \beta \Phi(g)$ ;
- (ii)  $\Phi(fg) = \Phi(f)\Phi(g)$  (the operation composition is on the right) and  $\Phi(\bar{f}) = (\Phi(f))^*$ ;
- (iii)  $\|\Phi(f)\| = \|f\| := \sup_{t \in Sp(A)} |f(t)|$ ;

(iv)  $\Phi(f_0) = 1_H$  and  $\Phi(f_1) = A$ , where  $f_0(t) = 1$  and  $f_1(t) = t$ , for  $t \in Sp(A)$ .

With this notation we define

$$f(A) := \Phi(f), \text{ for all } f \in C(Sp(A)),$$

and we call it the continuous functional calculus for a selfadjoint operator  $A$ .

If  $A$  is a selfadjoint operator and  $f$  is a real valued continuous function on  $Sp(A)$  then  $f(t) \geq 0$  for any  $t \in Sp(A)$  implies that  $f(A) \geq 0$ , i.e.  $f(A)$  is a positive operator on  $H$ . Moreover, if both  $f$  and  $g$  are real valued continuous functions on  $Sp(A)$  then the following important property holds:

(P)  $f(t) \geq g(t)$  for any  $t \in Sp(A)$ , implies that  $f(A) \geq g(A)$  in the operator order of  $B(H)$ .

Equivalently, we use (see [5], pp. 7-8):

Let  $U$  be a selfadjoint operator on the complex Hilbert space  $(H, \langle \cdot, \cdot \rangle)$  with the spectrum  $Sp(U)$  included in the interval  $[m, M]$  for some real numbers  $m < M$  and  $\{E_\lambda\}_\lambda$  be its spectral family.

Then for any continuous function  $f : [a, b] \rightarrow \mathbb{C}$ , where  $[m, M] \subset (a, b)$ , it is well known that we have the following spectral representation in terms of the Riemann-Stieljes integral:

$$\langle f(U)x, y \rangle = \int_{m-0}^M f(\lambda) d(\langle E_\lambda x, y \rangle),$$

for any  $x, y \in H$ . The function  $g_{x,y}(\lambda) := \langle E_\lambda x, y \rangle$  is of bounded variation on the interval  $[m, M]$ , and

$$g_{x,y}(m-0) = 0 \text{ and } g_{x,y}(M) = \langle x, y \rangle,$$

for any  $x, y \in H$ . Furthermore, it is known that  $g_x(\lambda) := \langle E_\lambda x, x \rangle$  is increasing and right continuous on  $[m, M]$ .

In this article we will be using a lot the formula

$$\langle f(U)x, x \rangle = \int_{m-0}^M f(\lambda) d(\langle E_\lambda x, x \rangle), \quad \forall x \in H.$$

As a symbol we can write

$$f(U) = \int_{m-0}^M f(\lambda) dE_\lambda.$$

Above,  $m = \min \{\lambda | \lambda \in Sp(U)\} := \min Sp(U)$ ,  $M = \max \{\lambda | \lambda \in Sp(U)\} := \max Sp(U)$ . The projections  $\{E_\lambda\}_{\lambda \in \mathbb{R}}$ , are called the spectral family of  $A$ , with the properties:

(a)  $E_\lambda \leq E_{\lambda'}$  for  $\lambda \leq \lambda'$ ;

(b)  $E_{m-0} = 0_H$  (zero operator),  $E_M = 1_H$  (identity operator) and  $E_{\lambda+0} = E_\lambda$  for all  $\lambda \in \mathbb{R}$ .

Furthermore

$$E_\lambda := \varphi_\lambda(U), \quad \forall \lambda \in \mathbb{R},$$

is a projection which reduces  $U$ , with

$$\varphi_\lambda(s) := \begin{cases} 1, & \text{for } -\infty < s \leq \lambda, \\ 0, & \text{for } \lambda < s < +\infty. \end{cases}$$

The spectral family  $\{E_\lambda\}_{\lambda \in \mathbb{R}}$  determines uniquely the self-adjoint operator  $U$  and vice versa.

For more on the topic see [9], pp. 256-266, and for more details see there pp. 157-266. See also [4].

Some more basics are given (we follow [5], pp. 1-5):

Let  $(H; \langle \cdot, \cdot \rangle)$  be a Hilbert space over  $\mathbb{C}$ . A bounded linear operator  $A$  defined on  $H$  is selfjoint, i.e.,  $A = A^*$ , iff  $\langle Ax, x \rangle \in \mathbb{R}, \forall x \in H$ , and if  $A$  is selfadjoint, then

$$\|A\| = \sup_{x \in H: \|x\|=1} |\langle Ax, x \rangle|.$$

Let  $A, B$  be selfadjoint operators on  $H$ . Then  $A \leq B$  iff  $\langle Ax, x \rangle \leq \langle Bx, x \rangle, \forall x \in H$ .

In particular,  $A$  is called positive if  $A \geq 0$ .

Denote by

$$\mathcal{P} := \left\{ \varphi(s) := \sum_{k=0}^n \alpha_k s^k \mid n \geq 0, \alpha_k \in \mathbb{C}, 0 \leq k \leq n \right\}.$$

If  $A \in \mathcal{B}(H)$  (the Banach algebra of all bounded linear operators defined on  $H$ , i.e. from  $H$  into itself) is selfadjoint, and  $\varphi(s) \in \mathcal{P}$  has real coefficients, then  $\varphi(A)$  is selfadjoint, and

$$\|\varphi(A)\| = \max \{ |\varphi(\lambda)|, \lambda \in Sp(A) \}.$$

If  $\varphi$  is any function defined on  $\mathbb{R}$  we define

$$\|\varphi\|_A := \sup \{ |\varphi(\lambda)|, \lambda \in Sp(A) \}.$$

If  $A$  is selfadjoint operator on Hilbert space  $H$  and  $\varphi$  is continuous and given that  $\varphi(A)$  is selfadjoint, then  $\|\varphi(A)\| = \|\varphi\|_A$ . And if  $\varphi$  is a continuous real valued function so it is  $|\varphi|$ , then  $\varphi(A)$  and  $|\varphi|(A) = |\varphi(A)|$  are selfadjoint operators (by [5], p. 4, Theorem 7).

Hence it holds

$$\|\|\varphi(A)\|\| = \|\|\varphi\|\|_A = \sup \{ \|\varphi(\lambda)\|, \lambda \in Sp(A) \}$$

$$= \sup \{ |\varphi(\lambda)|, \lambda \in Sp(A) \} = \|\varphi\|_A = \|\varphi(A)\|,$$

that is  $\|\varphi(A)\| = \|\varphi(A)\|$ .

For a selfadjoint operator  $A \in \mathcal{B}(H)$  which is positive, there exists a unique positive selfadjoint operator  $B := \sqrt{A} \in \mathcal{B}(H)$  such that  $B^2 = A$ , that is  $(\sqrt{A})^2 = A$ . We call  $B$  the square root of  $A$ .

Let  $A \in \mathcal{B}(H)$ , then  $A^*A$  is selfadjoint and positive. Define the "operator absolute value"  $|A| := \sqrt{A^*A}$ . If  $A = A^*$ , then  $|A| = \sqrt{A^2}$ .

For a continuous real valued function  $\varphi$  we observe the following:

$$|\varphi(A)| \text{ (the functional absolute value)} = \int_{m-0}^M |\varphi(\lambda)| dE_\lambda =$$

$$\int_{m-0}^M \sqrt{(\varphi(\lambda))^2} dE_\lambda = \sqrt{(\varphi(A))^2} = |\varphi(A)| \text{ (operator absolute value),}$$

where  $A$  is a selfadjoint operator.

That is we have

$$|\varphi(A)| \text{ (functional absolute value)} = |\varphi(A)| \text{ (operator absolute value).}$$

The next comes from [4], p. 3:

We say that a sequence  $\{A_n\}_{n=1}^\infty \subset \mathcal{B}(H)$  converges uniformly to  $A$  (convergence in norm), iff

$$\lim_{n \rightarrow \infty} \|A_n - A\| = 0,$$

and we denote it as  $\lim_{n \rightarrow \infty} A_n = A$ .

We will be using Hölder's-McCarthy, 1967 ([10]), inequality: Let  $A$  be a selfadjoint positive operator on a Hilbert space  $H$ . Then

$$\langle A^r x, x \rangle \leq \langle Ax, x \rangle^r,$$

for all  $0 < r < 1$  and  $x \in H : \|x\| = 1$ .

Let  $A, B \in \mathcal{B}(H)$ , then

$$\|AB\| \leq \|A\| \|B\|,$$

by Banach algebra property.

## 2 Main Results

Here we derive self adjoint operator-Korovkin type theorems via operator-Shisha-Mond type inequalities. This is a quantitative approach, studying the degree of operator-uniform approximation with rates of sequences of operator-positive linear operators in the operator order of  $\mathcal{B}(H)$ . We continue similarly with

important polynomial operators. Our approach is direct based on Gelfand's isometry.

All the functions we are dealing here are real valued. We assume that  $Sp(A) \subseteq [m, M]$ .

Let  $\{L_n\}_{n \in \mathbb{N}}$  be a sequence of positive linear operators from  $C([m, M])$  into itself (i.e. if  $f, g \in C([m, M])$  such that  $f \geq g$ , then  $L_n(f) \geq L_n(g)$ ). It is interesting to study the convergence of  $L_n \rightarrow I$  (unit operator, i.e.  $I(f) = f, \forall f \in C([m, M])$ ). By property (i) we have that

$$\Phi(L_n f - f) = \Phi(L_n f) - \Phi(f) = (L_n f)(A) - f(A), \quad (1)$$

and

$$\Phi(L_n 1 \pm 1) = \Phi(L_n 1) \pm \Phi(1) = (L_n 1)(A) \pm 1_H, \quad (2)$$

the last comes by property (iv).

And by property (iii) we obtain

$$\|\Phi(L_n f - f)\| = \|(L_n f)(A) - f(A)\| = \|L_n f - f\|, \quad (3)$$

and

$$\|\Phi(L_n 1 \pm 1)\| = \|(L_n 1)(A) \pm 1_H\| = \|L_n(1) \pm 1\|. \quad (4)$$

We need

**Theorem 1** (*Shisha and Mond ([12]), 1968*) Let  $\{L_n\}_{n \in \mathbb{N}}$  be a sequence of positive linear operators from  $C([m, M])$  into itself. For  $n = 1, 2, \dots$ , suppose  $L_n(1)$  is bounded. Let  $f \in C([m, M])$ . Then for  $n = 1, 2, \dots$ , we have

$$\|L_n f - f\|_\infty \leq \|f\|_\infty \|L_n 1 - 1\|_\infty + \|L_n(1) + 1\|_\infty \omega_1(f, \mu_n), \quad (5)$$

where

$$\mu_n := \left\| L_n \left( (t - x)^2 \right) (x) \right\|_\infty^{\frac{1}{2}}, \quad (6)$$

with

$$\omega_1(f, \delta) := \sup_{\substack{x, y \in [m, M] \\ |x - y| \leq \delta}} |f(x) - f(y)|, \quad \delta > 0, \quad (7)$$

and  $\|\cdot\|_\infty$  stands for the sup-norm over  $[m, M]$ .

In particular, if  $L_n(1) = 1$ , then (5) becomes

$$\|L_n(f) - f\|_\infty \leq 2\omega_1(f, \mu_n). \quad (8)$$

**Note:** (i) In forming  $\mu_n^2$ ,  $x$  is kept fixed, however  $t$  forms the functions  $t, t^2$  on which  $L_n$  acts.

(ii) One can easily find, for  $n = 1, 2, \dots$ ,

$$\mu_n^2 \leq \|(L_n(t^2))(x) - x^2\|_\infty + 2c \|(L_n(t))(x) - x\|_\infty + c^2 \|(L_n(1))(x) - 1\|_\infty, \quad (9)$$

where

$$c := \max(|m|, |M|).$$

So, if the Korovkin's assumptions are fulfilled, i.e. if  $L_n(id^2) \xrightarrow{u} id^2$ ,  $L_n(id) \xrightarrow{u} id$  and  $L_n(1) \xrightarrow{u} 1$ , as  $n \rightarrow \infty$ ,  $id$  is the identity map and  $u$  is the uniform convergence, then  $\mu_n \rightarrow 0$ , and then  $\omega_1(f, \mu_n) \rightarrow 0$ , as  $n \rightarrow +\infty$ , and we obtain from (5) that  $\|L_n f - f\|_\infty \rightarrow 0$ , i.e.  $L_n(f) \xrightarrow{u} f$ , as  $n \rightarrow \infty$ ,  $\forall f \in C([m, M])$ .

We give

**Theorem 2** *All as in Theorem 1. Then*

$$\|(L_n f)(A) - f(A)\| \leq \|f(A)\| \|(L_n 1)(A) - 1_H\| + \|(L_n(1))(A) + 1_H\| \omega_1(f, \mu_n), \quad (10)$$

where

$$\mu_n := \left\| L_n \left( (t - A)^2 \right) (A) \right\|^{\frac{1}{2}}. \quad (11)$$

In particular, if  $(L_n(1))(A) = 1_H$ , then

$$\|(L_n(f))(A) - f(A)\| \leq 2\omega_1(f, \mu_n). \quad (12)$$

Furthermore it holds

$$\mu_n^2 \leq \|(L_n(t^2))(A) - A^2\| + 2c \|(L_n(t))(A) - A\| + c^2 \|(L_n(1))(A) - 1_H\|. \quad (13)$$

So, if  $(L_n(t^2))(A) \rightarrow A^2$ ,  $(L_n(t))(A) \rightarrow A$ ,  $(L_n(1))(A) \rightarrow 1_H$ , uniformly, as  $n \rightarrow \infty$ , then by (13) and (10) we get  $(L_n(f))(A) \rightarrow f(A)$ , uniformly, as  $n \rightarrow \infty$ .

That is establishing the self adjoint operator Korovkin theorem with rates.

Next we follow [2], pp. 273-274.

**Theorem 3** *Let  $L_n : C([m, M]) \rightarrow C([m, M])$ ,  $n \in \mathbb{N}$ , be a sequence of positive linear operators,  $f \in C([m, M])$ ,  $g \in C([m, M])$  and it is an  $(1 - 1)$  function.*

*Assume  $\{L_n(1)\}_{n \in \mathbb{N}}$  is uniformly bounded. Then*

$$\|L_n(f) - f\| \leq \|f\| \|L_n(1) - 1\| + (1 + \|L_n(1)\|) \omega_g(f, \rho_n), \quad (14)$$

where

$$\omega_g(f, h) := \sup_{x, y} \{|f(x) - f(y)| : |g(x) - g(y)| \leq h\}, \quad (15)$$

$h > 0$ , with

$$\rho_n := \left( \left\| L_n \left( (g - g(y))^2 \right) (y) \right\| \right)^{\frac{1}{2}}. \quad (16)$$

Here  $\|\cdot\|$  stands for the supremum norm. If  $L_n(1) = 1$ , then (14) simplifies to

$$\|L_n(f) - f\| \leq 2\omega_g(f, \rho_n). \quad (17)$$

We also have that

$$\rho_n^2 \leq \|L_n(g^2) - g^2\| + 2\|g\| \|L_n(g) - g\| + \|g\|^2 \|L_n(1) - 1\|. \quad (18)$$

If  $L_n(1) \xrightarrow{u} 1$ ,  $L_n(g) \xrightarrow{u} g$ ,  $L_n(g^2) \xrightarrow{u} g^2$ , then  $\omega_g(f, \rho_n) \rightarrow 0$ , and then  $L_n(f) \xrightarrow{u} f$ , as  $n \rightarrow +\infty, \forall f \in C([m, M])$ , where  $u$  stands for uniform convergence, so we get a generalization of Korovkin theorem quantitatively, and clearly by  $L_n(1) \xrightarrow{u} 1$ , we get  $\|L_n(1)\| \leq K, \forall n \in \mathbb{N}$ , where  $K > 0$ .

We present

**Theorem 4** All as in Theorem 3. Then

$$\begin{aligned} & \|(L_n(f))(A) - f(A)\| \leq \\ & \|f(A)\| \|(L_n(1))(A) - 1_H\| + (1 + \|(L_n(1))(A)\|) \omega_g(f, \rho_n), \end{aligned} \quad (19)$$

with

$$\rho_n := \left( \left\| L_n \left( (g - g(A))^2 \right) (A) \right\| \right)^{\frac{1}{2}}. \quad (20)$$

If  $(L_n(1))(A) = 1_H$ , then

$$\|(L_n(f))(A) - f(A)\| \leq 2\omega_g(f, \rho_n). \quad (21)$$

It holds

$$\begin{aligned} & \rho_n^2 \leq \|(L_n(g^2))(A) - g^2(A)\| + \\ & 2\|g(A)\| \|(L_n(g))(A) - A\| + \|g(A)\|^2 \|(L_n(1))(A) - 1_H\|. \end{aligned} \quad (22)$$

If  $(L_n(1))(A) \rightarrow 1_H$ ,  $(L_n(g))(A) \rightarrow A$ ,  $(L_n(g^2))(A) \rightarrow g^2(A)$ , uniformly, as  $n \rightarrow +\infty$ , then  $(L_n(f))(A) \rightarrow f(A)$ , uniformly, as  $n \rightarrow +\infty$ .

We make

**Remark 5** Next we consider the general Bernstein positive linear polynomial operators from  $C([m, M])$  into itself, for  $f \in C([m, M])$  we define

$$(B_N f)(s) = \sum_{i=0}^N \binom{N}{i} f\left(m + i \left(\frac{M-m}{N}\right)\right) \left(\frac{s-m}{M-m}\right)^i \left(\frac{M-s}{M-m}\right)^{N-i}, \quad (23)$$

$\forall s \in [m, M]$ , see [13], p. 80.

Then by [13], p. 81, we get that

$$\|B_N f - f\|_\infty \leq \frac{5}{4} \omega_1 \left( f, \frac{M-m}{\sqrt{N}} \right), \quad (24)$$

$\forall N \in \mathbb{N}$ , i.e.  $B_N f \xrightarrow{u} f$ , as  $N \rightarrow +\infty$ ,  $\forall f \in C([m, M])$ , the convergence is given with rates.

We clearly have that

$$\|(B_N f)(A) - f(A)\| \leq \frac{5}{4} \omega_1 \left( f, \frac{M-m}{\sqrt{N}} \right), \quad (25)$$

$\forall N \in \mathbb{N}$ , i.e.  $(B_N f)(A) \rightarrow f(A)$ , uniformly, as  $N \rightarrow +\infty$ .

We need

**Notation 6** Let  $x \in [m, M]$ . Denote

$$c(x) := \max(x-m, M-x) = \frac{1}{2} [M-m + |M+m-2x|] > 0. \quad (26)$$

Let  $h > 0$  be fixed,  $n \in \mathbb{N}$ . Define (see [1], p. 210)

$$\Phi_{*n}(x) := \left( \frac{|x|^{n+1}}{(n+1)!h} + \frac{|x|^n}{2n!} + \frac{h|x|^{n-1}}{8(n-1)!} \right). \quad (27)$$

We need

**Theorem 7** ([1], p. 219) Let  $\{L_N\}_{N \in \mathbb{N}}$  be a sequence of positive linear operators from  $C([m, M])$  into itself,  $x \in [m, M]$ ,  $f \in C^n([m, M])$ .

Here  $c(x)$ ,  $\Phi_{*n}(x)$  as in Notation 6. Assume that  $\omega_1(f^{(n)}, h) \leq w$ , where  $w, h$  are fixed positive numbers,  $0 < h < M-m$ . Then

$$\begin{aligned} |(L_N(f))(x) - f(x)| &\leq |f(x)| |(L_N(1))(x) - 1| + \\ &\sum_{k=1}^n \frac{|f^{(k)}(x)|}{k!} \left| (L_N((t-x)^k))(x) \right| + \frac{w \Phi_{*n}(c(x))}{(c(x))^n} (L_N(|t-x|^n))(x). \end{aligned} \quad (28)$$

Inequality (28) is sharp, for details see [1], p. 220.

Clearly all functions involved in (28) are continuous, see also [3], i.e. both sides of (28) are continuous functions.

Using properties (P) and (ii) and (28) we derive

**Theorem 8** All as in Theorem 7. Then

$$\begin{aligned} |(L_N(f))(A) - f(A)| &\leq |f(A)| |(L_N(1))(A) - 1_H| + \\ &\sum_{k=1}^n \frac{|f^{(k)}(A)|}{k!} \left| (L_N((t-A)^k))(A) \right| + \frac{w \Phi_{*n}(c(A))}{(c(A))^n} (L_N(|t-A|^n))(A). \end{aligned} \quad (29)$$



**Remark 9** *Inequality (29) implies*

$$\begin{aligned} & \| (L_N(f))(A) - f(A) \| \leq \| f(A) \| \| (L_N(1))(A) - 1_H \| + \\ & \sum_{k=1}^n \frac{\| f^{(k)}(A) \|}{k!} \left\| \left( L_N \left( (t-A)^k \right) \right) (A) \right\| + w \left\| \frac{\Phi_{*n}(c(A))}{(c(A))^n} \right\| \| (L_N(|t-A|^n))(A) \|. \end{aligned} \quad (30)$$

**Remark 10** *(to Theorem 8 and (30)) Assume further*

$$\| L_N(1) \|_\infty \leq \mu, \quad \forall N \in \mathbb{N}; \quad \mu > 0. \quad (31)$$

*By Riesz representation theorem, for each  $s \in [m, M]$ , there exists a positive finite measure  $\mu_s$  on  $[m, M]$  such that*

$$(L_N(f))(s) = \int_{[m, M]} f(t) d\mu_{sN}(t), \quad \forall f \in C([m, M]). \quad (32)$$

*Therefore ( $k = 1, \dots, n-1$ )*

$$\left| (L_N(\cdot - s)^k)(s) \right| = \left| \int_{[m, M]} (\lambda - s)^k d\mu_{sN}(\lambda) \right| \leq \int_{[m, M]} |\lambda - s|^k d\mu_{sN}(\lambda)$$

*(by Hölder's inequality)*

$$\begin{aligned} & \leq \left( \int_{[m, M]} 1 d\mu_{sN}(\lambda) \right)^{\frac{n-k}{n}} \left( \int_{[m, M]} |\lambda - s|^n d\mu_{sN}(\lambda) \right)^{\frac{k}{n}} \\ & = ((L_N(1))(s))^{\frac{n-k}{n}} ((L_N(|\cdot - s|^n))(s))^{\frac{k}{n}} \leq \mu^{\frac{n-k}{n}} ((L_N(|\cdot - s|^n))(s))^{\frac{k}{n}}. \end{aligned} \quad (33)$$

*That is*

$$\left| (L_N(\cdot - s)^k)(s) \right| \leq \mu^{\frac{n-k}{n}} ((L_N(|\cdot - s|^n))(s))^{\frac{k}{n}}, \quad (34)$$

$k = 1, \dots, n-1$ .

*Of course it holds*

$$|(L_N(\cdot - s)^n)(s)| \leq (L_N|\cdot - s|^n)(s). \quad (35)$$

*By property (P) we obtain*

$$\left| (L_N(\cdot - A)^k)(A) \right| \leq \mu^{\frac{n-k}{n}} ((L_N(|\cdot - A|^n))(A))^{\frac{k}{n}}, \quad (36)$$

*for  $k = 1, \dots, n-1$ , and*

$$|(L_N(\cdot - A)^n)(A)| \leq (L_N|\cdot - A|^n)(A). \quad (37)$$

Therefore

$$\begin{aligned} \left\| \left( L_N (\cdot - A)^k \right) (A) \right\| &\leq \mu^{\frac{n-k}{n}} \left\| \left( L_N (|\cdot - A|^n) \right) (A) \right\|^{\frac{k}{n}} \\ &= \mu^{\frac{n-k}{n}} \sup_{x \in H: \|x\|=1} \left\langle \left( L_N (|\cdot - A|^n) \right) (A) \right\rangle^{\frac{k}{n}} x, x \end{aligned} \quad (38)$$

(by Hölder's-Mc Carthy inequality)

$$\begin{aligned} &\leq \mu^{\frac{n-k}{n}} \sup_{x \in H: \|x\|=1} \left\langle \left( L_N (|\cdot - A|^n) \right) (A) x, x \right\rangle^{\frac{k}{n}} \\ &= \mu^{\frac{n-k}{n}} \left( \sup_{x \in H: \|x\|=1} \left\langle \left( L_N (|\cdot - A|^n) \right) (A) x, x \right\rangle \right)^{\frac{k}{n}} \\ &= \mu^{\frac{n-k}{n}} \left\| L_N (|\cdot - A|^n) (A) \right\|^{\frac{k}{n}}. \end{aligned} \quad (39)$$

Therefore it holds

$$\left\| \left( L_N (t - A)^k \right) (A) \right\| \leq \mu^{\frac{n-k}{n}} \left\| L_N (|t - A|^n) (A) \right\|^{\frac{k}{n}}, \quad (40)$$

$k = 1, \dots, n-1$ , and of course

$$\left\| L_N (t - A)^n (A) \right\| \leq \left\| L_N (|t - A|^n) (A) \right\|. \quad (41)$$

Based on (40) and (41) and by assuming that  $(L_n(1))(A) \rightarrow 1_H$  and  $(L_N(|t - A|^n))(A) \rightarrow 0_H$ , uniformly, as  $N \rightarrow +\infty$ , we obtain by (30) that  $(L_N(f))(A) \rightarrow f(A)$ , uniformly as  $N \rightarrow +\infty$ .

We mention

**Theorem 11** ([1], p. 230) For any  $f \in C^1([0, 1])$  consider the Bernstein polynomials

$$(\beta_n(f))(t) := \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} t^k (1-t)^{n-k}, \quad t \in [0, 1].$$

Then

$$\|(\beta_n f) - f\|_\infty \leq \frac{0.78125}{\sqrt{n}} \omega_1\left(f', \frac{1}{4\sqrt{n}}\right). \quad (42)$$

We make

**Remark 12** The map

$$[m, M] \ni s = \varphi(t) = (M - m)t + m, \quad t \in [0, 1], \quad (43)$$

maps  $(1 - 1)$  and onto,  $[0, 1]$  onto  $[m, M]$ .

Let  $f \in C^1([m, M])$ , then

$$f(s) = f(\varphi(t)) = f((M-m)t + m), \quad (44)$$

and

$$\frac{df(s)}{dt} = (f(\varphi(t)))' = f'(\varphi(t))(M-m) = f'(s)(M-m). \quad (45)$$

By (42) we get that

$$\begin{aligned} & \|\beta_n(f((M-m)t + m) - f((M-m)t + m))\|_{\infty, [0,1]} \leq \\ & \frac{0.78125}{\sqrt{n}} \omega_1\left(f'(s)(M-m), \frac{1}{4\sqrt{n}}\right) = \frac{0.78125}{\sqrt{n}} (M-m) \omega_1\left(f'(s), \frac{1}{4\sqrt{n}}\right). \end{aligned} \quad (46)$$

However we have

$$\omega_1\left(f'(s), \frac{1}{4\sqrt{n}}\right) = \omega_1\left(f'((M-m)t + m), \frac{1}{4\sqrt{n}}\right) = \quad (47)$$

$$\sup_{\substack{t_1, t_2 \in [0,1] \\ |t_1 - t_2| \leq \frac{1}{4\sqrt{n}}}} |f'((M-m)t_1 + m) - f'((M-m)t_2 + m)| =$$

$$\sup_{\substack{s_1, s_2 \in [m, M] \\ |s_1 - s_2| \leq \frac{M-m}{4\sqrt{n}}}} |f'(s_1) - f'(s_2)| = \omega_1\left(f', \frac{M-m}{4\sqrt{n}}\right), \quad (48)$$

above notice that

$$\begin{aligned} |s_1 - s_2| &= |((M-m)t_1 + m) - ((M-m)t_2 + m)| = \\ &= (M-m)|t_1 - t_2| \leq \frac{M-m}{4\sqrt{n}}. \end{aligned} \quad (49)$$

So we have proved that

$$\omega_1\left(f'(s), \frac{1}{4\sqrt{n}}\right) = \omega_1\left(f', \frac{M-m}{4\sqrt{n}}\right). \quad (50)$$

Finally, we observe that

$$\begin{aligned} & (\beta_n(f((M-m)t + m))(t)) = \\ & \sum_{k=0}^n \left( f\left((M-m)\frac{k}{n} + m\right) \right) \binom{n}{k} t^k (1-t)^{n-k} = \\ & \sum_{k=0}^n \left( f\left((M-m)\frac{k}{n} + m\right) \right) \binom{n}{k} \left(\frac{s-m}{M-m}\right)^k \left(\frac{M-s}{M-m}\right)^{n-k} =: (B_n(f))(s), \end{aligned} \quad (51)$$

$s \in [m, M]$ .

The operators  $(B_n(f))(s)$  are the general Bernstein polynomials.

From (46) and (50), we derive that

$$\|(B_n f)(s) - f(s)\|_{\infty, [m, M]} \leq \frac{0.78125}{\sqrt{n}} (M - m) \omega_1 \left( f', \frac{M - m}{4\sqrt{n}} \right). \quad (52)$$

Based on the above and the property (iii), we can give

**Theorem 13** Let  $f' \in [m, M]$ . Then

$$\|(B_n f)(A) - f(A)\| \leq \frac{0.78125 (M - m)}{\sqrt{n}} \omega_1 \left( f', \frac{M - m}{4\sqrt{n}} \right). \quad (53)$$

I.e.  $(B_n f)(A) \rightarrow A$ , uniformly, with rates as  $n \rightarrow +\infty$ .

We make

**Remark 14** Let  $f \in C([m, M])$ , then the function  $f((M - m)t + m)$  is a continuous function in  $t \in [0, 1]$ .

Let  $r \in \mathbb{N}$ , we evaluate the modulus of smoothness ( $\delta > 0$ )

$$\begin{aligned} \omega_r(f((M - m)t + m), \delta) &= \\ \sup_{0 \leq h \leq \delta} \left\| \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} f((M - m)(t + kh) + m) \right\|_{\infty, [0, 1 - rh]} &= \\ \sup_{0 \leq h^* \leq \delta(M - m)} \left\| \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} f(s + kh^*) \right\|_{s, \infty, [m, M - rh^*]} & \\ (h^* = (M - m)h) & \\ &= \omega_r(f, (M - m)\delta), \end{aligned} \quad (54)$$

proving that

$$\omega_r(f((M - m)t + m), \delta) = \omega_r(f, (M - m)\delta), \quad (55)$$

for any  $r \in \mathbb{N}$ , and  $\delta > 0$ .

We need

**Theorem 15** ([11], p. 97) For  $f \in C([0, 1])$ ,  $n \in \mathbb{N}$ , we have

$$\|\beta_n(f) - f\| \leq \omega_2 \left( f, \frac{1}{\sqrt{n}} \right), \quad (56)$$

a sharp inequality.

We get

**Theorem 16** *Let  $f \in C([m, M])$ ,  $n \in \mathbb{N}$ . Then*

$$\|(B_n(f))(A) - f(A)\| = \|B_n(f) - f\|_\infty \leq \omega_2\left(f, \frac{M-m}{\sqrt{n}}\right). \quad (57)$$

We need

**Definition 17** ([11], p. 151) *Let  $f \in C([0, 1])$ ,  $n \in \mathbb{N}$ . We define the Durrmeyer type operators (the genuine Bernstein-Durrmeyer operators)*

$$\begin{aligned} (M_n^{-1,-1}(f))(x) &= f(0)(1-x)^n + f(1)x^n + \\ & (n-1) \sum_{k=1}^{n-1} p_{n,k}(x) \int_0^1 f(t) p_{n-2,k-1}(t) dt, \end{aligned} \quad (58)$$

where

$$p_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \quad n \in \mathbb{N}, \quad x \in [0, 1].$$

We will use

**Theorem 18** ([11], p. 155) *For  $f \in C([0, 1])$ ,  $n \in \mathbb{N}$ , we have*

$$\|M_n^{-1,-1}(f) - f\|_\infty \leq \frac{5}{4} \omega_2\left(f, \frac{1}{\sqrt{n+1}}\right). \quad (59)$$

We make

**Remark 19** *Let  $f \in C([m, M])$ , then  $f((M-m)t + m) \in C([0, 1])$ . Hence ( $s \in [m, M]$ ,  $t \in [0, 1]$ )*

$$\begin{aligned} (\overline{M}_n^{-1,-1}f)(s) &:= M_n^{-1,-1}(f((M-m)t + m))(t) \stackrel{(58)}{=} \\ & f(m) \left(\frac{M-s}{M-m}\right)^n + f(M) \left(\frac{s-m}{M-m}\right)^n + \\ & (n-1) \sum_{k=1}^{n-1} \binom{n}{k} \left(\frac{s-m}{M-m}\right)^k \left(\frac{M-s}{M-m}\right)^{n-k} \binom{n-2}{k-1} \int_m^M f(\overline{s}) \cdot \\ & \left(\frac{\overline{s}-m}{M-m}\right)^{k-1} \left(\frac{M-\overline{s}}{M-m}\right)^{n-k-1} d\overline{s}. \end{aligned} \quad (60)$$

We give

**Theorem 20** Let  $f \in C([m, M])$ ,  $n \in \mathbb{N}$ . Then

$$\left\| \left( \overline{M}_n^{-1, -1} f \right) (A) - f(A) \right\| = \left\| \overline{M}_n^{-1, -1} f - f \right\|_\infty \leq \frac{5}{4} \omega_2 \left( f, \frac{M-m}{\sqrt{n+1}} \right). \quad (61)$$

We need

**Definition 21** ([7]) For  $f \in C([0, 1])$ ,  $w \in \mathbb{N}$ , and  $0 \leq \beta \leq \gamma$ , we define the Stancu-type positive linear operators

$$\left( L_{w_0}^{(0\beta\gamma)} f \right) (x) = \sum_{k=0}^w f \left( \frac{k+\beta}{w+\gamma} \right) p_{w,k}(x), \quad x \in [0, 1], \quad (62)$$

$$p_{w,k}(x) = \binom{w}{k} x^k (1-x)^{w-k}.$$

We need

**Theorem 22** ([2], p. 516 and [7]) For  $\mathbb{N} \ni w > \lceil \gamma^2 \rceil$  ( $\lceil \cdot \rceil$  is the ceiling),  $f \in C([0, 1])$  we have:

$$\begin{aligned} \left\| L_{w_0}^{(0\beta\gamma)} f - f \right\|_\infty &\leq \left[ 3 + \frac{(w^3 + 4w\beta^2(w - \gamma^2))}{4(w - \gamma^2)(w + \gamma)^2} \right] \omega_2 \left( f, \frac{1}{\sqrt{w}} \right) \\ &\quad + \frac{2(\beta + \gamma)\sqrt{w}}{(w + \gamma)} \omega_1 \left( f, \frac{1}{\sqrt{w}} \right). \end{aligned} \quad (63)$$

We make

**Remark 23** Let  $f \in C([m, M])$ , then  $f((M-m)t + m) \in C([0, 1])$ . Hence ( $s \in [m, M]$ ,  $t \in [0, 1]$ )

$$\begin{aligned} \left( \overline{L}_{w_0}^{(0\beta\gamma)} f \right) (s) &:= L_{w_0}^{(0\beta\gamma)} (f((M-m)t + m))(t) \stackrel{(62)}{=} \\ &\sum_{k=0}^w f \left( (M-m) \left( \frac{k+\beta}{w+\gamma} \right) + m \right) \binom{w}{k} \left( \frac{s-m}{M-m} \right)^k \left( \frac{M-s}{M-m} \right)^{w-k}. \end{aligned} \quad (64)$$

We give

**Theorem 24** Let  $f \in C([m, M])$ ,  $w \in \mathbb{N}$ ,  $0 \leq \beta \leq \gamma$ . We take  $w > \lceil \gamma^2 \rceil$ . Then

$$\begin{aligned} \left\| \left( \overline{L}_{w_0}^{(0\beta\gamma)} f \right) (A) - f(A) \right\| &= \left\| \left( \overline{L}_{w_0}^{(0\beta\gamma)} f \right) - f \right\|_\infty \leq \\ &\left[ 3 + \frac{(w^3 + 4w\beta^2(w - \gamma^2))}{4(w - \gamma^2)(w + \gamma)^2} \right] \omega_2 \left( f, \frac{M-m}{\sqrt{w}} \right) + \frac{2(\beta + \gamma)\sqrt{w}}{(w + \gamma)} \omega_1 \left( f, \frac{M-m}{\sqrt{w}} \right). \end{aligned} \quad (65)$$

We make

**Remark 25** Next we assume that the spectrum of  $A$  is  $[0, 1]$ . For example, it could be  $Af = xf(x)$  on  $L^2([0, 1])$  which is a self adjoint operator and it has spectrum  $[0, 1]$ .

We need

**Definition 26** ([14]) Let  $f \in C([0, 1])$ , we define the special Stancu operator

$$S_n(f, x) = \frac{2(n!)}{(2n)!} \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} (nx)_k (n-nx)_{n-k}, \quad (66)$$

where  $(a)_0 = 1$ ,  $(a)_b = \sum_{k=0}^{b-1} (a-k)$ ,  $a \in \mathbb{R}$ ,  $b \in \mathbb{N}$ ,  $n \in \mathbb{N}$ ,  $x \in [0, 1]$ .

**Theorem 27** ([8], p. 75) Let  $f \in C([0, 1])$ ,  $n \in \mathbb{N}$ . Then

$$|((S_n - M_n^{-1,-1})(f))(x)| \leq c_1 \omega_4 \left( f, \sqrt[4]{\frac{3x(1-x)}{n(n+1)}} \right), \quad (67)$$

$\forall x \in [0, 1]$ , where  $c_1 > 0$  is an absolute constant independent of  $n$ ,  $f$  and  $x$ .

We obtain

**Theorem 28** Let  $f \in C([0, 1])$ ,  $n \in \mathbb{N}$ . Then

$$\|(S_n - M_n^{-1,-1})(A)\| = \|S_n - M_n^{-1,-1}\|_\infty \leq c_1 \omega_4 \left( f, \sqrt[4]{\frac{3}{4n(n+1)}} \right). \quad (68)$$

## References

- [1] G.A. Anastassiou, *Moments in probability and approximation theory*, Longman Scientific & Technical, Pitman Research Notes in Mathematics Series, No. 287, John Wiley & Sons, Inc., Essex, New York, 1993.
- [2] G.A. Anastassiou, *Intelligent Mathematics: Computational Analysis*, Springer, Heidelberg, New York, 2011.
- [3] G.A. Anastassiou, *Self adjoint operator Korovkin type quantitative approximations*, submitted for publication, 2016.
- [4] S.S. Dragomir, *Inequalities for functions of selfadjoint operators on Hilbert Spaces*, [ajmaa.org/RGMIA/monographs/InFuncOp.pdf](http://ajmaa.org/RGMIA/monographs/InFuncOp.pdf), 2011.

- [5] S. Dragomir, *Operator inequalities of Ostrowski and Trapezoidal type*, Springer, New York, 2012.
- [6] T. Furuta, J. Mićić Hot, J. Pečarić, Y. Seo, *Mond-Pečarić Method in Operator Inequalities. Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Element, Zagreb, 2005.
- [7] H.H. Gonska, J. Meier, *Quantitative theorems on approximation by Bernstein-Stancu operators*, Estratto da *Calcolo* 21 (fasc. IV), 1984, 317-335.
- [8] H. Gonska, P. Pitul, I. Rasa, *On Peano's form of the Taylor remainder, Voronovskaja's theorem and the commutator of positive linear operators*, In: O. Agratini, P. Blaga (eds.), *Proceedings of the International Conference on Numerical Analysis and Approximation theory*, Cluj-Napoca, Romania, July 5-8, 2006, Babes-Bolyai University, 55-80.
- [9] G. Helmberg, *Introduction to Spectral Theory in Hilbert Space*, John Wiley & Sons, Inc., New York, 1969.
- [10] C.A. McCarthy,  $c_p$ , *Israel J. Math.*, 5, 1967, 249-271.
- [11] R. Paltanea, *Approximation Theory using positive linear operators*, Birkhauser, Boston, 2004.
- [12] O. Shisha, B. Mond, *The degree of convergence of sequences of linear positive operators*, *Nat. Acad. of Sci. U.S.*, 60, 1968, 1196-1200.
- [13] L. Shumaker, *Spline functions basic theory*, Wiley-Interscience, New York, 1981.
- [14] D.D. Stancu, *Approximation of functions by a new class of linear polynomial operators*, *Rev. Roumaine Math. Pures Appl.* 13, 1968, 1173-1194.