

**HERMITE-HADAMARD TYPE INEQUALITIES OF SECOND
KIND FOR GENERALIZED RIEMANN-LIOUVILLE
FRACTIONAL INTEGRALS**

SILVESTRU SEVER DRAGOMIR^{1,2}

ABSTRACT. In this paper we establish some new Hermite-Hadamard type inequalities for the Generalized Riemann-Liouville fractional integrals $I_{a+,g}^\alpha f$ and $I_{b-,g}^\alpha f$, g , where g is a strictly increasing function on (a, b) , having a continuous derivative on (a, b) and under the assumption that the composite function $f \circ g^{-1}$ is convex on $(g(a), g(b))$. These provide some complementary results established in a previous paper. Some natural applications for Hadamard and Exponential fractional integrals are also provided.

1. INTRODUCTION

The following integral inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2},$$

which holds for any convex function $f : [a, b] \rightarrow \mathbb{R}$, is well known in the literature as the *Hermite-Hadamard inequality*.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the Theory of Special Means and in Information Theory for divergence measures, from which we would like to refer the reader to the monograph [11], the recent survey paper [9] and the references therein.

Let (a, b) with $-\infty \leq a < b \leq \infty$ be a finite or infinite interval of the real line \mathbb{R} and α a complex number with $\text{Re}(\alpha) > 0$. Also let g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . Following [14, p. 100], we introduce the *generalized left- and right-sided Riemann-Liouville fractional integrals* of a function f with respect to another function g on $[a, b]$ by

$$(1.1) \quad I_{a+,g}^\alpha f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \frac{g'(t) f(t) dt}{[g(x) - g(t)]^{1-\alpha}}, \quad a < x \leq b$$

and

$$(1.2) \quad I_{b-,g}^\alpha f(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \frac{g'(t) f(t) dt}{[g(t) - g(x)]^{1-\alpha}}, \quad a \leq x < b.$$

For $g(t) = t$ we have the classical *Riemann-Liouville fractional integrals*

$$(1.3) \quad J_{a+}^\alpha f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \frac{f(t) dt}{(x-t)^{1-\alpha}}, \quad a < x \leq b$$

1991 *Mathematics Subject Classification*. 26D15, 26D10, 26D07, 26A33.

Key words and phrases. Riemann-Liouville fractional integrals, Hadamard fractional integral, Hermite-Hadamard type inequalities, Convex functions, *GA*-convex functions.

and

$$(1.4) \quad J_{b-}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \frac{f(t) dt}{(t-x)^{1-\alpha}}, \quad a \leq x < b,$$

while for the logarithmic function $g(t) = \ln t$ we have the *Hadamard fractional integrals* [14, p. 111]

$$(1.5) \quad H_{a+}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \left[\ln \left(\frac{x}{t} \right) \right]^{\alpha-1} \frac{f(t) dt}{t}, \quad 0 \leq a < x \leq b$$

and

$$(1.6) \quad H_{b-}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \left[\ln \left(\frac{t}{x} \right) \right]^{\alpha-1} \frac{f(t) dt}{t}, \quad 0 \leq a < x < b.$$

Also, for $g(t) = \exp(\beta t)$, $\beta > 0$, we can consider the " *β -Exponential fractional integrals*"

$$(1.7) \quad E_{a+, \beta}^{\alpha} f(x) := \frac{\beta}{\Gamma(\alpha)} \int_a^x \frac{\exp(\beta t) f(t) dt}{[\exp(\beta x) - \exp(\beta t)]^{1-\alpha}}, \quad a < x \leq b$$

and

$$(1.8) \quad E_{b-, \beta}^{\alpha} f(x) := \frac{\beta}{\Gamma(\alpha)} \int_x^b \frac{\exp(\beta t) f(t) dt}{[\exp(\beta t) - \exp(\beta x)]^{1-\alpha}}, \quad a \leq x < b.$$

If g is a function which maps an interval I of the real line to the real numbers, and is both continuous and injective then we can define the *g -mean of two numbers* $a, b \in I$ as

$$M_g(a, b) := g^{-1} \left(\frac{g(a) + g(b)}{2} \right).$$

If $I = \mathbb{R}$ and $g(t) = t$ is the *identity function*, then $M_g(a, b) = A(a, b) := \frac{a+b}{2}$, the *arithmetic mean*. If $I = (0, \infty)$ and $g(t) = \ln t$, then $M_g(a, b) = G(a, b) := \sqrt{ab}$, the *geometric mean*. If $I = (0, \infty)$ and $g(t) = \frac{1}{t}$, then $M_g(a, b) = H(a, b) := \frac{2ab}{a+b}$, the *harmonic mean*. If $I = (0, \infty)$ and $g(t) = t^p$, $p \neq 0$, then $M_g(a, b) = M_p(a, b) := \left(\frac{a^p + b^p}{2} \right)^{1/p}$, the *power mean with exponent p* . Finally, if $I = \mathbb{R}$ and $g(t) = \exp t$, then

$$M_g(a, b) = LME(a, b) := \ln \left(\frac{\exp a + \exp b}{2} \right),$$

the *LogMeanExp function*.

In the recent paper [10] we obtained the following result:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function on $[a, b]$ and g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . If $f \circ g^{-1}$*

is convex on $(g(a), g(b))$, then for any $x \in (a, b)$ we have the inequalities

$$\begin{aligned}
(1.9) \quad & \frac{1}{\alpha} f \circ g^{-1} \left(\frac{\alpha}{\alpha+1} \frac{g(a) + g(b)}{2} + \frac{1}{\alpha+1} g(x) \right) \\
& \leq \int_0^1 s^{\alpha-1} f \circ g^{-1} \left(s \frac{g(a) + g(b)}{2} + (1-s) g(x) \right) ds \\
& \leq \frac{1}{2} \Gamma(\alpha) \left[\frac{I_{a+,g}^\alpha f(x)}{(g(x) - g(a))^\alpha} + \frac{I_{b-,g}^\alpha f(x)}{(g(b) - g(x))^\alpha} \right] \\
& \leq \frac{1}{\alpha+1} \left[\frac{f(a) + f(b)}{2} + \frac{1}{\alpha} f(x) \right].
\end{aligned}$$

In particular, we have the inequalities

$$\begin{aligned}
(1.10) \quad f(M_g(a, b)) & \leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(g(b) - g(a))^\alpha} [I_{a+,g}^\alpha f(M_g(a, b)) + I_{b-,g}^\alpha f(M_g(a, b))] \\
& \leq \frac{\alpha}{\alpha+1} \left[\frac{f(a) + f(b)}{2} + \frac{1}{\alpha} f(M_g(a, b)) \right] \leq \frac{f(a) + f(b)}{2}.
\end{aligned}$$

Motivated by the above results, in this paper we establish some Hermite-Hadamard type inequalities for the dual operator

$$\frac{1}{2} \Gamma(\alpha) \left[\frac{J_{x-,g}^\alpha f(a)}{(g(x) - g(a))^\alpha} + \frac{I_{x+,g}^\alpha f(b)}{(g(b) - g(x))^\alpha} \right], \quad x \in (a, b)$$

under the assumption that the composite function $f \circ g^{-1}$ is convex on $(g(a), g(b))$. We call these inequalities of second kind, to make a distinction between these and the inequalities (1.9) and (1.10). Some applications for Hadamard and Exponential fractional integrals are also provided.

2. THE MAIN RESULTS

We have the following result:

Theorem 2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function on $[a, b]$ and g be a strictly increasing function on (a, b) , having a continuous derivative g' on (a, b) . If $f \circ g^{-1}$ is convex on $(g(a), g(b))$, then for any $x \in (a, b)$ we have the inequalities*

$$\begin{aligned}
(2.1) \quad & \frac{1}{\alpha} f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha+1} + g(x) \frac{\alpha}{\alpha+1} \right) \\
& \leq \int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + s g(x) \right) ds \\
& \leq \frac{1}{2} \Gamma(\alpha) \left[\frac{I_{x+,g}^\alpha f(b)}{(g(b) - g(x))^\alpha} + \frac{I_{x-,g}^\alpha f(a)}{(g(x) - g(a))^\alpha} \right] \\
& \leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(x) \right].
\end{aligned}$$

Proof. Using the change of variable $u = g(t)$, then we have $du = g'(t) dt$, $t = g^{-1}(u)$ and

$$I_{a+,g}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_{g(a)}^{g(x)} \frac{f \circ g^{-1}(u) du}{[g(x) - u]^{1-\alpha}}$$

for $a < x \leq b$ and

$$I_{b-,g}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_{g(x)}^{g(b)} \frac{f \circ g^{-1}(u) du}{[u - g(x)]^{1-\alpha}}$$

for $a \leq x < b$.

Further, if we change the variable $u = (1-s)g(a) + sg(x)$, with $s \in [0, 1]$, then for $a < x \leq b$ we have

$$\begin{aligned} (2.2) \quad I_{a+,g}^\alpha f(x) &= \frac{1}{\Gamma(\alpha)} (g(x) - g(a))^\alpha \int_0^1 (1-s)^{\alpha-1} f \circ g^{-1}((1-s)g(a) + sg(x)) ds \\ &= \frac{1}{\Gamma(\alpha)} (g(x) - g(a))^\alpha \int_0^1 s^{\alpha-1} f \circ g^{-1}(sg(a) + (1-s)g(x)) ds \end{aligned}$$

where for the last equality we replaced s by $1-s$.

If we change the variable $u = (1-s)g(x) + sg(b)$, with $s \in [0, 1]$, then for $a \leq x < b$ we also have

$$(2.3) \quad I_{b-,g}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} (g(b) - g(x))^\alpha \int_0^1 s^{\alpha-1} f \circ g^{-1}((1-s)g(x) + sg(b)) ds.$$

Now, if we replace x with b in (2.2) and a with x , then we get

$$I_{x+,g}^\alpha f(b) = \frac{1}{\Gamma(\alpha)} (g(b) - g(x))^\alpha \int_0^1 s^{\alpha-1} f \circ g^{-1}(sg(x) + (1-s)g(b)) ds.$$

Also, if we replace x with a and b with x in (2.3), then we get

$$(2.4) \quad I_{x-,g}^\alpha f(a) = \frac{1}{\Gamma(\alpha)} (g(x) - g(a))^\alpha \int_0^1 s^{\alpha-1} f \circ g^{-1}((1-s)g(a) + sg(x)) ds.$$

By adding these two equalities, we get the following equality of interest

$$\begin{aligned} (2.5) \quad & \frac{1}{2} \Gamma(\alpha) \left[\frac{I_{x+,g}^\alpha f(b)}{(g(b) - g(x))^\alpha} + \frac{I_{x-,g}^\alpha f(a)}{(g(x) - g(a))^\alpha} \right] \\ &= \frac{1}{2} \int_0^1 s^{\alpha-1} [f \circ g^{-1}(sg(x) + (1-s)g(b)) + f \circ g^{-1}((1-s)g(a) + sg(x))] ds \end{aligned}$$

for any $x \in (a, b)$.

By the convexity of $f \circ g^{-1}$ on $(g(a), g(b))$ we get

$$\begin{aligned} & \frac{f \circ g^{-1}(sg(x) + (1-s)g(b)) ds + f \circ g^{-1}((1-s)g(a) + sg(x))}{2} \\ & \geq f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + sg(x) \right) \end{aligned}$$

for any $s \in [0, 1]$.

On multiplying this inequality by $s^{\alpha-1}$ and taking the integral over s , we get

$$\begin{aligned} & \int_0^1 s^{\alpha-1} \frac{f \circ g^{-1}(sg(x) + (1-s)g(b)) ds + f \circ g^{-1}((1-s)g(a) + sg(x))}{2} ds \\ & \geq \int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + sg(x) \right) ds, \end{aligned}$$

which proves the second inequality in (2.1).

By Jensen's inequality for the convex function $f \circ g^{-1}$ we also have

$$\begin{aligned} & \frac{\int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a)+g(b)}{2} + sg(x) \right) ds}{\int_0^1 s^{\alpha-1} ds} \\ & \geq f \circ g^{-1} \left(\frac{\int_0^1 s^{\alpha-1} \left((1-s) \frac{g(a)+g(b)}{2} + sg(x) \right) ds}{\int_0^1 s^{\alpha-1} ds} \right) \\ & = f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha+1} + g(x) \frac{\alpha}{\alpha+1} \right) \end{aligned}$$

giving that

$$\begin{aligned} & \int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + sg(x) \right) ds \\ & \geq \frac{1}{\alpha} f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha+1} + g(x) \frac{\alpha}{\alpha+1} \right), \end{aligned}$$

which proves the first inequality in (2.1).

By the convexity of $f \circ g^{-1}$ we also have

$$\begin{aligned} & \frac{f \circ g^{-1}(sg(x) + (1-s)g(b)) + f \circ g^{-1}((1-s)g(a) + sg(x))}{2} \\ & \leq \frac{1}{2} [sf \circ g^{-1}(g(x)) + (1-s)f \circ g^{-1}(g(b))] \\ & \quad + \frac{1}{2} [(1-s)f \circ g^{-1}(g(a)) + sf \circ g^{-1}(g(x))] \\ & = (1-s) \frac{f(a) + f(b)}{2} + sf(x) \end{aligned}$$

for any $x \in (a, b)$ and $s \in [0, 1]$.

On multiplying this inequality by $s^{\alpha-1}$ and taking the integral over s , we get

$$\begin{aligned} & \int_0^1 s^{\alpha-1} \frac{f \circ g^{-1}(sg(x) + (1-s)g(b)) + f \circ g^{-1}((1-s)g(a) + sg(x))}{2} ds \\ & \leq \int_0^1 s^{\alpha-1} \left[(1-s) \frac{f(a) + f(b)}{2} + sf(x) \right] ds \\ & = \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(x) \right] \end{aligned}$$

for any $x \in (a, b)$, which proves the last part of (2.1). \square

Corollary 1. *With the assumptions of Theorem 2, we have*

$$\begin{aligned} (2.6) \quad f(M_g(a, b)) & \leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(g(b) - g(a))^\alpha} \left[I_{M_g(a,b)+,g}^\alpha f(b) + I_{M_g(a,b)-,g}^\alpha f(a) \right] \\ & \leq \frac{\alpha}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(M_g(a, b)) \right] \leq \frac{f(a) + f(b)}{2}. \end{aligned}$$

Proof. If we take $x = M_g(a, b)$ in (2.1), then we get

$$\begin{aligned} & \frac{1}{\alpha} f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha + 1} + g(M_g(a, b)) \frac{\alpha}{\alpha + 1} \right) \\ & \leq \int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + sg(M_g(a, b)) \right) ds \\ & \leq \frac{1}{2} \Gamma(\alpha) \left[\frac{I_{M_g(a,b)+,g}^\alpha f(b)}{(g(b) - g(M_g(a, b)))^\alpha} + \frac{I_{M_g(a,b)-,g}^\alpha f(a)}{(g(M_g(a, b)) - g(a))^\alpha} \right] \\ & \leq \frac{1}{\alpha + 1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(M_g(a, b)) \right], \end{aligned}$$

that is equivalent to

$$\begin{aligned} \frac{1}{\alpha} f(M_g(a, b)) & \leq 2^{\alpha-1} \Gamma(\alpha) \left[\frac{I_{M_g(a,b)+,g}^\alpha f(b)}{(g(b) - g(a))^\alpha} + \frac{I_{M_g(a,b)-,g}^\alpha f(a)}{(g(b) - g(a))^\alpha} \right] \\ & \leq \frac{1}{\alpha + 1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(M_g(a, b)) \right], \end{aligned}$$

which proves the first two inequalities in (2.6).

By the convexity of $f \circ g^{-1}$ we also have

$$\begin{aligned} f(M_g(a, b)) & = f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \right) \leq \frac{(f \circ g^{-1})(g(a)) + f \circ g^{-1}(g(b))}{2} \\ & = \frac{f(a) + f(b)}{2}. \end{aligned}$$

Therefore

$$\frac{1}{\alpha + 1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(M_g(a, b)) \right] \leq \frac{1}{\alpha} \frac{f(a) + f(b)}{2},$$

which proves the last part of (2.6). \square

Corollary 2. *With the assumptions of Theorem 2, we have the integral inequalities*

$$\begin{aligned} (2.7) \quad & \frac{1}{\alpha} f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha + 1} + \frac{\alpha}{\alpha + 1} \frac{1}{b-a} \int_a^b g(x) dx \right) \\ & \leq \frac{1}{\alpha} \frac{1}{b-a} \int_a^b f \circ g^{-1} \left(\frac{g(a) + g(b)}{2} \frac{1}{\alpha + 1} + \frac{\alpha}{\alpha + 1} g(x) \right) dx \\ & \leq \frac{1}{b-a} \int_a^b \left(\int_0^1 s^{\alpha-1} f \circ g^{-1} \left((1-s) \frac{g(a) + g(b)}{2} + sg(x) \right) ds \right) dx \\ & \leq \frac{1}{2} \Gamma(\alpha) \frac{1}{b-a} \int_a^b \left[\frac{I_{x+,g}^\alpha f(b)}{(g(b) - g(x))^\alpha} + \frac{I_{x-,g}^\alpha f(a)}{(g(x) - g(a))^\alpha} \right] dx \\ & \leq \frac{1}{\alpha + 1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(x) dx \right]. \end{aligned}$$

The first inequality in (2.6) follows by Jensen's inequality for the convex function $f \circ g^{-1}$. The rest follows by taking the integral mean in (2.1).

Remark 1. In the case of $g(t) = t$, $t \in [a, b]$, and if f is convex on (a, b) , then we have by (2.1) and (2.6) the following inequalities for the classical Riemann-Liouville fractional integrals J_{a+}^α and J_{b-}^α that have been obtained in [8]

$$(2.8) \quad \begin{aligned} \frac{1}{\alpha} f \left(\frac{a+b}{2} \frac{1}{\alpha+1} + \frac{\alpha}{\alpha+1} x \right) &\leq \int_0^1 s^{\alpha-1} f \left((1-s) \frac{a+b}{2} + sx \right) ds \\ &\leq \frac{1}{2} \Gamma(\alpha) \left[\frac{J_{x+}^\alpha f(b)}{(b-a)^\alpha} + \frac{J_{x-}^\alpha f(a)}{(x-a)^\alpha} \right] \\ &\leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(x) \right] \end{aligned}$$

for any $x \in (a, b)$ and

$$(2.9) \quad \begin{aligned} f \left(\frac{a+b}{2} \right) &\leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[J_{\frac{a+b}{2}+}^\alpha f(b) + J_{\frac{a+b}{2}-}^\alpha f(a) \right] \\ &\leq \frac{\alpha}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f \left(\frac{a+b}{2} \right) \right] \leq \frac{f(a) + f(b)}{2}. \end{aligned}$$

We notice that the inequality (2.9) is an improvement of the result obtained by Sarikaya and Yildirim in [22], in which they prove the inequality for the first and last term.

From (2.7), we have for a convex function f that

$$(2.10) \quad \begin{aligned} \frac{1}{\alpha} f \left(\frac{a+b}{2} \right) &\leq \frac{1}{\alpha} \frac{1}{b-a} \int_a^b f \left(\frac{a+b}{2} \frac{1}{\alpha+1} + \frac{\alpha}{\alpha+1} x \right) dx \\ &\leq \frac{1}{b-a} \int_a^b \left(\int_0^1 s^{\alpha-1} f \left((1-s) \frac{a+b}{2} + sx \right) ds \right) dx \\ &\leq \frac{1}{2} \Gamma(\alpha) \frac{1}{b-a} \int_a^b \left[\frac{J_{x+}^\alpha f(b)}{(b-x)^\alpha} + \frac{J_{x-}^\alpha f(a)}{(x-a)^\alpha} \right] dx \\ &\leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(x) dx \right]. \end{aligned}$$

For some recent papers on Hermite-Hadamard inequalities for the Riemann-Liouville fractional integrals see [1], [3], [4], [12]-[27] and the references therein.

3. APPLICATIONS FOR GA-CONVEX FUNCTIONS

Let $I \subset (0, \infty)$ be an interval; a real-valued function $f : I \rightarrow \mathbb{R}$ is said to be *GA-convex* (concave) on I if

$$(3.1) \quad f(x^{1-\lambda} y^\lambda) \leq (\geq) (1-\lambda) f(x) + \lambda f(y)$$

for all $x, y \in I$ and $\lambda \in [0, 1]$.

Since the condition (3.1) can be written as

$$(3.2) \quad f \circ \exp((1-\lambda) \ln x + \lambda \ln y) \leq (\geq) (1-\lambda) f \circ \exp(\ln x) + \lambda f \circ \exp(\ln y),$$

then we observe that $f : I \rightarrow \mathbb{R}$ is *GA-convex* (concave) on I if and only if $f \circ \exp$ is convex (concave) on $\ln I := \{\ln z, z \in I\}$. If $I = [a, b]$ then $\ln I = [\ln a, \ln b]$.

It is known that the function $f(x) = \ln(1+x)$ is *GA-convex* on $(0, \infty)$ [2].

For real and positive values of x , the *Euler gamma* function Γ and its *logarithmic derivative* ψ , the so-called *digamma function*, are defined by

$$\Gamma(x) := \int_0^\infty t^{x-1} e^{-t} dt \text{ and } \psi(x) := \frac{\Gamma'(x)}{\Gamma(x)}.$$

It has been shown in [26] that the function $f : (0, \infty) \rightarrow \mathbb{R}$ defined by

$$f(x) = \psi(x) + \frac{1}{2x}$$

is *GA*-concave on $(0, \infty)$ while the function $g : (0, \infty) \rightarrow \mathbb{R}$ defined by

$$g(x) = \psi(x) + \frac{1}{2x} + \frac{1}{12x^2}$$

is *GA*-convex on $(0, \infty)$.

For some recent inequalities on *GA*-convex functions see [5]-[7].

If we assume that the function $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ is *GA*-convex on $[a, b]$, then by Theorem 2 for $g : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$, $g(t) = \ln t$, we have the following inequalities for Hadamard fractional integrals:

$$\begin{aligned} (3.3) \quad \frac{1}{\alpha} f \left([G(a, b)]^{\frac{1}{\alpha+1}} x^{\frac{\alpha}{\alpha+1}} \right) &\leq \int_0^1 s^{\alpha-1} f \left([G(a, b)]^{1-s} x^s \right) ds \\ &\leq \frac{1}{2} \Gamma(\alpha) \left[\frac{H_{a+}^\alpha f(x)}{[\ln(\frac{x}{a})]^\alpha} + \frac{H_{b-}^\alpha f(x)}{[\ln(\frac{b}{x})]^\alpha} \right] \\ &\leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(x) \right] \end{aligned}$$

for any $x \in (a, b)$, where $G(a, b) := \sqrt{ab}$ is the geometric mean of a, b .

In particular, we have

$$\begin{aligned} (3.4) \quad f(G(a, b)) &\leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{[\ln(\frac{b}{a})]^\alpha} [H_{a+}^\alpha f(G(a, b)) + H_{b-}^\alpha f(G(a, b))] \\ &\leq \frac{\alpha}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(G(a, b)) \right] \leq \frac{f(a) + f(b)}{2}, \end{aligned}$$

provided $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ is *GA*-convex.

We also have the integral inequalities

$$\begin{aligned} (3.5) \quad \frac{1}{\alpha} f \left([G(a, b)]^{\frac{1}{\alpha+1}} [I(a, b)]^{\frac{\alpha}{\alpha+1}} \right) &\leq \frac{1}{\alpha} \frac{1}{b-a} \int_a^b f \left([G(a, b)]^{\frac{1}{\alpha+1}} x^{\frac{\alpha}{\alpha+1}} \right) dx \\ &\leq \frac{1}{b-a} \int_a^b \left(\int_0^1 s^{\alpha-1} f \left([G(a, b)]^{1-s} x^s \right) ds \right) dx \\ &\leq \frac{1}{2} \Gamma(\alpha) \frac{1}{b-a} \int_a^b \left[\frac{H_{a+}^\alpha f(x)}{[\ln(\frac{x}{a})]^\alpha} + \frac{H_{b-}^\alpha f(x)}{[\ln(\frac{b}{x})]^\alpha} \right] dx \\ &\leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(x) \right] \end{aligned}$$

since $\ln I(a, b) = \frac{1}{b-a} \int_a^b \ln x dx$ and the *identric mean* $I(a, b)$ is defined by $I(a, b) = \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}}$ for $0 < a < b$.

4. APPLICATION FOR EXPONENTIAL FRACTIONAL INTEGRAL

Consider the "*Exponential fractional integrals*"

$$E_{a+}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \frac{\exp(t) f(t) dt}{[\exp(x) - \exp(t)]^{1-\alpha}}, \quad a < x \leq b$$

and

$$E_{b-}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \frac{\exp(t) f(t) dt}{[\exp(t) - \exp(x)]^{1-\alpha}}, \quad a \leq x < b.$$

Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function on $[a, b] \subset \mathbb{R}$. If $f \circ \ln$ is convex on $(\exp(a), \exp(b))$, then for any $x \in (a, b)$ we have by Theorem 2 the inequalities

$$\begin{aligned} (4.1) \quad & \frac{1}{\alpha} f \left[\ln \left(\frac{1}{\alpha+1} \frac{\exp a + \exp b}{2} + \frac{\alpha}{\alpha+1} \exp x \right) \right] \\ & \leq \int_0^1 s^{\alpha-1} f \left[\ln \left((1-s) \frac{\exp a + \exp b}{2} + s \exp x \right) \right] ds \\ & \leq \frac{1}{2} \Gamma(\alpha) \left[\frac{E_{a+}^{\alpha} f(x)}{(\exp x - \exp a)^{\alpha}} + \frac{E_{b-}^{\alpha} f(x)}{(\exp b - \exp x)^{\alpha}} \right] \\ & \leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(x) \right]. \end{aligned}$$

In particular, we have

$$\begin{aligned} (4.2) \quad & f(LME(a, b)) \\ & \leq \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(\exp b - \exp a)^{\alpha}} [E_{a+}^{\alpha} f(LME(a, b)) + E_{b-}^{\alpha} f(LME(a, b))] \\ & \leq \frac{\alpha}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + f(LME(a, b)) \right] \leq \frac{f(a) + f(b)}{2}, \end{aligned}$$

where $LME(a, b) := \ln \left(\frac{\exp a + \exp b}{2} \right)$, provided that $f \circ \ln$ is convex on $(\exp(a), \exp(b))$.

Finally, by (2.10) we have

$$\begin{aligned} (4.3) \quad & \frac{1}{\alpha} f \left[\ln \left(\frac{1}{\alpha+1} \frac{\exp a + \exp b}{2} + \frac{\alpha}{\alpha+1} \frac{\exp b - \exp a}{b-a} \right) \right] \\ & \leq \frac{1}{\alpha} \frac{1}{b-a} \int_a^b f \left[\ln \left(\frac{1}{\alpha+1} \frac{\exp a + \exp b}{2} + \frac{\alpha}{\alpha+1} \exp x \right) \right] dx \\ & \leq \frac{1}{b-a} \int_a^b \left(\int_0^1 s^{\alpha-1} f \left[\ln \left((1-s) \frac{\exp a + \exp b}{2} + s \exp x \right) \right] ds \right) dx \\ & \leq \frac{1}{2} \Gamma(\alpha) \frac{1}{b-a} \int_a^b \left[\frac{E_{a+}^{\alpha} f(x)}{(g(x) - g(a))^{\alpha}} + \frac{E_{b-}^{\alpha} f(x)}{(g(b) - g(x))^{\alpha}} \right] dx \\ & \leq \frac{1}{\alpha+1} \left[\frac{1}{\alpha} \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(x) dx \right], \end{aligned}$$

provided that $f \circ \ln$ is convex on $(\exp(a), \exp(b))$.

REFERENCES

- [1] P. Agarwal, M. Jleli and M. Tomar, Certain Hermite-Hadamard type inequalities via generalized k -fractional integrals, *Journal of Inequalities and Applications* **2017** (2017) :55.
- [2] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, Generalized convexity and inequalities, *J. Math. Anal. Appl.* **335** (2007) 1294–1308.
- [3] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.* **1**(1) (2010), 51-58.
- [4] J. Deng and J. Wang, Fractional Hermite-Hadamard inequalities for $(\alpha; m)$ -logarithmically convex functions. *J. Inequal. Appl.* **2013**, Article ID 364 (2013)
- [5] S. S. Dragomir, Inequalities of Hermite-Hadamard type for GA-convex functions, *RGMIA Res. Rep. Coll.* **18** (2015), Art. 30. [Online <http://rgmia.org/papers/v18/v18a30.pdf>].
- [6] S. S. Dragomir, Some new inequalities of Hermite-Hadamard type for GA-convex functions, *RGMIA Res. Rep. Coll.* **18** (2015), Art. 33. [Online <http://rgmia.org/papers/v18/v18a33.pdf>].
- [7] S. S. Dragomir, Inequalities of Jensen type for GA-convex functions, *RGMIA Res. Rep. Coll.* **18** (2015), Art. 35. [Online <http://rgmia.org/papers/v18/v18a35.pdf>].
- [8] S. S. Dragomir, Some inequalities of Hermite-Hadamard type for convex functions and Riemann-Liouville fractional integrals, *RGMIA Res. Rep. Coll.* **20** (2017), Art. 40. [Online <http://rgmia.org/papers/v20/v20a40.pdf>].
- [9] S. S. Dragomir, Ostrowski type inequalities for Lebesgue integral: a survey of recent results, *Australian J. Math. Anal. Appl.*, Volume **14**, Issue 1, Article 1, pp. 1-287, 2017. [Online <http://ajmaa.org/cgi-bin/paper.pl?string=v14n1/V14I1P1.tex>].
- [10] S. S. Dragomir, Hermite-Hadamard type inequalities of first kind for generalized Riemann-Liouville fractional integrals, *RGMIA Res. Rep. Coll.* **20** (2017), Art. 76. [Online <http://rgmia.org/papers/v20/v20a76.pdf>].
- [11] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, 2000. [Online http://rgmia.org/monographs/hermite_hadamard.html].
- [12] S.-R. Hwang, K.-L. Tseng and K.-C. Hsu, New inequalities for fractional integrals and their applications, *Turk. J. Math.* **40** (2016): 471-486.
- [13] A. Kashuri and R. Liko, Hermite-Hadamard type fractional integral inequalities for the generalized $(1; m)$ -preinvex functions, *Nonlinear Analysis and Differential Equations*, Vol. **4**, 2016, no. 8, 353 - 367.
- [14] A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*. North-Holland Mathematics Studies, 204. Elsevier Science B.V., Amsterdam, 2006. xvi+523 pp. ISBN: 978-0-444-51832-3; 0-444-51832-0
- [15] M. Kunt and I. Iscan, Hermite-Hadamard-Fejer type inequalities for p -convex functions via fractional integrals, *Communication in Mathematical Modeling and Applications*, **2** (2017), No. 1, 1-15.
- [16] M. A. Khan, Y. Khurshid and T. Ali, Hermite-Hadamard inequality for fractional integrals via η -convex functions, *Acta Math. Univ. Comenianae* Vol. **LXXXVI**, 1 (2017), pp. 153-164.
- [17] W. Liu and W. Wen, Some generalizations of different type of integral inequalities for MT-convex functions, *Filomat* **30**:2 (2016), 333–342
- [18] M. E. Özdemir and Ç. Yildiz, The Hadamard's inequality for quasi-convex functions via fractional integrals, *Annals of the University of Craiova, Mathematics and Computer Science Series*, Volume **40** (2), 2013, Pages 167-173.
- [19] S. Qaisar, M. Iqbal, S. Hussain, S. I. Butt and M. A. Meraj, New inequalities on Hermite-Hadamard utilizing fractional integrals, *Kragujevac J. Math.*, **42** (1) (2018), Pages 15–27.
- [20] M. Z. Sarikaya, H. Filiz and M. E. Kiris, On some generalized integral inequalities for Riemann-Liouville fractional integrals, *Filomat*, **29**:6 (2015), 1307–1314.
- [21] M. Z. Sarikaya, E. Set, H. Yaldiz, and N. Basak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, *Mathematical and Computer Modelling*, **57** (2013), pp.2403–2407, doi:10.1016/j.mcm.2011.12.048.
- [22] M. Z. Sarikaya and H. Yildirim, On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals, *Miskolc Mathematical Notes*, **17** (2016), No. 2, pp. 1049–1059.

- [23] D.-P. Shi, B.-Y. Xi and F. Qi, Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals of (α, m) -convex functions, *Fractional Differential Calculus*, Volume **4**, Number 1 (2014), 31–43
- [24] M. Tunc, On new inequalities for h -convex functions via Riemann-Liouville fractional integration, *Filomat* **27**:4 (2013), 559–565.
- [25] J.-R. Wang, X. Lia and Y. Zhou, Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals via s -convex functions and applications to special Means, *Filomat* **30**:5 (2016), 1143–1150.
- [26] X.-M. Zhang, Y.-M. Chu and X.-H. Zhang, The Hermite-Hadamard type inequality of GA-convex functions and its application, *Journal of Inequalities and Applications*, Volume **2010**, Article ID 507560, 11 pages.
- [27] Y. Zhang and J. Wang, On some new Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals, *J. Inequal. Appl.* **2013**, Article ID 220 (2013)

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.org/dragomir>

²DST-NRF CENTRE OF EXCELLENCE, IN THE MATHEMATICAL AND STATISTICAL SCIENCES, SCHOOL OF COMPUTER SCIENCE & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA