

**JENSEN'S TYPE INEQUALITIES FOR THE FINITE HILBERT
TRANSFORM OF CONVEX FUNCTIONS**

SILVESTRU SEVER DRAGOMIR^{1,2}

ABSTRACT. In this paper we obtain some new inequalities for the finite Hilbert transform of convex functions by the use of Jensen's integral inequality. Applications for exponential function are provided as well.

1. INTRODUCTION

Allover this paper, we consider the *finite Hilbert transform* on the open interval (a, b) defined by

$$(Tf)(a, b; t) := \frac{1}{\pi} PV \int_a^b \frac{f(\tau)}{\tau - t} d\tau := \lim_{\varepsilon \rightarrow 0+} \left[\int_a^{t-\varepsilon} + \int_{t+\varepsilon}^b \right] \frac{f(\tau)}{\pi(\tau - t)} d\tau$$

for $t \in (a, b)$ and for various classes of functions f for which the above Cauchy Principal Value integral exists, see [13, Section 3.2] or [17, Lemma II.1.1].

For several recent papers devoted to inequalities for the finite Hilbert transform (Tf) , see [2]-[10], [14]-[16] and [18]-[19].

Now, if we assume that the mapping $f : (a, b) \rightarrow \mathbb{R}$ is convex on (a, b) , then it is locally Lipschitzian on (a, b) and then the finite Hilbert transform of f exists in every point $t \in (a, b)$.

The following result concerning upper and lower bounds for the finite Hilbert transform of a convex function holds.

Theorem 1 (Dragomir et al., 2001 [1]). *Let $f : (a, b) \rightarrow \mathbb{R}$ be a convex function on (a, b) and $t \in (a, b)$. Then we have*

$$\begin{aligned} (1.1) \quad & \frac{1}{\pi} \left[f(t) \ln \left(\frac{b-t}{t-a} \right) + f(t) - f(a) + \varphi(t)(b-t) \right] \\ & \leq (Tf)(a, b; t) \\ & \leq \frac{1}{\pi} \left[f(t) \ln \left(\frac{b-t}{t-a} \right) + f(b) - f(t) + \varphi(t)(t-a) \right], \end{aligned}$$

where $\varphi(t) \in [f'_-(t), f'_+(t)]$, $t \in (a, b)$.

1991 *Mathematics Subject Classification.* 26D15; 26D10.

Key words and phrases. Finite Hilbert Transform, Convex functions .

Corollary 1. Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable convex function on (a, b) and $t \in (a, b)$. Then we have

$$(1.2) \quad \begin{aligned} & \frac{1}{\pi} \left[f(t) \ln \left(\frac{b-t}{t-a} \right) + f(t) - f(a) + f'(t)(b-t) \right] \\ & \leq (Tf)(a, b; t) \\ & \leq \frac{1}{\pi} \left[f(t) \ln \left(\frac{b-t}{t-a} \right) + f(b) - f(t) + f'(t)(t-a) \right]. \end{aligned}$$

We observe that if we take $t = \frac{a+b}{2}$, then we get from (1.2) that

$$(1.3) \quad \begin{aligned} & \frac{1}{\pi} \left[f \left(\frac{a+b}{2} \right) - f(a) + \frac{1}{2} f' \left(\frac{a+b}{2} \right) (b-a) \right] \\ & \leq (Tf) \left(a, b; \frac{a+b}{2} \right) \\ & \leq \frac{1}{\pi} \left[f(b) - f \left(\frac{a+b}{2} \right) + \frac{1}{2} f' \left(\frac{a+b}{2} \right) (b-a) \right]. \end{aligned}$$

More recently, we have established the following result as well:

Theorem 2 (Dragomir, 2018 [12]). Let $f : (a, b) \rightarrow \mathbb{R}$ be a convex function on (a, b) with finite lateral derivatives $f'_+(a)$ and $f'_-(b)$. Then for $t \in (a, b)$ we have

$$(1.4) \quad \begin{aligned} & \frac{1}{\pi} (b-a) f'_+(a) \leq \frac{1}{\pi} (b-a) \frac{f(t) - f(a)}{t-a} \\ & \leq (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) \\ & \leq \frac{1}{\pi} (b-a) \frac{f(b) - f(t)}{b-t} \leq \frac{1}{\pi} (b-a) f'_-(b). \end{aligned}$$

In particular,

$$(1.5) \quad \begin{aligned} & \frac{1}{\pi} (b-a) f'_+(a) \leq \frac{2}{\pi} (b-a) \frac{f \left(\frac{a+b}{2} \right) - f(a)}{b-a} \\ & \leq (Tf) \left(a, b; \frac{a+b}{2} \right) \\ & \leq \frac{2}{\pi} (b-a) \frac{f(b) - f \left(\frac{a+b}{2} \right)}{b-a} \leq \frac{1}{\pi} (b-a) f'_-(b). \end{aligned}$$

In this paper, by the use of Jensen's celebrated inequality, we obtain some new inequalities for the finite Hilbert transform of convex functions. Applications for exponential function are provided as well.

2. MAIN RESULTS

We have:

Theorem 3. Let $f : (a, b) \rightarrow \mathbb{R}$ be a convex function on (a, b) and $x : (a, b) \rightarrow \mathbb{R}$ a locally absolutely continuous function on (a, b) with $x'(s) \in (a, b)$ for almost every

$s \in (a, b)$. Then we have

$$(2.1) \quad f\left(\frac{1}{b-a}\left[\pi(Tx)(a, b; t) - x(t) \ln\left(\frac{b-t}{t-a}\right)\right]\right) \leq \frac{1}{b-a}\left[\pi T\left(\int_a^t f \circ x'(s) ds\right)(a, b; t) - \left(\int_a^t f \circ x'(s) ds\right) \ln\left(\frac{b-t}{t-a}\right)\right],$$

for any $t \in (a, b)$.

In particular,

$$(2.2) \quad f\left(\frac{\pi}{b-a}(Tx)\left(a, b; \frac{a+b}{2}\right)\right) \leq \frac{\pi}{b-a}T\left(\int_a^b f \circ x'(s) ds\right)\left(a, b; \frac{a+b}{2}\right).$$

Proof. Using Jensen's integral inequality for the convex function f we have

$$(2.3) \quad f\left(\frac{x(\tau) - x(t)}{\tau - t}\right) = f\left(\frac{\int_t^\tau x'(s) ds}{\tau - t}\right) \leq \frac{\int_t^\tau f \circ x'(s) ds}{\tau - t}$$

for any $\tau, t \in (a, b)$ with $\tau \neq t$.

Let $t \in (a, b)$ and $t - a > \varepsilon > 0$, then by integrating (2.3) over $\tau \in [a, t - \varepsilon]$ we get

$$(2.4) \quad \int_a^{t-\varepsilon} f\left(\frac{x(\tau) - x(t)}{\tau - t}\right) d\tau \leq \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t}\right) d\tau.$$

Using Jensen's integral inequality for the convex function f we get

$$(2.5) \quad f\left(\frac{1}{t-\varepsilon-a} \int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau\right) \leq \frac{1}{t-\varepsilon-a} \int_a^{t-\varepsilon} f\left(\frac{x(\tau) - x(t)}{\tau - t}\right) d\tau.$$

By using (2.4) and (2.5) we get

$$(2.6) \quad (t - \varepsilon - a) f\left(\frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau\right) \leq \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t}\right) d\tau$$

for $t \in (a, b)$ and $t - a > \varepsilon > 0$.

Let $t \in (a, b)$ and $b - t > \varepsilon > 0$, then by integrating (2.3) over $\tau \in [t + \varepsilon, b]$ we obtain

$$(2.7) \quad \int_{t+\varepsilon}^b f\left(\frac{x(\tau) - x(t)}{\tau - t}\right) d\tau \leq \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t}\right) d\tau.$$

Using Jensen's integral inequality for the convex function f we get

$$(2.8) \quad f\left(\frac{1}{b-t-\varepsilon} \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau\right) \leq \frac{1}{b-t-\varepsilon} \int_{t+\varepsilon}^b f\left(\frac{x(\tau) - x(t)}{\tau - t}\right) d\tau.$$

On utilising (2.7) and (2.8) we deduce

$$(2.9) \quad (b - t - \varepsilon) f\left(\frac{1}{b-t-\varepsilon} \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau\right) \leq \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t}\right) d\tau$$

for $t \in (a, b)$ and $b - t > \varepsilon > 0$.

If we add the inequalities (2.6) and (2.9) we get

$$(2.10) \quad \begin{aligned} & (t - \varepsilon - a) f \left(\frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) \\ & + (b - t - \varepsilon) f \left(\frac{1}{b - t - \varepsilon} \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) \\ & \leq \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t} \right) d\tau + \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t} \right) d\tau \end{aligned}$$

for $t \in (a, b)$ and $\min\{b - t, t - a\} > \varepsilon > 0$.

By the convexity of f we also have

$$(2.11) \quad \begin{aligned} & (t - \varepsilon - a) f \left(\frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) \\ & + (b - t - \varepsilon) f \left(\frac{1}{b - t - \varepsilon} \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) \\ & \geq (b - a - 2\varepsilon) f \left(\frac{1}{b - a - 2\varepsilon} \left[\int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \right) \end{aligned}$$

for $t \in (a, b)$ and $\min\{b - t, t - a\} > \varepsilon > 0$.

Therefore, by (2.10) and (2.11) we get

$$(2.12) \quad \begin{aligned} & (b - a - 2\varepsilon) f \left(\frac{1}{b - a - 2\varepsilon} \left[\int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \right) \\ & \leq \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t} \right) d\tau + \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau f \circ x'(s) ds}{\tau - t} \right) d\tau \\ & = \int_a^{t-\varepsilon} \left(\frac{\int_a^\tau f \circ x'(s) ds - \int_a^t f \circ x'(s) ds}{\tau - t} \right) d\tau \\ & + \int_{t+\varepsilon}^b \left(\frac{\int_a^\tau f \circ x'(s) ds - \int_a^t f \circ x'(s) ds}{\tau - t} \right) d\tau \end{aligned}$$

for $t \in (a, b)$ and $\min\{b - t, t - a\} > \varepsilon > 0$.

Taking the limit over $\varepsilon \rightarrow 0+$ in (2.12) we get

$$(2.13) \quad \begin{aligned} & f \left(\frac{1}{b - a} PV \left(\int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) \right) \\ & \leq \frac{1}{b - a} PV \int_a^b \left(\frac{\int_a^\tau f \circ x'(s) ds - \int_a^t f \circ x'(s) ds}{\tau - t} \right) d\tau, \end{aligned}$$

for $t \in (a, b)$, which is an inequality of interest in itself.

As for the mapping , $\mathbf{1}(t) = 1$, $t \in (a, b)$, we have

$$\begin{aligned} (T\mathbf{1})(a, b; t) &= \frac{1}{\pi} PV \int_a^b \frac{1}{\tau - t} d\tau \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0+} \left[\int_a^{t-\varepsilon} \frac{1}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{1}{\tau - t} d\tau \right] \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0+} \left[\ln |\tau - t|_a^{t-\varepsilon} + \ln (\tau - t)|_{t+\varepsilon}^b \right] \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0+} [\ln \varepsilon - \ln (t - a) + \ln (b - t) - \ln \varepsilon] \\ &= \frac{1}{\pi} \ln \left(\frac{b - t}{t - a} \right), \quad t \in (a, b). \end{aligned}$$

Then, obviously, for $g : (a, b) \rightarrow \mathbb{R}$ we have

$$\begin{aligned} (Tg)(a, b; t) &= \frac{1}{\pi} PV \int_a^b \frac{g(\tau) - g(t) + g(t)}{\tau - t} d\tau \\ &= \frac{1}{\pi} PV \int_a^b \frac{g(\tau) - g(t)}{\tau - t} d\tau + \frac{g(t)}{\pi} PV \int_a^b \frac{1}{\tau - t} d\tau \end{aligned}$$

from where we get the equality

$$(2.14) \quad (Tg)(a, b; t) - \frac{g(t)}{\pi} \ln \left(\frac{b - t}{t - a} \right) = \frac{1}{\pi} PV \int_a^b \frac{g(\tau) - g(t)}{\tau - t} d\tau$$

for all $t \in (a, b)$.

Using the equality (2.14) we have

$$PV \left(\int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) = \pi (Tx)(a, b; t) - x(t) \ln \left(\frac{b - t}{t - a} \right)$$

and

$$\begin{aligned} PV \int_a^b \left(\frac{\int_a^\tau f \circ x'(s) ds - \int_a^t f \circ x'(s) ds}{\tau - t} \right) d\tau \\ = \pi \left(T \left(\int_a^\cdot f \circ x'(s) ds \right) \right) (a, b; t) - \int_a^t f \circ x'(s) ds \ln \left(\frac{b - t}{t - a} \right) \end{aligned}$$

and by (2.13) we get the desired result (2.1). \square

We also have the reverse inequality:

Theorem 4. Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$ and $x : [a, b] \rightarrow \mathbb{R}$ an absolutely continuous function on $[a, b]$ with $x'(s) \in [a, b]$ for almost every $s \in (a, b)$. Then we have

$$\begin{aligned} (2.15) \quad &\frac{1}{b - a} \left[\pi T \left(\int_a^\cdot f \circ x'(s) ds \right) (a, b; t) - \left(\int_a^t f \circ x'(s) ds \right) \ln \left(\frac{b - t}{t - a} \right) \right] \\ &\leq \frac{f(a)}{b - a} \left[b - \frac{1}{b - a} \left(\pi (Tx)(a, b; t) - x(t) \ln \left(\frac{b - t}{t - a} \right) \right) \right] \\ &+ \frac{f(b)}{b - a} \left[\frac{1}{b - a} \left(\pi (Tx)(a, b; t) - x(t) \ln \left(\frac{b - t}{t - a} \right) \right) - a \right] \end{aligned}$$

for any $t \in (a, b)$.

In particular,

$$(2.16) \quad \begin{aligned} & \frac{\pi}{b-a} T \left(\int_a^b f \circ x'(s) ds \right) \left(a, b; \frac{a+b}{2} \right) \\ & \leq \frac{f(a)}{b-a} \left[b - \frac{\pi}{b-a} (Tx) \left(a, b; \frac{a+b}{2} \right) \right] \\ & + \frac{f(b)}{b-a} \left[\frac{\pi}{b-a} (Tx) \left(a, b; \frac{a+b}{2} \right) - a \right]. \end{aligned}$$

Proof. By the convexity of f we have

$$(2.17) \quad \begin{aligned} f(x'(s)) &= f\left(\frac{(b-x'(s))a+(x'(s)-a)b}{b-a}\right) \\ &\leq \frac{b-x'(s)}{b-a} f(a) + \frac{x'(s)-a}{b-a} f(b) \end{aligned}$$

for almost every $s \in [a, b]$.

Let $t, \tau \in [a, b]$ with $t \neq \tau$. Taking the integral mean $\frac{1}{\tau-t} \int_t^\tau$, which is a linear and isotonic functional, in the inequality (2.17), we get

$$(2.18) \quad \begin{aligned} & \frac{1}{\tau-t} \int_t^\tau f(x'(s)) ds \\ & \leq \frac{1}{\tau-t} \int_t^\tau \left[\frac{b-x'(s)}{b-a} f(a) + \frac{x'(s)-a}{b-a} f(b) \right] ds \\ & = \frac{b - \frac{1}{\tau-t} \int_t^\tau x'(s) ds}{b-a} f(a) + \frac{\frac{1}{\tau-t} \int_t^\tau x'(s) ds - a}{b-a} f(b) \\ & = \frac{b - \frac{x(\tau)-x(t)}{\tau-t}}{b-a} f(a) + \frac{\frac{x(\tau)-x(t)}{\tau-t} - a}{b-a} f(b) \\ & = \frac{f(a)}{b-a} \left[b - \frac{x(\tau)-x(t)}{\tau-t} \right] + \frac{f(b)}{b-a} \left[\frac{x(\tau)-x(t)}{\tau-t} - a \right] \end{aligned}$$

for any $t, \tau \in [a, b]$ with $t \neq \tau$.

Let $t \in (a, b)$ and $t-a > \varepsilon > 0$, then by integrating (2.18) over $\tau \in [a, t-\varepsilon]$ we get

$$(2.19) \quad \begin{aligned} & \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau-t} \right) d\tau \\ & \leq \int_a^{t-\varepsilon} \left(\frac{f(a)}{b-a} \left[b - \frac{x(\tau)-x(t)}{\tau-t} \right] + \frac{f(b)}{b-a} \left[\frac{x(\tau)-x(t)}{\tau-t} - a \right] \right) d\tau \\ & = \frac{f(a)}{b-a} \left[b(t-\varepsilon-a) - \int_a^{t-\varepsilon} \frac{x(\tau)-x(t)}{\tau-t} d\tau \right] \\ & + \frac{f(b)}{b-a} \left[\int_a^{t-\varepsilon} \frac{x(\tau)-x(t)}{\tau-t} d\tau - a(t-\varepsilon-a) \right]. \end{aligned}$$

Let $t \in (a, b)$ and $b - t > \varepsilon > 0$, then by integrating (2.18) over $\tau \in [t + \varepsilon, b]$ we obtain

$$\begin{aligned}
(2.20) \quad & \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau - t} \right) d\tau \\
& \leq \int_{t+\varepsilon}^b \left(\frac{f(a)}{b-a} \left[b - \frac{x(\tau) - x(t)}{\tau - t} \right] + \frac{f(b)}{b-a} \left[\frac{x(\tau) - x(t)}{\tau - t} - a \right] \right) d\tau \\
& = \frac{f(a)}{b-a} \left[b(b-t-\varepsilon) - \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \\
& \quad + \frac{f(b)}{b-a} \left[\int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau - a(b-t-\varepsilon) \right].
\end{aligned}$$

For $t \in (a, b)$ and $\min\{b-t, t-a\} > \varepsilon > 0$, we have by adding the inequalities (2.19) and (2.20) that

$$\begin{aligned}
(2.21) \quad & \int_a^{t-\varepsilon} \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau - t} \right) d\tau + \int_{t+\varepsilon}^b \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau - t} \right) d\tau \\
& \leq \frac{f(a)}{b-a} \left[b(t-\varepsilon-a) - \int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau \right. \\
& \quad \left. + b(b-t-\varepsilon) - \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \\
& \quad + \frac{f(b)}{b-a} \left[\int_a^{t-\varepsilon} \frac{x(\tau) - x(t)}{\tau - t} d\tau - a(t-\varepsilon-a) \right. \\
& \quad \left. + \int_{t+\varepsilon}^b \frac{x(\tau) - x(t)}{\tau - t} d\tau - a(b-t-\varepsilon) \right].
\end{aligned}$$

If we take $\varepsilon \rightarrow 0+$ in (2.21), then we obtain

$$\begin{aligned}
& PV \int_a^b \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau - t} \right) d\tau \\
& \leq \frac{f(a)}{b-a} \left[b(t-a) + b(b-t) - PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \\
& \quad + \frac{f(b)}{b-a} \left[PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau - a(t-a) - a(b-t) \right] \\
& = \frac{f(a)}{b-a} \left[b(b-a) - PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \\
& \quad + \frac{f(b)}{b-a} \left[PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau - a(b-a) \right],
\end{aligned}$$

which produces the following inequality of interest

$$(2.22) \quad PV \int_a^b \left(\frac{\int_t^\tau (f \circ x')(s) ds}{\tau - t} \right) d\tau \leq f(a) \left[b - \frac{1}{b-a} PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right] \\ + f(b) \left[\frac{1}{b-a} PV \int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau - a \right],$$

for any $t \in (a, b)$.

Now, since

$$PV \left(\int_a^b \frac{x(\tau) - x(t)}{\tau - t} d\tau \right) = \pi(Tx)(a, b; t) - x(t) \ln \left(\frac{b-t}{t-a} \right)$$

and

$$PV \int_a^b \left(\frac{\int_a^\tau f \circ x'(s) ds - \int_a^t f \circ x'(s) ds}{\tau - t} \right) d\tau \\ = \pi \left(T \left(\int_a^t f \circ x'(s) ds \right) \right) (a, b; t) - \int_a^t f \circ x'(s) ds \ln \left(\frac{b-t}{t-a} \right),$$

hence by (2.22) we get

$$\pi \left(T \left(\int_a^t f \circ x'(s) ds \right) \right) (a, b; t) - \int_a^t f \circ x'(s) ds \ln \left(\frac{b-t}{t-a} \right) \\ \leq f(a) \left[b - \frac{1}{b-a} \left(\pi(Tx)(a, b; t) - x(t) \ln \left(\frac{b-t}{t-a} \right) \right) \right] \\ + f(b) \left[\frac{1}{b-a} \left(\pi(Tx)(a, b; t) - x(t) \ln \left(\frac{b-t}{t-a} \right) \right) - a \right],$$

which proves (2.15). \square

3. EXAMPLES

If we consider the function $\exp t = e^t$, $t \in (a, b)$ a real interval, then

$$(3.1) \quad (T \exp)(a, b; t) = \frac{\exp(t)}{\pi} [E_i(b-t) - E_i(a-t)],$$

where E_i is defined by

$$E_i(x) := PV \int_{-\infty}^x \frac{\exp(s)}{s} ds, \quad x \in \mathbb{R}.$$

Indeed, we have

$$E_i(b-t) - E_i(a-t) = PV \int_{a-t}^{b-t} \frac{\exp(s)}{s} ds = PV \int_a^b \frac{\exp(\tau-t)}{\tau-t} ds \\ = \exp(-t) \pi(T \exp)(a, b; t)$$

and the equality (3.1) is proved.

Let $f : (a, b) \rightarrow \mathbb{R}$ be a convex function on (a, b) . Then we have by (2.1) and (2.15) for $x(t) = \exp t$, that

$$\begin{aligned} (3.2) \quad & f\left(\frac{\exp(t)}{b-a}\left[E_i(b-t) - E_i(a-t) - \ln\left(\frac{b-t}{t-a}\right)\right]\right) \\ & \leq \frac{1}{b-a}\left[\pi T\left(\int_a^t f \circ \exp(s) ds\right)(a, b; t) - \left(\int_a^t f \circ \exp(s) ds\right) \ln\left(\frac{b-t}{t-a}\right)\right] \\ & \leq \frac{f(a)}{b-a}\left[b - \frac{\exp(t)}{b-a}\left(E_i(b-t) - E_i(a-t) - \ln\left(\frac{b-t}{t-a}\right)\right)\right] \\ & \quad + \frac{f(b)}{b-a}\left[\frac{\exp(t)}{b-a}\left(E_i(b-t) - E_i(a-t) - \ln\left(\frac{b-t}{t-a}\right)\right) - a\right] \end{aligned}$$

for $t \in (a, b)$.

In particular, we have

$$\begin{aligned} (3.3) \quad & f\left(\frac{\exp\left(\frac{a+b}{2}\right)}{b-a}\left[E_i\left(\frac{b-a}{2}\right) - E_i\left(-\frac{b-a}{2}\right)\right]\right) \\ & \leq \frac{1}{b-a}\left[\pi T\left(\int_a^b f \circ \exp(s) ds\right)\left(a, b; \frac{a+b}{2}\right)\right] \\ & \leq \frac{f(a)}{b-a}\left[b - \frac{\exp\left(\frac{a+b}{2}\right)}{b-a}\left(E_i\left(\frac{b-a}{2}\right) - E_i\left(-\frac{b-a}{2}\right)\right)\right] \\ & \quad + \frac{f(b)}{b-a}\left[\frac{\exp\left(\frac{a+b}{2}\right)}{b-a}\left(E_i\left(\frac{b-a}{2}\right) - E_i\left(-\frac{b-a}{2}\right)\right) - a\right]. \end{aligned}$$

If we take $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \exp(\alpha x)$, $\alpha \neq 0$, then for an absolutely continuous function $x : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ we have from (2.1) and (2.15) that

$$\begin{aligned} (3.4) \quad & \exp\left(\frac{\alpha}{b-a}\left[\pi(Tx)(a, b; t) - x(t) \ln\left(\frac{b-t}{t-a}\right)\right]\right) \\ & \leq \frac{1}{b-a}\left[\pi T\left(\int_a^t \exp(\alpha x'(s)) ds\right)(a, b; t) - \left(\int_a^t \exp(\alpha x'(s)) ds\right) \ln\left(\frac{b-t}{t-a}\right)\right] \\ & \leq \frac{\exp a}{b-a}\left[b - \frac{1}{b-a}\left(\pi(Tx)(a, b; t) - x(t) \ln\left(\frac{b-t}{t-a}\right)\right)\right] \\ & \quad + \frac{\exp b}{b-a}\left[\frac{1}{b-a}\left(\pi(Tx)(a, b; t) - x(t) \ln\left(\frac{b-t}{t-a}\right)\right) - a\right] \end{aligned}$$

for any $t \in (a, b)$.

In particular,

$$\begin{aligned} (3.5) \quad & \exp\left(\frac{\alpha}{b-a}\left[\pi(Tx)\left(a, b; \frac{a+b}{2}\right)\right]\right) \\ & \leq \frac{1}{b-a}\left[\pi T\left(\int_a^b \exp(\alpha x'(s)) ds\right)\left(a, b; \frac{a+b}{2}\right)\right] \\ & \leq \frac{\exp a}{b-a}\left[b - \frac{\pi}{b-a}(Tx)\left(a, b; \frac{a+b}{2}\right)\right] + \frac{\exp b}{b-a}\left[\frac{\pi}{b-a}(Tx)\left(a, b; \frac{a+b}{2}\right) - a\right]. \end{aligned}$$

REFERENCES

- [1] N. M. Dragomir, S. S. Dragomir and P. M. Farrell, Some inequalities for the finite Hilbert transform. *Inequality Theory and Applications*. Vol. I, 113–122, Nova Sci. Publ., Huntington, NY, 2001.
- [2] N. M. Dragomir, S. S. Dragomir and P. M. Farrell, Approximating the finite Hilbert transform via trapezoid type inequalities. *Comput. Math. Appl.* **43** (2002), no. 10-11, 1359–1369.
- [3] N. M. Dragomir, S. S. Dragomir, P. M. Farrell and G. W. Baxter, On some new estimates of the finite Hilbert transform. *Libertas Math.* **22** (2002), 65–75.
- [4] N. M. Dragomir, S. S. Dragomir and P. M. Farrell and G. W. Baxter, A quadrature rule for the finite Hilbert transform via trapezoid type inequalities. *J. Appl. Math. Comput.* **13** (2003), no. 1-2, 67–84.
- [5] N. M. Dragomir, S. S. Dragomir and P. M. Farrell P. M.; Baxter, G. W. A quadrature rule for the finite Hilbert transform via midpoint type inequalities. *Fixed point theory and applications*. Vol. 5, 11–22, Nova Sci. Publ., Hauppauge, NY, 2004.
- [6] S. S. Dragomir, Inequalities for the Hilbert transform of functions whose derivatives are convex. *J. Korean Math. Soc.* **39** (2002), no. 5, 709–729.
- [7] S. S. Dragomir, Approximating the finite Hilbert transform via an Ostrowski type inequality for functions of bounded variation. *J. Inequal. Pure Appl. Math.* **3** (2002), no. 4, Article 51, 19 pp
- [8] S. S. Dragomir, Approximating the finite Hilbert transform via Ostrowski type inequalities for absolutely continuous functions. *Bull. Korean Math. Soc.* **39** (2002), no. 4, 543–559.
- [9] S. S. Dragomir, Some inequalities for the finite Hilbert transform of a product. *Commun. Korean Math. Soc.* **18** (2003), no. 1, 39–57.
- [10] S. S. Dragomir, Sharp error bounds of a quadrature rule with one multiple node for the finite Hilbert transform in some classes of continuous differentiable functions. *Taiwanese J. Math.* **9** (2005), no. 1, 95–109.
- [11] S. S. Dragomir, Inequalities and approximations for the Finite Hilbert transform: a survey of recent results, Preprint *RGMIA Res. Rep. Coll.* **21** (2018), Art. 30, 90 pp. [Online <http://rgmia.org/papers/v21/v21a30.pdf>].
- [12] S. S. Dragomir, Inequalities for the finite Hilbert transform of convex functions, Preprint *RGMIA Res. Rep. Coll.* **21** (2018), Art. 31.
- [13] F. D. Gakhov, *Boundary Value Problems* (English translation), Pergamon Press, Oxford, 1966.
- [14] W. Liu and X. Gao, Approximating the finite Hilbert transform via a companion of Ostrowski's inequality for function of bounded variation and applications. *Appl. Math. Comput.* **247** (2014), 373–385.
- [15] W. Liu, X. Gao and Y. Wen, Approximating the finite Hilbert transform via some companions of Ostrowski's inequalities. *Bull. Malays. Math. Sci. Soc.* **39** (2016), no. 4, 1499–1513.
- [16] W. Liu and N. Lu, Approximating the finite Hilbert transform via Simpson type inequalities and applications. *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* **77** (2015), no. 3, 107–122.
- [17] S. G. Mikhlin and S. Prössdorf, *Singular Integral Operators* (English translation), Springer Verlag, Berlin, 1986.
- [18] S. Wang, X. Gao and N. Lu, A quadrature formula in approximating the finite Hilbert transform via perturbed trapezoid type inequalities. *J. Comput. Anal. Appl.* **22** (2017), no. 2, 239–246.
- [19] S. Wang, N. Lu and X. Gao, A quadrature rule for the finite Hilbert transform via Simpson type inequalities and applications. *J. Comput. Anal. Appl.* **22** (2017), no. 2, 229–238.

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.org/dragomir>

²DST-NRF CENTRE OF EXCELLENCE IN THE MATHEMATICAL, AND STATISTICAL SCIENCES, SCHOOL OF COMPUTER SCIENCE, & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND,, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA.