

Certain fractional integral inequalities using generalized Katugampola fractional integral operator

Asha B.Nale¹, Satish K.Pachal², Vaijanath L.Chinchane³

^{1,2}Department of Mathematics,

Dr. Babasaheb Ambedkar Marathwada University,
Aurangabad-431 004, INDIA.

E-mail ID: ashabnale@gmail.com

E-mail ID: drskpanchal@gmail.com

³Department of Mathematics,

Deogiri Institute of Engineering and Management
Studies Aurangabad-431005, INDIA

E-mail ID: chinchane85@gmail.com

Abstract

The purpose of this paper is to obtain some new fractional integral inequalities involving convex functions by applying generalized Katugampola fractional integral operator.

Keywords : generalized Katugampola fractional integral, convex functions and inequality.

Mathematics Subject Classification : 26A99, 26D10.

1 Introduction

In the last years, a number of researchers have found inequalities and application by means results about fractional integrals such as Riemann-Liouville fractional integral operator, Hadamard integral operator, Saigo integral operator, Erdelyi-Kober integral operator, Katugampola fractional integral, see [1, 2, 3, 5, 7, 8, 9, 10, 11, 12, 15, 16, 17, 19, 21]. Fractional inequalities play major role in the development of fractional differential, integral equations and other fields of sciences and technology. Recently, In 2019, V.L. Chinchane and D.B. Pachpatte have studied inequalities using Saigo fractional integral, see [6].

Theorem 1.1 *Let f, h be two positive continuous functions on $[0, \infty)$ and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing, f is increasing on $[0, \infty)$ and for any convex function ϕ , $\phi(0) = 0$, then for $t > 0$, $\alpha > \max\{0, -\beta\}$, $\beta < 1$, $\beta - 1 < \eta < 0$, we have*

$$\frac{I_{0,t}^{\alpha,\beta,\eta}[f(t)]}{I_{0,t}^{\alpha,\beta,\eta}[h(t)]} \geq \frac{I_{0,t}^{\alpha,\beta,\eta}[\phi(f(t))]}{I_{0,t}^{\alpha,\beta,\eta}[\phi(h(t))]} \quad (1.1)$$

and

Theorem 1.2 *Let f, h be two positive continuous functions on $[0, \infty)$ and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing, f is increasing on $[0, \infty)$ and for any convex function ϕ , $\phi(0) = 0$, then we have inequality*

$$\frac{I_{0,t}^{\alpha,\beta,\eta}[f(t)] I_{0,t}^{\psi,\delta,\zeta}[\phi(h(t))] + I_{0,t}^{\psi,\delta,\zeta}[f(t)] I_{0,t}^{\alpha,\beta,\eta}[\phi(h(t))]}{I_{0,t}^{\alpha,\beta,\eta}[h(t)] I_{0,t}^{\psi,\delta,\zeta}[\phi(f(t))] + I_{0,t}^{\psi,\delta,\zeta}[h(t)] I_{0,t}^{\alpha,\beta,\eta}[\phi(f(t))]} \geq 1, \quad (1.2)$$

where for all $t > 0$, $\alpha > \max\{0, -\beta\}$, $\psi > \max\{0, -\delta\}$, $\beta < 1$, $\beta - 1 < \eta < 0$, $\delta < 1$, $\delta - 1 < \zeta < 0$.

In the literature, some fractional inequalities are obtain by using Generalized Katugampola fractional integral, see [1, 4, 14, 18]. Motivated by above work in this paper we have obtain some new inequalities using generalized Katugampola fractional integral for convex functions.

2 Preliminaries

Here, we devoted to basic concepts of Generalized Katugampola fractional integral, see [1, 13, 20].

Definition 2.1 *Consider the space $X_c^p(a, b)$ ($c \in \mathbb{R}, 1 \leq p \leq \infty$), of those complex valued Lebesgue measurable functions f on (a, b) for which the norm $\|f\|_{X_c^p} < \infty$, such that*

$$\|f\|_{X_c^p} = \left(\int_a^b |x^c f|^p \frac{dx}{x} \right)^{\frac{1}{p}}, \quad (1 \leq p < \infty)$$

and

$$\|f\|_{X_c^p} = \sup_{x \in (a,b)} [x^c |f|].$$

In particular, when $c = \frac{1}{p}$, the space $X_c^p(a, b)$ coincides with the space $L^p(a, b)$

Definition 2.2 *The left and right sided fractional integrals of a function f where $f \in X_c^p(a, b)$, $\alpha > 0$ and $\beta, \rho, \eta, k \in \mathbb{R}$, are defined respectively by*

$${}^\rho \mathcal{J}_{a+; \eta, k}^{\alpha, \beta} f(x) = \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_a^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} f(\tau) d\tau, \quad 0 \leq a < x < b \leq \infty, \quad (2.1)$$

and

$${}^\rho \mathcal{J}_{b-; \eta, k}^{\alpha, \beta} f(x) = \frac{\rho^{1-\beta} x^{k\rho}}{\Gamma(\alpha)} \int_x^b \frac{\tau^{k+\rho-1}}{(\tau^\rho - x^\rho)^{1-\alpha}} f(\tau) d\tau, \quad 0 \leq a < x < b \leq \infty, \quad (2.2)$$

if the integral exist.

To represent and discuss our new results in this paper we use the left sided fractional integrals, the right sided fractional can be proved similarly, also we consider $a = 0$, in (2.1), to obtain

$${}^{\rho}\mathcal{J}_{\eta,k}^{\alpha,\beta} f(x) = \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^{\rho} - \tau^{\rho})^{1-\alpha}} f(\tau) d\tau. \quad (2.3)$$

The above fractional integrals has the following composition (index) formulae

$$\begin{aligned} {}^{\rho}\mathcal{J}_{a+;\eta_1,k_1}^{\alpha_1,\beta_1} {}^{\rho}\mathcal{J}_{a+;\eta_2,-\rho\eta_1}^{\alpha_2,\beta_2} f(x) &= {}^{\rho}\mathcal{J}_{a+;\eta_2,k_1}^{\alpha_1+\alpha_2,\beta_1+\beta_2} f(x), \\ {}^{\rho}\mathcal{J}_{b-;\eta_1,-\rho\eta_2}^{\alpha_1,\beta_1} {}^{\rho}\mathcal{J}_{b-;\eta_2,k_2}^{\alpha_2,\beta_2} f(x) &= {}^{\rho}\mathcal{J}_{a+;\eta_1,k_2}^{\alpha_1+\alpha_2,\beta_1+\beta_2} f(x). \end{aligned} \quad (2.4)$$

For the convenience of establishing our results we define the following function as in [?]: let $x > 0, \alpha > 0, \rho, k, \beta, \eta \in \mathbb{R}$, then

$$\Lambda_{x,k}^{\rho,\beta}(\alpha, \eta) = \frac{\Gamma(\eta + 1)}{\Gamma(\eta + \alpha + 1)} \rho^{-\beta} x^{k+\rho(\eta+\alpha)}. \quad (2.5)$$

Remark 2.1 *The fractional integral (2.1) contain five well-known fractional integral as its particular cases, see [1, 13, 20]*

1. *Setting $k = 0, \eta = 0, a = 0$ and taking the limit $\rho \rightarrow 1$ in (2.1), the integral operator (2.1) reduces to the Riemann-Liouville fractional integral.*
2. *Setting $k = 0, \eta = 0$ and taking the limit $\rho \rightarrow 1$ in (2.1), the integral operator (2.1) reduces to the Liouville fractional integral.*
3. *Setting $\beta = \alpha, \eta = 0$ and taking the limit $\rho \rightarrow 0^+$ with L' Hospital rule in (2.1), the integral operator (2.1) reduces to the Hadamard fractional integral.*
4. *Setting $\beta = 0, k = -\rho(\alpha+\eta)$ in (2.1), the integral operator (2.1) reduces to the Erdelyi-Kober fractional integral.*
5. *Setting $\beta = \alpha, k = 0$ and $\eta = 0$ in (2.1), the integral operator (2.1) reduces to the Katugampola fractional integral.*

3 Fractional integral inequalities involving convex functions

In this section, we prove some fractional integral inequalities involving convex function using generalized Katugampola fractional integral,

Theorem 3.1 *Let p, r be two positive continuous functions on $[0, \infty)$ and $p \leq r$ on $[0, \infty)$. If $\frac{p}{r}$ is decreasing, p is increasing on $[0, \infty)$ and for any convex function Φ , $\Phi(0) = 0$, then for all $\alpha \geq 0$, $t > 0, \beta, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{{}_\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)]}{{}_\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)]} \geq \frac{{}_\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\Phi(p(t))]}{{}_\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\Phi(r(t))]} \quad (3.1)$$

Proof:- If the function Φ is convex with $\Phi(0) = 0$, then the function $\frac{\Phi(t)}{t}$ is increasing. Since p is increasing, then $\frac{\Phi(p(t))}{r(t)}$ is also increasing. Clearly $\frac{p(t)}{r(t)}$ is decreasing, for all $\tau, \sigma \in [0, \infty)$, and

$$\left(\frac{\Phi(p(\tau))}{p(\tau)} - \frac{\Phi(p(\sigma))}{p(\sigma)} \right) \left(\frac{p(\sigma)}{r(\sigma)} - \frac{p(\tau)}{r(\tau)} \right) \geq 0, \quad (3.2)$$

which implies that

$$\frac{\Phi(p(\tau))}{p(\tau)} \frac{p(\sigma)}{r(\sigma)} + \frac{\Phi(p(\sigma))}{p(\sigma)} \frac{p(\tau)}{r(\tau)} - \frac{\Phi(p(\tau))}{p(\tau)} \frac{p(\tau)}{r(\tau)} - \frac{\Phi(p(\sigma))}{p(\sigma)} \frac{p(\sigma)}{r(\sigma)} \geq 0. \quad (3.3)$$

Multiplying equation (3.3) by $r(\tau)r(\sigma)$, we have

$$\begin{aligned} \frac{\Phi(p(\tau))}{p(\tau)} p(\sigma)r(\tau) + \frac{\Phi(p(\sigma))}{p(\sigma)} p(\tau)r(\sigma) - \frac{\Phi(p(\tau))}{p(\tau)} p(\tau)r(\sigma) \\ - \frac{\Phi(p(\sigma))}{p(\sigma)} p(\sigma)r(\tau) \geq 0. \end{aligned} \quad (3.4)$$

Multiplying both sides of (3.4) by $\frac{\rho^{1-\beta} t^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(t^\rho - \tau^\rho)^{1-\alpha}}$, $\tau \in (0, t)$ which is positive, and integrating obtained result with respect to τ from 0 to t , we have

$$\begin{aligned} p(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(x)} r(t) \right] + \frac{\Phi(p(\sigma))}{p(\sigma)} r(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] \\ - r(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} p(t) \right] - \frac{\Phi(p(\sigma))}{p(\sigma)} p(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] \geq 0. \end{aligned} \quad (3.5)$$

Multiplying both sides of (3.5) by $\frac{\rho^{1-\beta} t^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(t^\rho - \sigma^\rho)^{1-\alpha}}$, $\sigma \in (0, t)$ which is positive, and integrating obtained result with respect to σ from 0 to t , we have

$$\begin{aligned} & [p(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right] + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] \\ & \geq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} p(t) \right] + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} p(t) \right] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)]. \end{aligned} \quad (3.6)$$

It follows that

$${}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)r(t)} \right] \geq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(f(t))}{p(t)} p(t) \right], \quad (3.7)$$

$$\frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)]} \geq \frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} p(t) \right]}}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right]}. \quad (3.8)$$

Since $p \leq r$ on $[0, \infty)$ and function $\frac{\Phi(t)}{t}$ is increasing, then for $\tau, \sigma \in [0, \infty)$, we have

$$\frac{\Phi(p(\tau))}{p(\tau)} \leq \frac{\Phi(r(\tau))}{r(\tau)}. \quad (3.9)$$

Multiplying (3.9) by $\frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} r(\tau)$, $\tau \in (0, t)$ which is positive, and integrating equation (3.9) on both side with respective τ from 0 to t , we get

$$\begin{aligned} & \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_a^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \frac{\Phi(p(\tau))}{p(\tau)} r(\tau) d\tau \leq \\ & \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_a^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \frac{\Phi(r(\tau))}{r(\tau)} r(\tau) d\tau, \end{aligned} \quad (3.10)$$

which implies that,

$${}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right] \leq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\Phi(r(t))}{r(t)} r(t) \right]. \quad (3.11)$$

Hence, from (3.8) and (3.11) we obtain required inequality (3.1).

Theorem 3.2 *Let p, r be two positive continuous functions on $[0, \infty)$ and $p \leq r$ on $[0, \infty)$. If $\frac{p}{r}$ is decreasing, p is increasing on $[0, \infty)$ and for any convex function Φ , $\Phi(0) = 0$, then for all $k \geq 0$, $t > 0$, $\alpha, \theta \geq 0$, $t > 0, \beta, \pi, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\phi(r(t))] + {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [p(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\Phi(r(t))]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\Phi(p(t))] + {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [r(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\Phi(p(t))]} \geq 1, \quad (3.12)$$

Proof: If function Φ is convex with $\Phi(0) = 0$, then $\frac{\Phi(t)}{t}$ is increasing. Since p is increasing, then $\frac{\Phi(p(t))}{p(t)}$ is also increasing. Clearly $\frac{p(t)}{r(t)}$ is decreasing, for all $\tau, \sigma \in [0, t]$ $t > 0$. Multiplying equation (3.5) by $\frac{\rho^{1-\pi} t^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(t^\rho - \sigma^\rho)^{1-\theta}}$, $\sigma \in (0, t)$, $t \in \mathbb{R}$, which remains positive. Now integrating obtained result with respect to σ from 0 to t , we have

$$\begin{aligned} & {}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} [p(t)] {}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} \left[\frac{\Phi(f(t))}{p(t)} r(t) \right] + {}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right] {}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [p(t)] \\ & \geq {}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [r(t)] {}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} [\Phi(p(t))] + {}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [\Phi(p(t))] {}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} [r(t)]. \end{aligned} \quad (3.13)$$

Since $p \leq r$ on $[0, \infty)$ and as function $\frac{\Phi(t)}{t}$ is increasing, for $\tau, \sigma \in [0, t]$ $t > 0$, we have

$$\frac{\Phi(p(\tau))}{p(\tau)} \leq \frac{\Phi(r(\tau))}{r(\tau)}. \quad (3.14)$$

Multiplying both sides of (3.14) by $\frac{\rho^{1-\pi} t^k}{\Gamma(\theta)} \frac{\tau^{\rho(\eta+1)-1}}{(t^\rho - \tau^\rho)^{1-\theta}}$, $\tau \in (0, t)$, $t \in \mathbb{R}$ positive, and integrating obtained result with respect to τ from 0 to t , we have

$${}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} \left[\frac{\Phi(p(t))}{p(t)} r(t) \right] \leq {}^\rho \mathcal{J}_{\eta, k}^{\theta, \pi} [\Phi(p(t))]. \quad (3.15)$$

Hence, using (3.13) and (3.15), we obtain required inequality (3.12).

Now, we prove our main result.

Theorem 3.3 *Let p , r and q be three positive continuous functions on $[0, \infty)$ and $p \leq r$ on $[0, \infty)$. If $\frac{p}{r}$ is decreasing, p and q are increasing functions on $[0, \infty)$, and for any convex function ϕ such that $\phi(0) = 0$, then for all $\alpha \geq 0$, $t > 0, \beta, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{{}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [p(t)]}{{}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [r(t)]} \geq \frac{{}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [\phi(p(t))q(t)]}{{}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [\phi(r(t))q(t)]}, \quad (3.16)$$

Proof: Since $p \leq r$ on $[0, \infty)$ and function $\frac{\phi(t)}{t}$ is increasing, then for $\tau, \sigma \in [0, t]$, $t > 0$, we have

$$\frac{\phi(p(\tau))}{p(\tau)} \leq \frac{\phi(r(\tau))}{r(\tau)}. \quad (3.17)$$

Multiplying both sides of (3.17) by $\frac{\sigma^{1-\beta} t^k}{\Gamma(\alpha)} \frac{\tau^{\sigma(\eta+1)-1}}{(t^\sigma - \tau^\sigma)^{1-\alpha}} r(\tau)$ positive, and integrating obtained result with respect to τ from 0 to t , we have

$${}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} \left[\frac{\phi(p(t))}{p(t)} r(t) q(t) \right] \leq {}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} [\phi(r(t)) q(t)]. \quad (3.18)$$

On the other hand, since the fact that the function ϕ is convex with $\phi(0) = 0$. Then the function $\frac{\phi(t)}{t}$ is increasing. Since p is increasing, $\frac{\phi(p(t))}{p(t)}$ is also increasing. Clearly we can say that $\frac{p(t)}{r(t)}$ is decreasing, for all $\tau, \sigma \in [0, t]$ $t > 0$

$$\left(\frac{\phi(p(\tau))}{p(\tau)}q(\tau) - \frac{\phi(p(\sigma))}{p(\sigma)}q(\sigma) \right) (p(\sigma)r(\tau) - p(\tau)r(\sigma)) \geq 0, \quad (3.19)$$

which implies that

$$\begin{aligned} \frac{\phi(p(\tau))q(\tau)}{p(\tau)}p(\sigma)r(\tau) + \frac{\phi(p(\sigma))q(\sigma)}{p(\sigma)}p(\tau)r(\sigma) - \frac{\phi(p(\tau))q(\tau)}{p(\tau)}p(\tau)r(\sigma) \\ - \frac{\phi(p(\sigma))q(\sigma)}{p(\sigma)}p(\sigma)r(\tau) \geq 0. \end{aligned} \quad (3.20)$$

Hence, we can write

$$\begin{aligned} p(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\phi(p(t))}{p(t)}r(t)q(t) \right] + \frac{\phi(p(\sigma))}{p(\sigma)}r(\sigma)q(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] \\ - r(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(p(t))q(t)] - \frac{\phi(p(\sigma))}{p(\sigma)}p(\sigma)q(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] \geq 0, \end{aligned} \quad (3.21)$$

with the same argument as before, we have

$$\frac{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)]}{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)]} \geq \frac{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(p(t))q(t)]}{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\phi(p(t))}{p(t)}r(t)q(t) \right]}. \quad (3.22)$$

Hence, using equation (3.18) and (3.22), we obtain (3.16).

Now, we give generalization of Theorem 3.3.

Theorem 3.4 *Let p, r and q be three positive continuous functions on $[0, \infty)$ and $p \leq r$ on $[0, \infty)$. If $\frac{p}{r}$ is decreasing, p and q are increasing functions on $[0, \infty)$, and for any convex function ϕ such that $\phi(0) = 0$, then for all $k \geq 0$, $t > 0$, $\alpha, \theta \geq 0$, $t > 0, \beta, \pi, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\phi(r(t))q(t)] + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [p(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(r(t))q(t)]}{\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\phi(p(t))q(t)] + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [r(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(p(t))q(t)]} \geq 1. \quad (3.23)$$

Proof:- Multiplying equation (3.21) by $\frac{\rho^{1-\pi} t^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(t^\rho - \sigma^\rho)^{1-\theta}}$, $\sigma \in (0, t)$, $t \in \mathbb{R}$, which remains positive. Then integrate the resulting identity with respect to σ from 0 to t , we have

$$\begin{aligned} & {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [p(t)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\phi(p(t))q(t)}{p(t)} r(t) \right] + {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} \left[\frac{\phi(p(t))q(t)}{p(t)} r(t) \right] \times \\ & {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p(t)] \geq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [r(t)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\phi(p(t))q(t)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(p(t))q(t)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [r(t)], \end{aligned} \quad (3.24)$$

and since $p \leq r$ on $[0, \infty)$ and use the fact that $\frac{\phi(t)q(t)}{t}$ is increasing, we obtain

$${}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} \left[\frac{\phi(p(t))q(t)}{p(t)} r(t) \right] \leq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [\phi(r(t))q(t)], \quad (3.25)$$

and

$${}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} \left[\frac{\phi(p(t))q(t)}{p(t)} r(t) \right] \leq {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [\phi(r(t))q(t)]. \quad (3.26)$$

Hence, from equation (3.24), (3.25) and (3.26), we obtain (3.23).

4 Other fractional integral inequalities

In [9], authors have proved the inequalities using Riemann-Liouville fractional integral. Now, we prove the similar results using generalized Katugampola fractional integral

Theorem 4.1 *Let p, q be two positive and continuous functions on $[0, \infty)$ such that p is decreasing and q is increasing on $[0, \infty)$, then for all $\alpha \geq 0$, $t > 0, \beta, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p^l(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [p^m(t)]} \geq \frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [q^n p^l(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [q^n p^m(t)]}. \quad (4.1)$$

Proof:- Consider $\sigma, \tau \in (0, t)$, we have

$$(q^n(\sigma) - q^n(\tau)) (p^l(\tau)p^m(\sigma) - p^m(\tau)p^l(\sigma)) \geq 0,$$

which implies that

$$q^n(\sigma)p^l(\tau)p^m(\sigma) + q^m(\tau)p^m(\tau)p^l(\sigma) \geq q^m(\sigma)p^m(\tau)p^l(\sigma) + q^m(\tau)p^m(\sigma)p^l(\tau), \quad (4.2)$$

Multiplying both sides of (4.2) by $\frac{\rho^{1-\beta}t^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(t^\rho-\tau^\rho)^{1-\alpha}}$ which is positive, and integrating obtained result with respect to τ from 0 to t , we have

$$q^n(\sigma)p^m(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^l(t)] + p^l(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^m(t)] \geq g^r(\sigma) f^p(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[f^q(t)] + p^m(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^l(t)]. \quad (4.3)$$

Now, multiplying both side of (4.3) by $\frac{\rho^{1-\beta}t^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(t^\rho-\sigma^\rho)^{1-\alpha}}$ which is positive from (2.4). Now integrating obtained result with respect to σ from 0 to t , we have

$${}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^m(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^l(t)] \geq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^l(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^m(t)], \quad (4.4)$$

which gives the inequality 4.1.

Theorem 4.2 *Let p, q be two positive and continuous functions on $[0, \infty)$ such that p is decreasing and q is increasing on $[0, \infty)$. then for all $k \geq 0, t > 0, \alpha, \theta \geq 0, t > 0, \beta, \pi, \rho, \eta, k \in \mathbb{R}$ we have,*

$$\frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^l(t)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi}[q^n p^m(t)] + {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi}[p^l(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^m(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^m(t)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi}[q^n p^m(t)] + {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi}[p^m(t)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^m(t)]} \geq 1. \quad (4.5)$$

Proof:- Multiplying equation (4.3) by $\frac{\rho^{1-\pi}t^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(t^\rho-\sigma^\rho)^{1-\theta}}$ ($\sigma \in (0, t), t > 0$), which remains positive. Then integrate the resulting identity with respect to σ from 0 to t , we obtain the result 4.5.

Theorem 4.3 *Let p, q be two positive and continuous functions on $[0, \infty)$ such that p is decreasing and q is increasing on $[0, \infty)$, Such that*

$$(p^n(\tau)q^n(\sigma) - p^n(\sigma)q^n(\tau)) (p^{l-n}(\tau) - p^{l-n}(\sigma)) \geq 0,$$

then for all $\alpha \geq 0, t > 0, \beta, \sigma, \eta, k \in \mathbb{R}$ we have,

$$\frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^{n+l}(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[p^{n+m}(t)]} \geq \frac{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^l(t)]}{{}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[q^n p^m(t)]}, \quad (4.6)$$

$n > 0$.

Proof:- Consider $\tau, \sigma \in (0, t)$, we get

$$(p^n(\tau)q^n(\sigma) - p^n(\sigma)q^n(\tau)) (p^m(\sigma)p^l(\tau) - p^m(\tau)p^l(\sigma)) \geq 0.$$

and using the same arguments as in Theorem [4.1]. we obtain the result

References

- [1] T.A Aljaaidi and D. B. Pachpatte, *Osme Grüss-type inequalities using generalized Katugampola fractional integral*, AIMS Mathematics,5(2),(2020),1011-1024.
- [2] S. Belarbi and Z. Dahmani, *On some new fractional integral inequality*, J. Inequal. Pure and Appl. Math. Art.86, 10(3), (2009), 1-5.
- [3] D. Baleanu, S. D. Purohit and J. C. Prajapati, *Integral inequalities involving generalized Erdélyi-Kober fractional integral operators*, Open. Math. 14, (2016), 89-99.
- [4] V. L. Chinchane, *New approach to Minkowski fractional inequalities using generalized K-fractional integral operator*, Journal of the Indian. Math. Soc., 1-2(85),(2018) 32-41.
- [5] V. L. Chinchane and D. B. Pachpatte, *On some Grüss-type fractional inequalities using Saigo fractional integral operator*, Journal of Mathematics, Article ID 527910, Vol.2014 (2014), 1-9.
- [6] V. L. Chinchane and D. B. Pachpatte, *Note on fractional integral inequality involving convex function using Saigo fractional integral*, Indian Journal of Mathematics, Vol.1(61),(2019),27-39.
- [7] V. L. Chinchane and D. B. Pachpatte, *On new fractional integral inequalities involving convex functions using Hadamard fractional integral*, Indian Journal of Mathematics, Vol.2(31),(2016),183-192.
- [8] Z. Dahmani, *A note on some new fractional results involving convex functions*, Acta Math. Univ. Comenianae, 81(2),(2012), 241-246.
- [9] Z. Dahmani and N. Bedjaoui, *Some generalized integral inequalities*, J. Advan. Res. Appl. Math, 3(4),(2011), 58-66.
- [10] Z. Dahmani and H. Metakkel El Ard, *Generalization of some integral inequalities using Riemann-Liouville operator*, Int. J. Open Problem Compt. Math., 4(4), (2011), 40-46.
- [11] A. A. George, *Fractional Differentiation Inequalities*, Springer Publishing Company, New York, 2009.
- [12] U. N. Katugampola, *A new approach to generalized fractional derivatives*, Bull. Math. Anal. Appl., 6(4), (2014), 1-15.

- [13] U. N. Katugampola, *new fractional integral unifying six existing fractional integral*, arXiv:1612.08596 [math.CA] (2016), 1-6.
- [14] S. Kilinc and H.Yildirim, *Generalized fractional integral inequalities involving Hypergeometric operators*, Int. J. Pure Appl. Math., 101(1), (2015), 71-82.
- [15] V. Kiryakova, *On two Saigo's fractional integral operator in the class of univalent functions*, Fract. Calc. Appl. Anal., 9(2),(2006).
- [16] A. R. Prabhakaran and K. Srinivasa Rao, *Saigo operator of fractional integration of Hypergeometric functions*, Int. J. Pure Appl. Math., 81(5), (2012), 755-763.
- [17] S. D. Purohit and R. K. Raina, *Chebyshev type inequalities for the Saigo fractional integral and their q - analogues*, J. Math. Inequal., 7(2),(2013), 239-249.
- [18] S. D. Purohit and R. K. Raina, *Certain fractional integral inequalities involving the Gauss hypergeometric function*, Rev. Tec. Ing. Univ. Zulia 37(2), (2014), 167-175.
- [19] M. Saigo, *A remark on integral operators involving the Grüss hypergeometric functions*, Math. Rep. Kyushu Univ., 11(1978), 135-143.
- [20] J. Vanterler da C. Sousa D. S. Oliveira E. Capelas de Oliveira *Grüss-type inequalities by means of generalized fractional integrals*, Bull Braz. Math. Soc. New 50,(2019). 1029-1047.
- [21] S.G.Somko, A.A.Kilbas and O.I.Marichev, *Fractional Integral and Derivative Theory and Application*, Gordon and Breach, Yverdon, Switzerland, 1993.