

INEQUALITIES AND BOUNDS FOR THE p -GENERALIZED TRIGONOMETRIC FUNCTIONS

LI YIN AND XIU-LI LIN

ABSTRACT. In this paper, we mainly show some bounds and inequalities for the p -generalized trigonometric functions by using classical analysis.

1. INTRODUCTION

It is well known that the level sets of the functions

$$T_p(x, y) = (|x|^p + |y|^p)^{\frac{1}{p}}, (x, y) \in \mathbb{R}^2, p > 0$$

can be easily described by the equation $r_p = c$ if one makes use of polar or standard triangle coordinates in the cases $p = 2$ and $p = 1$, respectively. In [16], Richter considered r_1 and r_2 as special cases of the p -generalized radius coordinate $r_p = (|x|^p + |y|^p)^{\frac{1}{p}}, p > 0$ and the functions $\sin \varphi$, $\cos \varphi$ as special cases of certain p -generalized trigonometric functions. In fact, Richter dealt with $l_{n,p}$ -spherical and simplicial coordinates. Here, we only discussed the $l_{2,p}$ -generalized trigonometric functions (or called p -generalized trigonometric functions) as a generalization to classical trigonometric functions. Next, we show some definitions and formulas found by Richter. For the details, the reader may refer to references ([16, 17]).

Definition 1.1. *The p -generalized sine and cosine values of an angle $\varphi \in (0, 2\pi)$ between the directions of the positive x -axes and the line through the points $(0, 0)$ and $(x, y) \in \mathbb{R}^2$ are defined for each $p > 0$ as*

$$\sin_p \varphi = \frac{y}{(|x|^p + |y|^p)^{\frac{1}{p}}}$$

and

$$\cos_p \varphi = \frac{x}{(|x|^p + |y|^p)^{\frac{1}{p}}}.$$

Obviously, it holds $|\sin_p \varphi| \leq 1$, $|\cos_p \varphi| \leq 1$ and

$$|\sin_p \varphi|^p + |\cos_p \varphi|^p = 1.$$

It is easy to see that for each $p > 0, \varphi \in (0, 2\pi)$,

$$\sin_p \varphi = \frac{\sin \varphi}{N_p \varphi} \tag{1.1}$$

2010 *Mathematics Subject Classification.* 33B10.

Key words and phrases. p -generalized trigonometric functions; Turán type inequalities; inequality.

This work was jointly supported by Natural Foundation of Shandong Province (Grant No. ZR2018MF023), and by the Science Foundation of Binzhou University (Grant No. BZXLYG1903 and BZXYL1506).

and

$$\cos_p \varphi = \frac{\cos \varphi}{N_p \varphi} \quad (1.2)$$

where $N_p \varphi = (|\sin \varphi|^p + |\cos \varphi|^p)^{\frac{1}{p}}$. For $\varphi \neq \frac{k\pi}{2}, k = 1, 2, 3$, the first derivatives of \sin_p and \cos_p are

$$\sin'_p \varphi = \cos_p \varphi \frac{|\cos \varphi|^{p-2}}{(N_p \varphi)^p} \quad (1.3)$$

and

$$\cos'_p \varphi = -\sin_p \varphi \frac{|\sin \varphi|^{p-2}}{(N_p \varphi)^p}. \quad (1.4)$$

In particular, for $p > 0$ and $x \in (0, \frac{\pi}{2})$, we have

$$\sin'_p x = \frac{(\cos x)^{p-1}}{((\sin x)^p + (\cos x)^p)^{\frac{p+1}{p}}}, \quad (1.5)$$

$$\cos'_p x = -\frac{(\sin x)^{p-1}}{((\sin x)^p + (\cos x)^p)^{\frac{p+1}{p}}} \quad (1.6)$$

and

$$\frac{\sin'_p x}{\cos'_p x} = -(\cot x)^{p-1}. \quad (1.7)$$

The generalized tangent function $\tan_p x$ is defined as

$$\tan_p x = \frac{\sin_p x}{\cos_p x}, \quad x \in \mathbb{R} \setminus \left\{ k\pi + \frac{\pi}{2} : k \in \mathbb{Z} \right\}. \quad (1.8)$$

It is obvious that $\tan_p x = \tan x$. From 1.8, it follows that

$$\tan'_p x = 1 + |\tan x|^2, \quad x \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right). \quad (1.9)$$

It is clear that all these generalized functions coincide with the classical ones when $p = 2$.

In this paper, we mainly show some bounds and classical inequalities and Turán type inequalities to parameters for the p -generalized trigonometric functions. This paper presents an attempt to study these functions, although many of the results are not perfect. We hope to provide some basic conclusions and research directions for interested researchers.

2. TURÁN TYPE INEQUALITIES

Lemma 2.1 ([14, Lemma 1]). *The two variable power mean $M_p(a, b)$ is concave in p for $p \geq 1$ and convex in p for $p \leq -1$. That is, $\frac{\partial^2}{\partial p^2} [M_p(a, b)] \leq 0$ for $p \geq 1$ and $\frac{\partial^2}{\partial p^2} [M_p(a, b)] \geq 0$ for $p \leq -1$ with equality if and only if $a = b$.*

Theorem 2.1. *For fixed $x \in (0, \frac{\pi}{2})$ and $k > 1$, the function $p \mapsto 2^{\frac{k}{p}} \sin_p x$ is log-convex on $p \in [1, \infty)$. In particular, for fixed $x \in (0, \frac{\pi}{2})$ and $k > 1, p, q \geq 1$, we have*

$$\sin_{\frac{p+q}{2}} x \leq 2^{\frac{k}{2p} + \frac{k}{2q} - \frac{2k}{p+q}} \sin_p^{\frac{1}{2}} x \sin_q^{\frac{1}{2}} x. \quad (2.1)$$

Proof. By using the formula 1.1, we easily obtain

$$2^{\frac{k}{p}} \sin_p x = \frac{2^{\frac{k}{p}} \sin x}{(\sin^p x + \cos^p x)^{\frac{1}{p}}} = \frac{2^{\frac{k-1}{p}}}{\left(\frac{1+\cot^p x}{2}\right)^{\frac{1}{p}}} = \frac{2^{\frac{k-1}{p}}}{M_p(1, \lambda)}$$

and

$$\log \left(2^{\frac{k}{p}} \sin_p x \right) = \frac{k-1}{p} \log 2 - \log M_p(1, \lambda)$$

where $\lambda = \cot x$. On one hand, we have

$$\frac{d}{dp} \left(\frac{k-1}{p} \log 2 \right) = -\frac{k-1}{p^2} \log 2$$

and

$$\frac{d^2}{dp^2} \left(\frac{k-1}{p} \log 2 \right) = \frac{2(k-1)}{p^3} \log 2 > 0.$$

This implies the function $2^{\frac{k-1}{p}}$ is strictly log-convex for $p \geq 1$ and $k > 1$. On the other hand, simple computation yields

$$[M_{pt}(1, \lambda)]^t = M_p(1, \lambda^t),$$

and

$$t \log (M_{pt}(1, \lambda)) = \log (M_p(1, \lambda^t)). \quad (2.2)$$

Differentiate (2.2) with respect to p while holding λ fixed, we get

$$t^2 [\log (M_{pt}(1, \lambda))] = [\log (M_p(1, \lambda^t))]'$$

and

$$t^3 [\log (M_{pt}(1, \lambda))]'' = [\log (M_p(1, \lambda^t))]''.$$

Putting $p = 1$, we have

$$t^3 [\log (M_t(1, \lambda))]'' \leq 0$$

by using Lemma 2.1 and the fact that the concave function meant log-concave function. So it is easy to see that the function $2^{\frac{k}{p}} \sin_p x$ is log-convex on $p \in [1, \infty)$. This completes the proof. \square

Corollary 2.1. For fixed $x \in (0, \frac{\pi}{2})$, the function $p \mapsto 2^{\frac{1}{p}} \sin_p x$ is log-convex on $p \in [1, \infty)$. In particular, for fixed $x \in (0, \frac{\pi}{2})$ and $p, q \geq 1$, we have

$$\sin_{\frac{p+q}{2}} x \leq 2^{\frac{1}{2p} + \frac{1}{2q} - \frac{2}{p+q}} \sin_{\frac{1}{2}} x \sin_{\frac{1}{2}} x. \quad (2.3)$$

Proof. By taking $k = 1$, we easily complete the proof. For the sake of simplicity, we omit the details. \square

Completely similar to proof of Theorem 2.1, we easily obtain the following Theorem 2.2.

Theorem 2.2. For fixed $x \in (0, \frac{\pi}{2})$ and $k > 1$, the function $p \mapsto 2^{\frac{k}{p}} \cos_p x$ is log-convex on $p \in [1, \infty)$. In particular, for fixed $x \in (0, \frac{\pi}{2})$ and $k > 1, p, q \geq 1$, we have

$$\cos_{\frac{p+q}{2}} x \leq 2^{\frac{k}{2p} + \frac{k}{2q} - \frac{2k}{p+q}} \cos_{\frac{1}{2}} x \cos_{\frac{1}{2}} x. \quad (2.4)$$

Corollary 2.2. For fixed $x \in (0, \frac{\pi}{2})$, the function $p \mapsto 2^{\frac{1}{p}} \cos_p x$ is log-convex on $p \in [1, \infty)$. In particular, for fixed $x \in (0, \frac{\pi}{2})$ and $p, q \geq 1$, we have

$$\cos_{\frac{p+q}{2}} x \leq 2^{\frac{1}{2p} + \frac{1}{2q} - \frac{2}{p+q}} \cos_{\frac{1}{p}} x \cos_{\frac{1}{q}} x. \quad (2.5)$$

3. BOUNDS FOR THE p -GENERALIZED TRIGONOMETRIC FUNCTIONS

Lemma 3.1 ([12]). Let $x \in (0, \frac{\pi}{2})$ and $n \geq 2$. Then

$$\sin^n x + \cos^n x \leq 1. \quad (3.1)$$

Lemma 3.2 ([12, c_p -inequality]). Let $a, b \in \mathbb{R}$ and $p > 0$. Then

$$(|a| + |b|)^p \leq c_p (|a|^p + |b|^p) \quad (3.2)$$

where $c_p = \begin{cases} 1, & 0 < p \leq 1, \\ 2^{p-1}, & p > 1. \end{cases}$

Lemma 3.3 ([12]). For $x \in (0, 1)$, we have

$$x < \tan x < \frac{x}{\sqrt{1-x^2}}.$$

Theorem 3.1. For fixed $x \in (0, \frac{\pi}{2})$ and $p \geq 2$, the following inequalities hold true:

- (i) $\sin x \leq \sin_p x \leq 2^{\frac{p-1}{p}} \sin_1 x$;
- (ii) $\cos x \leq \cos_p x \leq 2^{\frac{p-1}{p}} \cos_1 x$.

Proof. By using Lemma 3.1, Lemma 3.2 and definitions of the functions $\sin_p x$, $\cos_p x$, we have

$$\sin x \leq \sin_p x \leq \frac{\sin x}{2^{\frac{1-p}{p}} (\sin x + \cos x)} = 2^{\frac{p-1}{p}} \sin_1 x$$

and

$$\cos x \leq \cos_p x \leq \frac{\cos x}{2^{\frac{1-p}{p}} (\sin x + \cos x)} = 2^{\frac{p-1}{p}} \cos_1 x. \quad \square$$

Theorem 3.2. For fixed $x \in (0, \frac{\pi}{2})$ and $p \geq 2$, the following inequalities hold true:

- (i) $(\cos x)^{p-1} \leq \sin'_p x \leq 2^{\frac{p^2-1}{p}} \frac{(\cos x)^{p-1}}{(\sin x + \cos x)^{p+1}}$;
- (ii) $-2^{\frac{p^2-1}{p}} \frac{(\sin x)^{p-1}}{(\sin x + \cos x)^{p+1}} \leq \cos'_p x \leq -(\sin x)^{p-1}$.

Proof. By using Lemma 3.1, Lemma 3.2 and the formulas (1.5), (1.6), we easily complete the proof. □

Theorem 3.3. For $x \in (0, \frac{\pi}{2})$ and $k, l > 0$, we have

$$0 \leq \sin_p^k x \cos_p^l x \leq \left(\frac{k^k l^l}{(k+l)^{k+l}} \right)^{\frac{1}{p}}. \quad (3.3)$$

Proof. Since the function $\ln x$ is strictly concave on $(0, \infty)$, we take $\alpha = \frac{k}{l+k}, \beta = \frac{l}{l+k}$, $x_1 = \frac{1}{k} \sin_p^p x$ and $x_2 = \frac{1}{l} \cos_p^l x$ in Jessen inequality

$$\alpha \ln x_1 + \beta \ln x_2 \leq \ln(\alpha x_1 + \beta x_2).$$

This implies the inequality (3.3). □

Theorem 3.4. For $x \in (0, \frac{\pi}{2})$, the following inequalities hold true:

- (i) $(\sin_p x)^{\cos_p x} \leq (\cos_p x)^{\sin_p x}$, for $x \in (0, \frac{\pi}{4})$;
- (ii) $(\cos_p x)^{\sin_p x} \leq (\sin_p x)^{\cos_p x}$, for $x \in (\frac{\pi}{4}, \frac{\pi}{2})$;
- (iii) $(\sin_p x)^{\sin_p x} \leq (\cos_p x)^{\cos_p x}$, for $x \in (0, \frac{\pi}{4})$.

Proof. Considering to $\frac{\ln x}{x}$ is strictly increasing on $(0, \infty)$ and the inequalities

$$x \in \left(0, \frac{\pi}{4}\right), \sin_p x \leq \cos_p x,$$

$$x \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right), \cos_p x \leq \sin_p x,$$

we easily obtain proofs of (i) and (ii).

Due to (iii), the formula $(x \ln x)' = 1 + \ln x > 0$ for $x \in (0, \frac{\sqrt{2}}{2})$ implies that the function $x \ln x$ is strictly increasing. So, we have

$$\sin_p x \ln(\sin_p x) \leq \cos_p x \ln(\cos_p x).$$

The proof is complete. \square

Theorem 3.5. For fixed $x \in (0, \frac{\pi}{2})$ and $p \geq 0$, we have

$$(\sin_p x)^{p \sin_p^2 x} + (\cos_p x)^{p \cos_p^2 x} \geq \sqrt{2}. \quad (3.4)$$

Proof. The convexity of function x^x implies the above inequality. we omit the detail for the sake of simplicity. \square

Theorem 3.6. For fixed $x \in (0, 1)$ and $p \geq 0$, we have

$$\frac{\sqrt{1-x^2}}{\left[x^p + (\sqrt{1-x^2})^p\right]^{\frac{1}{p}}} < \cos_p x < \frac{1}{(1+x^p)^{\frac{1}{p}}}. \quad (3.5)$$

Proof. By using Lemma 3.3 and the formula $\frac{1}{\cos_p^2 x} - 1 = \tan^2 x$, we easily complete the proof. \square

4. CLASSICAL INEQUALITIES

Lemma 4.1 ([4, Lemma 3, p246]). Let us consider the function $f : (a, \infty) \rightarrow \mathbb{R}$, where $a \geq 0$. If the function g , defined by $g(x) = \frac{1}{x}(f(x) - 1)$, is increasing on (a, ∞) , then for the function h , defined by $h(x) = f(x^2)$, we have the following Grünbaum type inequality

$$1 + h(z) \geq h(x) + h(y), \quad (4.1)$$

where $x, y \geq a$ and $z^2 = x^2 + y^2$. If the function g is decreasing, then the inequality (2.1) is reversed.

Theorem 4.1. For fixed $x, y, z \in (0, \frac{\pi}{2})$ and $x^2 + y^2 = z^2$, we have

$$z^2 \sin_p(z^2) \geq x^2 \sin_p(x^2) + y^2 \sin_p(y^2) \quad (4.2)$$

and

$$z^2 \cos_p(z^2) \geq x^2 \cos_p(x^2) + y^2 \cos_p(y^2). \quad (4.3)$$

Proof. We only prove the inequality (4.2). Proof of inequality (4.3) is completely analogous to inequality (4.2). In fact, putting $f(x) = x \sin_p(x) + 1$, we get the function $g(x) = \frac{f(x)-1}{x} = \sin_p(x)$ is increasing on $(0, \frac{\pi}{2})$. by using Lemma (4.1), we easily obtain inequality (4.2). \square

Lemma 4.2 ([15, Mitrinović-Adamović inequality]). *Let $x \in (0, \frac{\pi}{2})$. Then*

$$\left(\frac{\sin x}{x}\right)^3 \geq \cos x.$$

Theorem 4.2. *For $p > 0$ and $x \in (0, \frac{\pi}{2})$, we have*

$$\frac{\sin_p^3 x}{x^3} \geq \frac{\cos_p^3 x}{\cos^2 x}. \quad (4.4)$$

Proof. Using the definition of $\sin_p x$, $\cos_p x$ and Lemma 4.1, we have

$$\begin{aligned} \frac{\sin_p^3 x}{x^3} &= \frac{\sin^3 x}{x^3 (\sin^p x + \cos^p x)^{\frac{3}{p}}} \\ &\geq \left[\frac{\cos x}{(\sin^p x + \cos^p x)^{\frac{1}{p}}} \right]^3 \frac{1}{\cos^2 x} = \frac{\cos_p^3 x}{\cos^2 x}. \end{aligned}$$

This completes the proof. \square

The above theorem generalized Mitrinović-Adamović inequality. The next theorem shows the Huugens-type inequality for the p -generalized trigonometric functions.

Theorem 4.3. *For $p > 0$ and $x \in (0, \frac{\pi}{2})$, we have*

$$\frac{3 \sin_p x}{x} + \frac{\cos^2 x}{\cos_p^3 x} > 4. \quad (4.5)$$

Proof. The weighted AG inequality shows that

$$ta + (1-t)b > a^t b^{1-t}$$

for $a > 0, b > 0$ and $0 < t < 1$. Taking $t = \frac{3}{4}$, $a = \frac{\sin_p x}{x}$ and $b = \frac{\cos^2 x}{\cos_p^3 x}$ and applying Theorem 4.3, we get

$$\frac{3 \sin_p x}{4 x} + \frac{1 \cos^2 x}{4 \cos_p^3 x} > \left(\frac{\sin_p x}{x}\right)^{\frac{3}{4}} \left(\frac{\cos^2 x}{\cos_p^3 x}\right)^{\frac{1}{4}} \geq 1.$$

So, we complete the proof. \square

Theorem 4.4. *For $t, p > 0$ and $x \in (0, \frac{\pi}{2})$, the following inequalities hold true:*

- (i) $\left[1 + \left(\frac{\sin_p x}{x}\right)^{2t}\right] \left[1 + \left(\frac{\tan_p x}{x}\right)^t\right] > 4;$
- (ii) $\left(\frac{\sin_p x}{x}\right)^{2t} + \left(\frac{\tan_p x}{x}\right)^t \geq 2 \sqrt{\left[1 + \left(\frac{\sin_p x}{x}\right)^{2t}\right] \left[1 + \left(\frac{\tan_p x}{x}\right)^t\right]} - 2 > 2;$
- (iii) $\left[1 + \left(\frac{\sin_p x}{x}\right)^t\right]^2 \left[1 + \left(\frac{\tan_p x}{x}\right)^t\right] > 8;$
- (iiii) $2 \left(\frac{\sin_p x}{x}\right)^t + \left(\frac{\tan_p x}{x}\right)^t \geq 3 \sqrt[3]{\left[1 + \left(\frac{\sin_p x}{x}\right)^t\right] \left[1 + \left(\frac{\tan_p x}{x}\right)^t\right]} - 3 > 3.$

Proof. By using the known inequality (See the reference [18])

$$a + b \geq 2\sqrt{(1+a)(1+b)} - 2 > 2\sqrt{ab}$$

and taking $a = \left(\frac{\sin_p x}{x}\right)^{2t}$, $b = \left(\frac{\tan_p x}{x}\right)^t$ and applying Theorem 3.1, Theorem 4.2, we can obtain (i) and (ii).

Completely similar, by using formula (See the reference [18])

$$2a + b \geq 3\sqrt[3]{(1+a)^2(1+b)} - 3 > 3\sqrt[3]{a^2b}$$

and putting $a = \left(\frac{\sin_p x}{x}\right)^{2t}$, $b = \left(\frac{\tan_p x}{x}\right)^t$, we easily obtain the expected results (iii) and (iiii). \square

Remark 4.1. Inequality (i) of Theorem 4.4 could be called multiplicative Wilker inequality, and likewise inequality (iii) of Theorem 4.4 as a multiplicative Huygens inequality.

5. COMMENTS AND OPEN PROBLEMS

The another generalized trigonometric and hyperbolic functions depending on a parameter $p > 1$ were studied by P. Lindqvist in a highly cited paper (see [13]). Motivated by this work, many authors have studied the equalities and inequalities related to generalized trigonometric and hyperbolic functions in [5, 6, 7]. Recently, in [19], S. Takeuchi has investigated the (p, q) -trigonometric functions depending on two parameters and in which the case of $p = q$ coincides with the p -function of Lindqvist, and for $p = q = 2$ they coincide with familiar elementary functions. These functions are defined differently from the above p -trigonometric functions, they are all defined by integration. These functions have been thoroughly studied, such as the multiple angle formula, classical inequalities, monotonicity of parameters, relations with hypergeometric functions, and generalized elliptic integrals defined by these functions. The reader may refer references [1, 2, 3, 8, 9, 10, 11, 20, 21, 22, 23, 24]. As an example, we introduce generalized trigonometric function with one parameter to illustrate the difference between them.

For $1 < q < \infty$ and $0 \leq x \leq 1$, the arcsine may be generalized as

$$\arcsin_q x = \int_0^x \frac{1}{(1-t^q)^{1/q}} dt$$

and

$$\frac{\pi_q}{2} = \arcsin_q 1 = \int_0^1 \frac{1}{(1-t^q)^{1/q}} dt.$$

The inverse of \arcsin_q on $[0, \frac{\pi_q}{2}]$ is called the generalized sine function, denoted by \sin_q and may be extended to $(-\infty, \infty)$. See [7] and closely related references therein.

For $x \in [0, \frac{\pi_q}{2}]$, the generalized cosine function $\cos_q x$ is defined by

$$\cos_q x = \frac{d \sin_q x}{dx}.$$

It is easy to see that

$$\cos_q x = (1 - \sin_q^q x)^{1/q}$$

and

$$\frac{d \cos_q x}{dx} = -\cos_q^{2-q} x \sin_q^{q-1} x.$$

Very naturally, we can define the following generalized elliptic integrals. We repeat the definition of complete q -elliptic integrals of the first kind $K_q(k)$ and of the second kind $E_q(k)$: for $k \in (0, 1)$

$$K_q(k) := \int_0^{\frac{\pi_q}{2}} \frac{d\theta}{(1 - k^q \sin_q^q \theta)^{1 - \frac{1}{q}}} = \int_0^1 \frac{dt}{(1 - t^q)^{\frac{1}{q}} (1 - k^q t^q)^{1 - \frac{1}{q}}},$$

$$E_q(k) := \int_0^{\frac{\pi_q}{2}} (1 - k^q \sin_q^q \theta)^{\frac{1}{q}} d\theta = \int_0^1 \left(\frac{1 - k^q t^q}{1 - t^q} \right)^{\frac{1}{q}} dt.$$

Takeuchi did a detailed, in-depth study of integrals of this kind. he showed Legendre's relation for $K_q(k)$ and $E_q(k)$ and observed relationship between the complete p -elliptic integrals and the Gaussian hyperbolic functions. As applications of complete p -elliptic, he also gave a computation formula of π_q with $q = 3$ and an elementary proof of Ramanujan's cubic transformation. The reader can see [22] and closely related references therein. In contrast to Lindqvist's trigonometric functions and Takeuchi's generalized elliptic integrals, similar elliptic integrals can be defined using the definition of p -trigonometric functions, but very unfortunately, it is difficult to study. Here are some questions for further study.

- Open Problem 5.1.** (i) Give the complete monotonicity and concavity and convexity of these functions $\sin_p x, \cos_p x$ and their inverse;
- (ii) Give the complete monotonicity of these functions $\sin_p x, \cos_p x$ and their inverse to parameter p ;
- (iii) Give some classical inequality of these functions $\sin_p x, \cos_p x$ and their inverse, such as Wilker type inequality, Cusa-Huygens type inequality, Kober inequality, Lazarević inequality et. al.;
- (iiii) Give series representation, integral representation of $\sin_p x, \cos_p x$ and their inverse to parameter p ;
- (iiiii) Give representation of hypergeometric function of $\sin_p x, \cos_p x$ and their inverse to parameter p .

By using the definition of p -trigonometric functions, we may define similar p -elliptic integrals $K_p(k)$ and $E_p(k)$.

- Open Problem 5.2.** (i) Give Legendre identity of p -elliptic integrals $K_p(k)$ and $E_p(k)$;
- (ii) Study multiple angle formula of p -elliptic integrals $K_p(k)$ and $E_p(k)$;
- (iii) Give the complete monotonicity and concavity and convexity of p -elliptic integrals $K_p(k)$ and $E_p(k)$ to parameter p ;
- (iiii) Give representation of hypergeometric function of p -elliptic integrals $K_p(k)$ and $E_p(k)$.

REFERENCES

- [1] Á. BARICZ, B. A. BHAYO, T. K. POGÁNY: *Functional inequalities for generalized inverse trigonometric and hyperbolic functions*. J. Math. Anal. Appl., **417** (2014), 244-259. <http://arxiv.org/abs/1401.4863> 7

- [2] Á. BARICZ, B. A. BHAYO, M. VUORINEN: *Turán type inequalities for generalized inverse trigonometric functions*. Filomat, **29**, No. 2 (2015), 303-313. <http://arxiv.org/abs/1209.1696>. 7
- [3] Á. BARICZ, B. A. BHAYO, R. KLÉN: *Convexity properties of generalized trigonometric and hyperbolic functions*. Aequat. Math., **89** (2015), 473-484. <http://arxiv.org/abs/1301.0699>. 7
- [4] Á. BARICZ, B. A. BHAYO, T. K. POGÁNY: *Convexity properties of generalized trigonometric and hyperbolic functions*, J. Math. Anal. Appl., **417**,(2014), 244-259 5
- [5] B. A. BHAYO, M. VUORINEN: *Inequalities for eigenfunctions of the p -Laplacian*, Issues of Analysis, **2**, No. 20 (2013), 13-35. <http://arxiv.org/abs/1101.3911> 7
- [6] B.A. BHAYO, M. VUORINEN: *Power mean inequalities generalized trigonometric functions*. Math. Vesnik, **67**, No. 1 (2015), 17-25. <http://arxiv.org/abs/1209.0983> 7
- [7] R. KLÉN, M. VUORINEN, X.-H. ZHANG: *Inequalities for the generalized trigonometric and hyperbolic functions*. J. Math. Anal. Appl., **409** (2014), 521-529. <http://arxiv.org/abs/1210.6749> 7
- [8] B. A. BHAYO AND L. YIN: *Logarithmic mean inequality for generalized trigonometric and hyperbolic functions*. Acta. Univ. Sapientiae Math., **6**, No. 2 (2014), 135-145. <http://arxiv.org/abs/1404.6732> 7
- [9] B. A. BHAYO AND L. YIN: *On the generalized convexity and concavity*. Problemy Analiza-Issues of Analysis, **22**, No. 1 (2015), 1-9. <http://arxiv.org/abs/1411.6586> 7
- [10] B. A. BHAYO AND L. YIN: *On the conjecture of generalized trigonometric and hyperbolic functions*. Math. Pannon., Vol. 24, No. 2 (2013), 1-8. <http://arxiv.org/abs/1402.7331> 7
- [11] D. B. KARP AND E. G. PRILEPKINA: *Parameter convexity and concavity of generalized trigonometric functions*. J. Math. Anal. Appl., **421**, No. 1 (2015), 370-382. <http://arxiv.org/abs/1402.3357> 7
- [12] J. C. KUANG: *Applied inequalities(Second edition)*. Shan Dong Science and Technology Press. Jinan, 2002. 4
- [13] P. LINDQVIST: *Some remarkable sine and cosine functions*. Ricerche di Math., **XLIV** (1995), 269-290. 7
- [14] T. J. MILDORF: *A sharp bound on the two variable powers mean*, Mathematical Reflections, 2006, 2, pp3-7. 2
- [15] D. S. MITRINOVIC, *Analytic inequalities*, Springer-verlag, New York, 1970. 6
- [16] W. D. Richter, *Generalized spherical and simplicial coordinates*, J. Math. Anal. Appl., **336**(2007), 1187-1202. 1
- [17] W. D. Richter, *Continuous $l_{n,p}$ -symmetric distributions*, Lithuanian Math. J., **49**(1)(2009), 93-108. 1
- [18] J. SÁNDOR: *On some new Wilker and Huygens type trigonometric-hyperbolic inequalities*, Proceedings of the Jangjeon Mathematical Society, **15**, No. 2 (2012). <http://arxiv.org/abs/1105.0859> 7
- [19] S. TAKEUCHI: *Generalized Jacobian elliptic functions and their application to bifurcation problems associated with p -Laplacian*. J. Math. Anal. Appl., **385** (2012), 24-35. 7
- [20] T. KAMIYA AND S. TAKEUCHI: *Complete (p, q) -elliptic integrals with application to a family of means*. J. Classical Anal., **10**, No. 1:15-25 <http://arxiv.org/abs/1507.01383> 7
- [21] S. TAKEUCHI: *The complete p -elliptic integrals and a computation formula of π_p for $p = 4$* . Ramanujan Journal , 2018 , 46 (2) :309-321 <http://arxiv.org/abs/1503.02394> 7
- [22] S. TAKEUCHI: *A new form of the generalized complete elliptic integrals*. Kodai J. Math., **39**, No. 1 (2016), 202-226. <http://arxiv.org/abs/1411.4778> 7, 8
- [23] S. TAKEUCHI: *Multiple-angle formulas of generalized trigonometric functions with two parameters*. <http://arxiv.org/abs/1603.06709> 7
- [24] S. TAKEUCHI: *Legendre-type relations for generalized complete elliptic integrals*. Journal of Classical Analysis, **9**, No. 1 (2016), 35-42. <http://arxiv.org/abs/1606.05115> 7

(Yin) COLLEGE OF SCIENCE, BINZHOU UNIVERSITY, BINZHOU CITY, SHANDONG PROVINCE,
256603, CHINA

E-mail address: yinli.79@163.com

(Lin) COLLEGE OF MATHEMATICS SCIENCE, QUFU NORMAL UNIVERSITY, QUFU CITY, SHANDONG
PROVINCE, 273165, CHINA

E-mail address: 58681180@qq.com