

Grüss-type fractional inequality via Caputo-Fabrizio integral operator

Asha B. Nale¹ Satish K. Panchal² Vaijanath L. Chinchane³,

^{1,2}Department of Mathematics,

Dr. Babasaheb Ambedkar Marathwada

University, Aurangabad-431 004, INDIA.

³Department of Mathematics,

Deogiri Institute of Engineering and Management

Studies Aurangabad-431005, INDIA

ashabnale@gmail.com/drskpanchal@gmail.com/chinchane85@gmail.com

Abstract

In this article, the main objective is to establish the Grüss-type fractional integral inequalities by employing the Caputo-Fabrizio fractional integral.

Mathematics Subject Classification: 26D10, 26D33.

Keywords: Grüss-type inequality, Caputo-Fabrizio fractional integral and integral inequality.

1 Introduction

Grüss inequality which establishes a connection between the integral of the product of two functions and the product of the integrals of the two functions. In 1935, G. Grüss proved the following well known classical integral inequality, see [24, 27].

Theorem 1.1 [27] *Let $f, g : [a, b] \rightarrow R$ be two integrable functions such that $\phi \leq f(x) \leq \Phi$ and $\gamma \leq g(x) \leq \Gamma$ for all $x \in [a, b]$; ϕ, Φ, γ and Γ are constant, then*

$$\left| \frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{b-a} \int_a^b f(x)dx \cdot \frac{1}{b-a} \int_a^b g(x)dx \right| \leq \frac{1}{4}(\Gamma - \gamma)(\Phi - \phi), \quad (1.1)$$

where the constant $\frac{1}{4}$ is sharp.

During the last few years, several numerous generalizations, variants and extensions of the Grüss inequality have appeared in the literature, see [18, 19,

20, 22, 24, 25, 26, 27, 29, 30, 31, 40] and the references cited therein. Chinchane and Pachpatte [10], investigated some new fractional integral inequalities of the Grüss-type by considering the Saigo fractional integral operator. In [1, 21, 34, 35, 36] authors obtained some the Grüss-type inequalities by using different types of fractional integral operators. Fractional calculus is generalization of traditional calculus into non-integer differential and integral order. Fractional calculus is very important due to it's various application in field of science and technology, see [2, 4, 32, 37]. In [5, 6], M. Caputo and M. Fabrizio introduced a new fractional derivative and application of new time and Spatial fractional derivative with exponential kernels. In literature very little work is reported on fractional integral inequalities using Caputo and Caputo-Fabrizio integral operator. X. Wang et al. [39] presented some properties of Caputo–Fabrizio fractional integral operator in the setting of-convex function. Recently, Gustava Nchama and et al. [28], proposed some fractional integral inequalities using the Caputo-Fabrizio fractional integral. Recently, many researchers have worked on fractional integral inequalities using the Riemann-Liouville, Hadamard and q-fractional integral, see [3, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 23, 38]. In [16], Dahmani and et al. gave the following fractional integral inequality using the Riemann-Liouville fractional integral. Motivated from [5, 6, 10, 16, 28, 39], our purpose in this paper is to propose some new results using the Caputo-Fabrizio integral operator. The paper has been organized as follows, in Section 2, we recall some auxiliary results related to the Caputo-Fabrizio integral operator. In Section 3, we investigate the Grüss-type fractional integral inequality using the Caputo-Fabrizio integral operator, in Section 4, we give the concluding remarks.

2 Preliminaries

In this section, we give some auxiliary results of fractional calculus that will be useful in this paper.

Definition 2.1 [6, 28] *Let $\alpha \in \mathbb{R}$ such that $0 < \alpha < 1$. The Caputo-Fabrizio fractional integral of order α of a function f is defined by*

$$\mathcal{I}_{0,x}^{\alpha} f(x) = \frac{1}{\alpha} \int_0^x e^{-(\frac{1-\alpha}{\alpha})(x-s)} f(s) ds. \quad (2.1)$$

Definition 2.2 [6, 28] *Let $\alpha, a \in \mathbb{R}$ such that $0 < x < 1$. The Caputo-Fabrizio fractional derivative of order α of a function f is defined by*

$$\mathcal{I}_{a,x}^{\alpha} f(x) = \frac{1}{1-\alpha} \int_a^x e^{\frac{-\alpha}{1-\alpha}(x-s)} f'(s) ds. \quad (2.2)$$

Definition 2.3 Let $\alpha > 0, \beta, \eta \in \mathbb{R}$, then the Saigo fractional integral $\mathcal{I}_{0,x}^{\alpha,\beta,\eta}[f(x)]$ of order α for a real valued continuous function $f(x)$ is defined by

$$\mathcal{I}_{0,x}^{\alpha,\beta,\eta} f(x) = \frac{x^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^x (x-\tau)^{\alpha-1} F_1(\alpha+\beta, -\eta; \alpha; 1-\frac{\tau}{x}) f(\tau) d\tau, \quad (2.3)$$

where the function $F_1(-)$ is the Gaussian hypergeometric function defined by

$$F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!},$$

and $(a)_n$ is the pochhammer symbol

$$(a)_n = a(a+1)\dots(a+n-1), (a)_0 = 1.$$

Definition 2.4 The Hadamard fractional integral is defined by

$${}^H\mathcal{I}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_1^x (\log \frac{x}{\tau})^{\alpha-1} f(\tau) \frac{d\tau}{\tau} \text{ for } \operatorname{Re}(\alpha) > 0, x > 1. \quad (2.4)$$

Definition 2.5 The Riemann-Liouville fractional integral is defined by

$$\mathcal{I}_{0,x}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-\tau)^{\alpha-1} f(\tau) d\tau. \quad (2.5)$$

3 Grüss-type fractional integral inequality

In this section, we investigate Grüss-type fractional integral inequalities involving the Caputo-Fabrizio fractional integer operator, for which assume that

(H_1) There exist two integrable function $\Phi_1(x), \Phi_2(x)$ on $[0, \infty[$, such that

$$\Phi_1(x) \leq u(x) \leq \Phi_2(x), \text{ for all } x \in [0, \infty[.$$

(H_2) There exist two integrable function $\Psi_1(x), \Psi_2(x)$ on $[0, \infty[$, such that

$$\Psi_1(x) \leq v(x) \leq \Psi_2(x), \text{ for all } x \in [0, \infty[.$$

Theorem 3.1 Suppose that u be an integrable function defined on $[0, \infty[$, consider the condition (H_1) hold. Then for all $x > 0, \alpha, \beta > 0$, we have

$$\begin{aligned} \mathcal{I}_{0,t}^\beta \Phi_1(x) \mathcal{I}_{0,t}^\alpha u(x) + \mathcal{I}_{0,t}^\alpha \Phi_2(x) \mathcal{I}_{0,t}^\beta u(x) &\geq \\ \mathcal{I}_{0,t}^\alpha \Phi_2(x) \mathcal{I}_{0,t}^\beta \Phi_1(x) + \mathcal{I}_{0,t}^\alpha u(x) \mathcal{I}_{0,t}^\beta u(x). & \end{aligned} \quad (3.1)$$

Proof:- From condition (H_1) , for all $\rho, \sigma \geq 0$, we obtain

$$(\Phi_2(\rho) - u(\rho))(u(\sigma) - \Phi_1(\sigma)) \geq 0. \quad (3.2)$$

that is

$$\Phi_2(\rho)u(\sigma) - \Phi_2(\rho)\Phi_1(\sigma) - u(\rho)u(\sigma) + u(\rho)\Phi_1(\sigma) \geq 0, \quad (3.3)$$

which implies that

$$\Phi_2(\rho)u(\sigma) + u(\rho)\Phi_1(\sigma) \geq \Phi_2(\rho)\Phi_1(\sigma) + u(\rho)u(\sigma). \quad (3.4)$$

Multiplying (3.4) by $\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}$, which is positive because $\rho \in (0, x)$, $x > 0$.

$$\begin{aligned} & u(\sigma)\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}\Phi_2(\rho) + \Phi_1(\sigma)\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}u(\rho) \\ & \geq \Phi_1(\sigma)\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}\Phi_2(\rho) + u(\sigma)\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}u(\rho). \end{aligned} \quad (3.5)$$

Now, integrating (3.5) with respect to ρ from 0 to x , we have

$$\begin{aligned} & \frac{u(\sigma)}{\alpha} \int_0^t e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}\Phi_2(\rho)d\rho + \frac{\Phi_1(\sigma)}{\alpha} \int_0^t e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}u(\rho)d\rho \\ & \geq \frac{\Phi_1(\sigma)}{\alpha} \int_0^t e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}\Phi_2(\rho)d\rho + \frac{u(\sigma)}{\alpha} \int_0^t e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}u(\rho)d\rho, \end{aligned} \quad (3.6)$$

therefore

$$\begin{aligned} & u(\sigma)\mathcal{I}_{0,t}^\alpha \Phi_2(x) + \Phi_1(\sigma)\mathcal{I}_{0,t}^\alpha u(x) \\ & \geq \Phi_1(\sigma)\mathcal{I}_{0,t}^\alpha \Phi_2(x) + u(\sigma)\mathcal{I}_{0,t}^\alpha u(x). \end{aligned} \quad (3.7)$$

Now, multiplying (3.7) by $\frac{1}{\beta}e^{-(\frac{1-\beta}{\beta})(x-\sigma)}$, which is positive because $\sigma \in (0, x)$, $x > 0$. Then integrating obtained result with respect to σ from 0 to x , we obtain

$$\begin{aligned} & \frac{\mathcal{I}_{0,x}^\alpha \Phi_2(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)}u(\sigma)d\sigma + \frac{\mathcal{I}_{0,x}^\alpha u(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)}\Phi_1(\sigma)d\sigma \\ & \geq \frac{\mathcal{I}_{0,t}^\alpha \Phi_2(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)}\Phi_1(\sigma)d\sigma + \frac{\mathcal{I}_{0,t}^\alpha u(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)}u(\sigma)d\sigma. \end{aligned} \quad (3.8)$$

This completes the proof.

Remark 3.1 If u be an integrable function defined on $[0, \infty[$, such that $\gamma \leq u(x) \leq \Gamma$, for all $x \in [0, \infty[$ and $\gamma, \Gamma \in \mathbb{R}$. Then for all $x > 0$ and $\alpha, \beta > 0$, we have

$$\begin{aligned} & \gamma \left(\frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u(x) + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\beta u(x) \\ & \geq \Gamma \gamma \left(\frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta u(x). \end{aligned} \quad (3.9)$$

Theorem 3.2 If u and v be two integrable functions defined on $[0, \infty[$, Suppose that (H_1) and (H_2) holds. Then for all $x > 0$, $\alpha, \beta > 0$, the following inequalities satisfied

$$\begin{aligned} (h1) \quad & \mathcal{I}_{0,x}^\beta \Psi_1(x) \mathcal{I}_{0,x}^\alpha u(x) + \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\beta v(x) \\ & \geq \mathcal{I}_{0,x}^\beta \Psi_1(x) \mathcal{I}_{0,x}^\alpha \Phi_2(x) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x). \\ (h2) \quad & \mathcal{I}_{0,x}^\beta \Phi_1(x) \mathcal{I}_{0,x}^\alpha v(x) + \mathcal{I}_{0,x}^\alpha \Psi_2(x) \mathcal{I}_{0,x}^\beta u(x) \\ & \geq \mathcal{I}_{0,x}^\beta \Phi_1(x) \mathcal{I}_{0,x}^\alpha \Psi_2(x) + \mathcal{I}_{0,x}^\beta u(x) \mathcal{I}_{0,x}^\alpha v(x). \\ (h3) \quad & \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\beta \Psi_2(x) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x) \\ & \geq \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\beta v(x) + \mathcal{I}_{0,x}^\beta \Psi_2(x) \mathcal{I}_{0,x}^\alpha u(x). \\ (h4) \quad & \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\beta \Psi_1(x) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x) \\ & \geq \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\beta v(x) + \mathcal{I}_{0,x}^\beta \Psi_1(x) \mathcal{I}_{0,x}^\alpha u(x). \end{aligned} \quad (3.10)$$

Proof To prove (h1), we use condition (H_1) and (H_2) , for all $x \in [0, \infty[$, we have

$$(\Phi_2(\rho) - u(\rho))(v(\sigma) - \Psi_1(\sigma)) \geq 0, \quad (3.11)$$

which implies that

$$\Phi_2(\rho)v(\sigma) + u(\rho)\Psi_1(\sigma) \geq \Phi_2(\rho)\Psi_1(\sigma) + u(\rho)u(\sigma). \quad (3.12)$$

Multiplying (3.12) by $\frac{1}{\alpha}e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)}$, which is positive because $\rho \in (0, x)$, $x > 0$

$$\begin{aligned} & v(\sigma) \frac{1}{\alpha} e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} \Phi_2(\rho) + \Psi_1(\sigma) \frac{1}{\alpha} e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} u(\rho) \\ & \geq \Psi_1(\sigma) \frac{1}{\alpha} e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} \Phi_2(\rho) + v(\sigma) \frac{1}{\alpha} e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} u(\rho). \end{aligned} \quad (3.13)$$

Integrating (3.13) with respect to ρ from 0 to x , we get

$$\begin{aligned} & \frac{v(\sigma)}{\alpha} \int_0^t e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} \Phi_2(\rho) d\rho + \frac{\Psi_1(\sigma)}{\alpha} \int_0^t e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} u(\rho) d\rho \\ & \geq \frac{\Psi_1(\sigma)}{\alpha} \int_0^t e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} \Phi_2(\rho) d\rho + \frac{v(\sigma)}{\alpha} \int_0^t e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} u(\rho) d\rho, \end{aligned} \quad (3.14)$$

therefore

$$v(\sigma)\mathcal{I}_{0,x}^\alpha\Phi_2(x) + \Psi_1(\sigma)\mathcal{I}_{0,x}^\alpha u(x) \geq \Psi_1(\sigma)\mathcal{I}_{0,x}^\alpha\Phi_2(x) + v(\sigma)\mathcal{I}_{0,x}^\alpha u(x). \quad (3.15)$$

Multiplying both sides of (3.15) by $\frac{1}{\beta}e^{-(\frac{1-\beta}{\beta})(x-\sigma)}$, which is positive because $\sigma \in (0, x)$, $x > 0$. Then integrating resulting identity with respect to σ over 0 to x , we obtain

$$\begin{aligned} & \frac{\mathcal{I}_{0,x}^\alpha\Phi_2(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)} v(\sigma) d\sigma + \frac{\mathcal{I}_{0,x}^\alpha u(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)} \Psi_1(\sigma) d\sigma \\ & \geq \frac{\mathcal{I}_{0,x}^\alpha\Phi_2(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)} v \Psi_1(\sigma) d\sigma + \frac{\mathcal{I}_{0,x}^\alpha u(x)}{\beta} \int_0^x e^{-(\frac{1-\beta}{\beta})(x-\sigma)} v(\sigma) d\sigma. \end{aligned} \quad (3.16)$$

This gives desired inequality (h1).

To prove (h2)-(h4), we use the following inequalities respectively

$$\begin{aligned} & (\Psi_2(\rho) - v(\rho)) (u(\sigma) - \Phi_1(\sigma)) \geq 0. \\ & (\Phi_2(\rho) - u(\rho)) (v(\sigma) - \Psi_2(\sigma)) \leq 0. \\ & (\Phi_2(\rho) - u(\rho)) (v(\sigma) - \Psi_1(\sigma)) \leq 0. \end{aligned}$$

Remark 3.2 *If u and v be two integrable function defined on $[0, \infty[$, Assume that (H_3) There exist real constant $\Gamma, \gamma, \Gamma', \gamma'$ such that*

$$\gamma \leq u(x) \leq \Gamma \text{ and } \gamma' \leq v(x) \leq \Gamma' \quad \forall x \in [0, \infty[. \quad (3.17)$$

Then for all $x > 0$, $\alpha, \beta > 0$, the following inequalities satisfied

$$\begin{aligned}
(h_1) & \left(\gamma' \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u(x) + \left(\Gamma \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\beta v(x) \\
& \geq \left(\gamma' \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \left(\Gamma \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x). \\
(h_2) & \left(\gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha v(x) + \left(\Gamma' \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\beta u(x) \\
& \geq \left(\gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \left(\Gamma' \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\beta u(x) \mathcal{I}_{0,x}^\alpha v(x). \\
(h_3) & \left(\Gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \left(\Gamma' \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x) \\
& \geq \left(\Gamma \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\beta v(x) + \left(\gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u(x). \\
(h_4) & \left(\gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \left(\gamma' \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\beta v(x) \\
& \geq \left(\gamma \frac{1}{1-\beta} \left[1 - e^{-\left(\frac{1-\beta}{\beta}\right)x} \right] \right) \mathcal{I}_{0,x}^\beta v(x) + \left(\gamma' \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u(x).
\end{aligned} \tag{3.18}$$

Lemma 3.3 If u be an integrable function on $[0, \infty)$, and $\Phi_1(x)$, $\Phi_2(x)$ be two integrable functions on $[0, \infty)$. Assume that the condition H_1 holds. Then for all $x > 0$, $\alpha > 0$, we have

$$\begin{aligned}
& \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u^2(x) - \left(\mathcal{I}_{0,x}^\alpha u(x) \right)^2 \\
& = \left(\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x) \right) \left(\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x) \right) \\
& - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha [(\Phi_2(x) - u(x))(u(x) - \Phi_1(x))] \\
& + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_1 u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_2 u(x) - \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha \Phi_2(x) - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x).
\end{aligned} \tag{3.19}$$

Proof:- Since u be an integrable function on $[0, \infty)$. For all $\rho, \sigma > 0$, we have

$$\begin{aligned}
& (\Phi_2(\sigma) - u(\sigma)) (u(\rho) - \Phi_1(\rho)) + (\Phi_2(\rho) - u(\rho)) (u(\sigma) - \Phi_1(\sigma)) \\
& - (\Phi_2(\rho) - u(\rho)) (u(\rho) - \Phi_1(\rho)) - (\Phi_2(\sigma) - u(\sigma)) (u(\sigma) - \Phi_1(\sigma)) \\
& = u^2(\rho) + u^2(\sigma) - 2u(\rho)u(\sigma) + \Phi_2(\sigma)u(\rho) + \Phi_1(\rho)u(\sigma) - \Phi_1(\rho)\Phi_2(\sigma) \\
& + \Phi_1(\rho)u(\sigma) + \Phi_1(\sigma)u(\rho) - \Phi_1(\sigma)\Phi_2(\rho) - \Phi_2(\rho)u(\rho) + \Phi_1(\rho)\Phi_2(\rho) \\
& - \Phi_1(\rho)u(\rho) - \Phi_2(\sigma)u(\sigma) + \Phi_1(\sigma)\Phi_2(\sigma) - \Phi_1(\sigma)u(\sigma).
\end{aligned} \tag{3.20}$$

Multiplying both sides of (3.20) by $\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\rho)}$, which is positive because $\rho \in (0, x)$, $x > 0$, integrating obtained result with respect to ρ from 0 to x , we have

$$\begin{aligned}
& (\Phi_2(\sigma) - u(\sigma)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x)) + (\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (u(\sigma) - \Phi_1(\sigma)) \\
& - \mathcal{I}_{0,x}^\alpha [(\Phi_2(x) - u(x)) (u(x) - \Phi_1(x))] - \frac{(\ln x)^\alpha}{\Gamma(\alpha + 1)} (\Phi_2(\sigma) - u(\sigma)) (u(\sigma) - \Phi_1(\sigma)) \\
& = \mathcal{I}_{0,x}^\alpha u^2(x) + u^2(\sigma) \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) - 2u(\sigma)\mathcal{I}_{0,x}^\alpha u(x) + \Phi_2(\sigma)\mathcal{I}_{0,x}^\alpha u(x) \\
& + u(\sigma)\mathcal{I}_{0,x}^\alpha \Phi_1(x) - \Phi_2(\sigma)\mathcal{I}_{0,x}^\alpha \Phi_1(x) + u(\sigma)\mathcal{I}_{0,x}^\alpha \Phi_2(x) \\
& + \Phi_1(\sigma)\mathcal{I}_{0,x}^\alpha u(x) - \Phi_1(\sigma)\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha \Phi_2 u(x) \\
& + \mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x) - \mathcal{I}_{0,x}^\alpha \Phi_1 u(x) - \Phi_2(\sigma)u(\sigma) \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \\
& + \Phi_1(\sigma)\Phi_2(\sigma) \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) - \Phi_1(\sigma)u(\sigma) \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right).
\end{aligned} \tag{3.21}$$

Again, multiplying (3.21) by $\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\sigma)}$, which is positive because $\sigma \in (0, x)$, $x > 0$, integrating obtained result with respect to ρ from 0 to x , we

have

$$\begin{aligned}
& (\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x)) \\
& + (\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x)) \\
& - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha [(\Phi_2(x) - u(x)) (u(x) - \Phi_1(x))] \\
& - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha [(\Phi_2(x) - u(x)) (u(x) - \Phi_1(x))] \\
& = \mathcal{I}_{0,x}^\alpha u^2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha u^2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) - 2\mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha u(x) + \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha \Phi_1(x) - \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha \Phi_1(x) \\
& + \mathcal{I}_{0,x}^\alpha u(x) {}_H D_{1,x}^{-\alpha} \Phi_2(x) + \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha \Phi_2(x) \\
& - \mathcal{I}_{0,x}^\alpha \Phi_2 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \\
& - \mathcal{I}_{0,x}^\alpha \Phi_1 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \\
& - \mathcal{I}_{0,x}^\alpha \Phi_2 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + \mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \\
& - \mathcal{I}_{0,x}^\alpha \Phi_1 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right), \tag{3.22}
\end{aligned}$$

which implies that

$$\begin{aligned}
& 2 (\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x)) \\
& - 2 \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha [(\Phi_2(x) - u(x)) (u(x) - \Phi_1(x))] \\
& = 2\mathcal{I}_{0,x}^\alpha u^2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) - 2(\mathcal{I}_{0,x}^\alpha u(x))^2 + 2\mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + 2\mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha \Phi_1(x) - 2\mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha \Phi_1(x) \\
& - 2\mathcal{I}_{0,x}^\alpha \Phi_2 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) + 2\mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \\
& - 2\mathcal{I}_{0,x}^\alpha \Phi_1 u(x) \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right). \tag{3.23}
\end{aligned}$$

This completes the proof.

If $\Phi_1(x) = \gamma$ and $\Phi_2(x) = \Gamma$; $\gamma, \Gamma \in \mathbb{R}$ for all $x \in [0, \infty)$, then inequality (3.19) reduces to following lemma.

Lemma 3.4 *If $\gamma, \Gamma \in \mathbb{R}$, and $u(x)$ be an integrable function on $[0, \infty)$ and satisfying the condition $\gamma \leq u(x) \leq \Gamma$. Then for all $x > 0$, $\alpha > 0$, we have*

$$\begin{aligned} & \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u^2(x) - (\mathcal{I}_{0,x}^\alpha u(x))^2 = \left(\Gamma \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) - \mathcal{I}_{0,x}^\alpha u(x) \right) \\ & \left(\mathcal{I}_{0,x}^\alpha u(x) - \left(\gamma \frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \right) - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha ((\Gamma - u(x))(u(x) - \gamma)). \end{aligned} \quad (3.24)$$

Theorem 3.5 *Let u and v be two integrable functions on $[0, \infty)$, and $\Phi_1(x)$, $\Phi_2(x)$, $\Psi_1(x)$ and $\Psi_2(x)$ are four integrable functions on $[0, \infty)$ satisfying the conditions H_1 and H_2 on $[0, \infty)$. Then for all $x > 0$, $\alpha > 0$, we have*

$$\begin{aligned} & \left| \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha uv(x) - \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha v(x) \right| \\ & \leq \sqrt{R(u, \Phi_1(x), \Phi_2(x)) R(v, \Psi_1(x), \Psi_2(x))}. \end{aligned} \quad (3.25)$$

where $R(a, b, c)$ is defined by

$$\begin{aligned} R(a, b, c) &= (\mathcal{I}_{0,x}^\alpha c(x) - \mathcal{I}_{0,x}^\alpha a(x)) (\mathcal{I}_{0,x}^\alpha a(x) - \mathcal{I}_{0,x}^\alpha b(x)) \\ &+ \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha ba(x) - \mathcal{I}_{0,x}^\alpha b(x) \mathcal{I}_{0,x}^\alpha a(x) \\ &+ \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha ca(x) - \mathcal{I}_{0,x}^\alpha c(x) \mathcal{I}_{0,x}^\alpha a(x) \\ &+ \mathcal{I}_{0,x}^\alpha b(x) \mathcal{I}_{0,x}^\alpha c(x) + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha bc(x). \end{aligned} \quad (3.26)$$

Proof Let u and v be two functions defined on $[0, \infty)$ satisfying the condition H_1 and H_2 . Define

$$H(\rho, \sigma) := (u(\rho) - u(\sigma)) (v(\rho) - v(\sigma)); \rho, \sigma \in (0, x), x > 0, \quad (3.27)$$

it follows that

$$H(\rho, \sigma) := u(\rho)v(\rho) - u(\rho)v(\sigma) - u(\sigma)v(\rho) + u(\sigma)v(\sigma). \quad (3.28)$$

Now, multiplying (3.28) by $\frac{1}{\alpha} e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)}$, which is positive because $\rho \in (0, x)$, $x > 0$, integrating obtained result with respect to ρ from 0 to x , we have

$$\begin{aligned} & \frac{1}{\alpha} \int_0^x e^{-\left(\frac{1-\alpha}{\alpha}\right)(x-\rho)} H(\rho, \sigma) d\rho \\ &= \mathcal{I}_{0,x}^\alpha uv(x) - u(\tau) \mathcal{I}_{0,x}^\alpha u(x) - u(\sigma) \mathcal{I}_{0,x}^\alpha v(x) + u(\sigma)v(\sigma) \mathcal{I}_{0,x}^\alpha (1). \end{aligned} \quad (3.29)$$

Multiplying (3.29) by $\frac{1}{\alpha}e^{-(\frac{1-\alpha}{\alpha})(x-\sigma)}$, which is positive because $\sigma \in (0, x)$, $x > 0$, integrating obtained result with respect to σ from 0 to x , we have

$$\begin{aligned} & \frac{1}{\alpha^2} \int_0^t \int_0^t e^{-(\frac{1-\alpha}{\alpha})(x-\rho)} e^{-(\frac{1-\alpha}{\alpha})(x-\sigma)} H(\rho, \sigma) d\rho d\sigma \\ & = 2 \left(\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha uv(x) - \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha v(x) \right). \end{aligned} \quad (3.30)$$

Applying the Cauchy-Schwarz inequality to (3.30), we have

$$\begin{aligned} & \left(\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha uv(x) - \mathcal{I}_{0,x}^\alpha u(x) \mathcal{I}_{0,x}^\alpha v(x) \right)^2 \leq \\ & \left(\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha u^2(x) - (\mathcal{I}_{0,x}^\alpha u(x))^2 \right) \times \\ & \left(\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha v^2(x) - \mathcal{I}_{0,x}^\alpha v(x)^2 \right), \end{aligned} \quad (3.31)$$

since $(\Phi_2(x) - u(t))(u(t) - \Phi_1(x)) \geq 0$ and $(\Psi_2(x) - v(t))(v(t) - \Psi_1(x)) \geq 0$, we have

$$\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha (\Phi_2(x) - u(t))(u(t) - \Phi_1(x)) \geq 0, \quad (3.32)$$

and

$$\left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha (\Psi_2(x) - v(t))(v(t) - \Psi_1(x)) \geq 0. \quad (3.33)$$

Thus, we have

$$\begin{aligned} & \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha u^2(x) - (\mathcal{I}_{0,x}^\alpha u(x))^2 \\ & \leq (\mathcal{I}_{0,x}^\alpha \Phi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x)) \\ & + \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_1 u(x) - \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha u(x) \\ & + \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_2 u(x) - \mathcal{I}_{0,x}^\alpha \Phi_2(x) \mathcal{I}_{0,x}^\alpha u(x) \\ & + \mathcal{I}_{0,x}^\alpha \Phi_1(x) \mathcal{I}_{0,x}^\alpha \Phi_2(x) - \left(\frac{1}{1-\alpha} \left[1 - e^{-(\frac{1-\alpha}{\alpha})x} \right] \right) \mathcal{I}_{0,x}^\alpha \Phi_1 \Phi_2(x) \\ & = R(u, \Phi_1, \Phi_2), \end{aligned} \quad (3.34)$$

and

$$\begin{aligned}
& \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha u^2(x) - (\mathcal{I}_{0,x}^\alpha u(x))^2 \\
& \leq (\mathcal{I}_{0,x}^\alpha \Psi_2(x) - \mathcal{I}_{0,x}^\alpha u(x)) (\mathcal{I}_{0,x}^\alpha u(x) - \mathcal{I}_{0,x}^\alpha \Psi_1(x)) \\
& + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Psi_1 u(x) - \mathcal{I}_{0,x}^\alpha \Psi_1(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Psi_2 u(x) - \mathcal{I}_{0,x}^\alpha \Psi_2(x) \mathcal{I}_{0,x}^\alpha u(x) \\
& + \mathcal{I}_{0,x}^\alpha \Psi_1(x) \mathcal{I}_{0,x}^\alpha \Psi_2(x) - \left(\frac{1}{1-\alpha} \left[1 - e^{-\left(\frac{1-\alpha}{\alpha}\right)x} \right] \right) \mathcal{I}_{0,x}^\alpha \Psi_1 \Psi_2(x) \\
& = R(u, \Psi_1, \Psi_2).
\end{aligned} \tag{3.35}$$

Combining (3.31), (3.34) and (3.35), we get (3.25).

4 Concluding Remarks

G. Nchama et al. [28], investigated some integral inequalities by considering Caputo-Fabrizio fractional integral operator. In [6] M. Caputo and Fabrizio introduced a new fractional differential and integral operator. Motivated by the above work, here we studied Grüss-type inequalities and other fractional inequalities by considering Caputo-Fabrizio fractional integral operator. By the help of this study we establish more general inequalities than in the classical cases. The inequalities investigated in this paper give some contribution to the fields of fractional calculus and Caputo-Fabrizio fractional integral operator. These inequalities are expected to lead to some application for finding bounds and uniqueness of solutions in fractional differential equations.

References

- [1] T. A. Aljaaidi and D. B. Pachpatte, *Some Grüss-type inequalities using generalized Katugampola fractional integral*, AIMS Mathematics, 2(5), (2020), 1011-1024. doi: 10.3934/math.2020070.
- [2] D. Baleanu, J. A. T. Machado and C. J. Luo, *Fractional Dynamic and Control*, Springer, 2012, pp 159-171.
- [3] S. Belarbi and Z. Dahmani, *On some new fractional integral inequality*, J. Inequal. Pure and Appl. Math., 10(3)(2009), Art.86, 5 pp.

- [4] P. L. Butzer, A. A. Kilbas and J. J. Trujillo, *Fractional Calculus in the Mellin setting and Hadardmard-type fractional integrals*, J. Math. Anal. Appl., (2002), 1-27.
- [5] M. Caputo and M. Fabrizio, *A new Defination of Fractional Derivative without singular kernel*, Progr., Fract., Differ., Appl., 1(2),(2015), 73-85.
- [6] M. Caputo and M. Fabrizio, *Applications of new time and Spatial fractional derivative with exponential kernels*, Progr., Fract., Differ., Appl., 2(2),(2016), 7-8.
- [7] V. L. Chinchane and D. B. Pachpatte, *A note on some integral inequalities via Hadamard integral*, J. Fractional Calculus Appl., 4(11)(2013), 1-5.
- [8] V. L. Chinchane and D. B. Pachpatte, *On some integral inequalities using Hadamard fractional integral*, Malaya J. Math., 1(1)(2012), 62-66.
- [9] V. L. Chinchane and D. B. Pachpatte, *Some new integral inequalities using Hadamard fractional integral operator*, Adv. Inequal. Appl., 2014, 2014:12.
- [10] V. L. Chinchane and D. B. Pachpatte, *On some Grüss-type fractional inequalities using Saigo fractional integral operator*, Journal of Mathematics, vol. 2014, Article ID 527910, 9 pages, 2014. <https://doi.org/10.1155/2014/527910>.
- [11] Z. Dahmani, *On Minkowski and Hermit-Hadamard integral inequalities via fractional integration*, Ann. Funct. Anal., 1(1)(2010), 51-58.
- [12] Z. Dahmani, *New inequalities in fractional integrals*, Int. J. Nonlinear Sci., 9, 4(2010), 493-497.
- [13] Z. Dahmani, *The Riemann-Liouville operator to generate some new inequalities*, Int. J. Nonlinear Sci., 12 (4)(2011), 452-455.
- [14] Z. Dahmani, *Some results associate with fractional integrals involving the extended Chebyshev*, Acta Univ. Apulensis Math. Inform., 27(2011), 217-224.
- [15] Z. Dahmani and A. Benzidane, *New inequalities using Q-fractional theory*, Bull. Math. Anal. Appl., 4(1)(2012), 190-196.

- [16] Z. Dahmani, L. Tabharit and S. Taf *New generalisation of Grüss inequality using Riemann-Liouville fractional integrals*, Bull. Math. Anal. Appl., 2(3)(2012), 92-99.
- [17] Z. Denton and A. S. Vatsala, *Fractional integral inequalities and application*, Comput. Math. Appl., 59(2010), 1087-1094.
- [18] S. S. Dragomir, *Some integral inequalities of Grüss type*, Indian J. Pur. Appl. Math., 31 (4) (2002), 397-415.
- [19] S. S. Dragomir, *A companion of the Grüss inequality and applications*, Appl. Math. Lett., 17 (2004), no. 4, 429-435.
- [20] S. S. Dragomir and S. Wang, *An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for home numerical quadrature rules*, Comput. Math. Appl., 33 (1997), no. 11, 15-20.
- [21] G. Farid, A. U. Rehman, V. N. Mishra, S. Mehmood, *Fractional integral inequalities of Grüss type via generalized Mittag-Leffler function*, Int. J. Anal. Appl., 17 (4) (2019), 548-558.
- [22] N. Elezovic, L. Marangunic and J. Pecaric, *Some improvements of Grüss type inequality*, J. Math. Inequal., 1 (2007), no. 3, 425-436.
- [23] A. A. George, *Fractional Differentiation Inequalities*, Springer Publishing Company, New York, 2009.
- [24] G. Grüss, *Über das maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx$* , Math. Z., 39 (1935), 215-226.
- [25] A. M. Mercer, P. Mercer, *New proofs of the Grüss inequality*, Aust. J. Math. Anal. Appl., 1(2) (2004), Art. 12.
- [26] A. M. Mercer, *An improvement of the Grüss inequality*, JIPAM. J. Inequal. Pure Appl. Math., 6 (2005), no. 4, Article 93.
- [27] Yu Miao, Fei Han and Jianyong Mu, *A New Ostrowski-Grüss type inequality*, Kragujevac Journal of Mathematics, 37(2) (2013), Pages 307-317.
- [28] G. A. M. Nchama, A. L. Mecias and M. R. Richard, *The Caputo-Fabrizio fractional integral to generate some new inequalities*, Inf. Sci. Lett. 2(8), (2019), Pages 73-80.

- [29] B. G. Pachpatte, *On multidimensional Grüss type inetegral inequalities*, J. Inequal. Pure and Appl. Math., Vol 03, Iss. 2, Art. 27, (2002), 1-7.
- [30] B. G. Pachpatte, *A note on Chebyshev-Grüss inequalities for differential equations*, Tamsui Oxf. J. Math. Sci., 22(1), (2006), 29-36.
- [31] G. H. Peng and Y. Miao, *A note on Grüss type inequality*, Appl. Math. Sci., (Ruse) 3 (2009), no. 8, 399-402.
- [32] I. Podlubny, *Fractional Differential Equations: An Introduction to Fractional Derivatives to method of their solution and some of their application*, Academic Press, San Diego CA, 1999.
- [33] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Application of Fractional Differential Equations*, Elsevier, Amersterdam, 2006.
- [34] G. Rahman, K. S. Nisar, S. Rashid and T. Abdeljawad, *Certain Grüss-type inequalities via tempered fractional integrals concerning another function*, J. Inequal. Appl., 2020, 147 (2020). <https://doi.org/10.1186/s13660-020-02420-x>.
- [35] S. Rashid, F. Jarad and M. A. Noor, *Grüss-type integrals inequalities via generalized proportional fractional operators*, RACSAM 114, 93 (2020). <https://doi.org/10.1007/s13398-020-00823-5>.
- [36] E. Set, M. Tomar and M. Z. Sarikaya, *On generalized Grüss type inequalities for k -fractional integrals*, Applied Mathematics and Computation 269, (2015), 29-34. DOI: 10.1016/j.amc.2015.07.026.
- [37] S. G. Somko, A. A. Kilbas and O. I. Marichev, *Fractional Integral and Derivative Theory and Application*, Gordon and Breach, Yverdon, Switzerland, 1993.
- [38] W. Yang, *Some new fractional quantum integral inequalities*, Appl. Math. Latt., 25, 6(2012), 963-969.
- [39] X. Wang, M. S. Saleem, K. N. Aslam, X. Wu and T. Zhou *On Caputo–Fabrizio fractional integral inequalities of Hermite–Hadamard type for modified -convex Functions*, Journal of Mathematics, vol. 2020, Article ID 8829140, 17 pages, 2020. <https://doi.org/10.1155/2020/8829140>.
- [40] C. Zhu, W. Yang and Q. Zhao *Some new fractional q - integral Grüss-type inequalities and other inequalities*, J. Inequal. Appl., 2012:299.