

## A THEOREM OF ROLEWICZ'S TYPE FOR MEASURABLE EVOLUTION FAMILIES IN BANACH SPACES

CONSTANTIN BUŞE AND SEVER S. DRAGOMIR

ABSTRACT. Let  $\mathbb{R}_+$  be the set of all non-negative real numbers,  $X$  be a Banach space and  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\}$  be a strongly measurable, exponentially bounded evolution family of bounded linear operators acting on  $X$ , satisfying a certain measurability condition as in Theorem 1 below. Let  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a non-decreasing function such that  $\varphi(t) > 0$  for all  $t > 0$ . We prove that if there exists  $M_\varphi > 0$  such that

$$\sup_{s \geq 0} \int_s^\infty \varphi(\|U(t, s)x\|) dt = M_\varphi < \infty, \text{ for all } x \in X, \|x\| \leq 1,$$

then  $\mathcal{U}$  is uniformly exponentially stable. For  $\varphi$  continuous and  $\mathcal{U}$  strongly continuous and exponentially bounded, this result is due to S. Rolewicz. The proofs uses the relatively recent techniques involving evolution semigroups theory.

Let  $X$  be a real or complex Banach space and  $\mathcal{L}(X)$  the Banach algebra of all linear and bounded operators on  $X$ . Let  $\mathbf{T} = \{T(t) : t \geq 0\} \subset \mathcal{L}(X)$  be a strongly continuous semigroup on  $X$  and  $\omega_0(\mathbf{T}) = \lim_{t \rightarrow \infty} \frac{\ln(\|T(t)\|)}{t}$  be its growth bound. The Datko-Pazy theorem ([2], [4]) states that  $\omega_0(\mathbf{T}) < 0$  if and only if for all  $x \in X$  the maps  $t \mapsto \|T(t)x\|$  belongs to  $L^p(\mathbb{R}_+)$  for some  $1 \leq p < \infty$ . Let  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a non-decreasing function such that  $\varphi(t) > 0$  for all  $t > 0$ . If for each  $x \in X, \|x\| \leq 1$  the maps  $t \mapsto \varphi(\|T(t)x\|)$  belongs to  $L^1(\mathbb{R}_+)$  then  $\mathbf{T}$  is exponentially stable, i.e.  $\omega_0(\mathbf{T})$  is negative. This later result is due to J. van Neerven [9, Theorem 3.2.2.]. The Datko-Pazy Theorem follows from this by taking  $\varphi(t) = t^p$  ( $t \geq 0$ ). Moreover, it is easily to see that the above Neerven's result remain true if we replace the strongly continuity assumption about  $\mathbf{T}$  with strongly measurability and exponentially boundedness assumptions about  $\mathbf{T}$ .

A family  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\} \subset \mathcal{L}(X)$  is called an *evolution family* of bounded linear operators on  $X$  if  $U(t, t) = I$  (the identity operator on  $X$ ) and  $U(t, \tau)U(\tau, s) = U(t, s)$  for all  $t \geq \tau \geq s \geq 0$ . Such a family is said to be *strongly continuous* if for every  $x \in X$ , the maps

$$(1) \quad (t, s) \mapsto U(t, s)x : \{(t, s) : t \geq s \geq 0\} \rightarrow X.$$

are continuous, and *exponentially bounded* if there are  $\omega > 0$  and  $K_\omega > 0$  such that

$$(2) \quad \|U(t, s)\| \leq K_\omega e^{\omega(t-s)} \text{ for all } t \geq s \geq 0.$$

If  $\mathbf{T} = \{T(t) : t \geq 0\} \subset \mathcal{L}(X)$  is a strongly continuous semigroup on  $X$ , then the family  $\{U(t, s) : t \geq s \geq 0\}$  given by  $U(t, s) = T(t-s)$  is a strongly continuous

---

*Date:* May 14, 2001.

*1991 Mathematics Subject Classification.* 47A30, 93D05, 35B35, 35B40, 46A30.

*Key words and phrases.* Evolution family of bounded linear operators, evolution operator semigroup, Rolewicz's theorem, exponential stability.

and exponentially bounded evolution family on  $X$ . Conversely, if  $\mathcal{U}$  is a strongly continuous evolution family on  $X$  and  $U(t, s) = U(t - s, 0)$  for all  $t \geq s \geq 0$  then the family  $\mathbf{T} = \{T(t) : t \geq 0\}$  is a strongly continuous semigroup on  $X$ . For more details about the strongly continuous semigroups and other references we refer to [4], [3]. The Datko-Pazy theorem can be also obtained from the following result given by S. Rolewicz ([5], [6]).

Let  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a continuous and nondecreasing function such that  $\varphi(0) = 0$  and  $\varphi(t) > 0$  for all  $t > 0$ . If  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\} \subset \mathcal{L}(X)$  is a strongly continuous and exponentially bounded evolution family on the Banach space  $X$  such that

$$(3) \quad \sup_{s \geq 0} \int_s^\infty \varphi(\|U(t, s)x\|) dt = M_\varphi < \infty, \quad \text{for all } x \in X, \|x\| \leq 1,$$

then  $\mathcal{U}$  is uniformly exponentially stable, that is (2) holds with some  $\omega < 0$ .

A shorter proof of the Rolewicz theorem was given by Q. Zheng [7] who removed the continuity assumption about  $\varphi$ . Other proofs (the semigroup case) of Rolewicz's theorem were offered by W. Littman [8] and J. van Neervan [9, pp. 81-82]. Some related results have been obtained by K.M. Przyłuski [11], G. Weiss [16] and J. Zabczyk [12].

A family  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\} \subset \mathcal{L}(X)$  is called *strongly measurable* if for all  $x \in X$  the maps given in (1) are measurable.

In this note we prove the following:

**Theorem 1.** Let  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a nondecreasing function such that  $\varphi(t) > 0$  for all  $t > 0$  and  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\} \subset \mathcal{L}(X)$  be a strongly measurable and exponentially bounded evolution family of operators on the Banach space  $X$ . If  $\mathcal{U}$  satisfies the conditions:

- (i) there exists  $M_\varphi > 0$  such that

$$(4) \quad \int_\xi^\infty \varphi(\|U(t, \xi)x\|) dt \leq M_\varphi < \infty, \quad \forall x \in X, \|x\| \leq 1, \forall \xi \geq 0,$$

- (ii) for all  $f \in L^1(\mathbb{R}_+, X)$  the maps

$$(5) \quad t \mapsto U(\cdot, \cdot - t)f(\cdot) : \mathbb{R}_+ \rightarrow L^1(\mathbb{R}_+, X)$$

are measurable, then  $\mathcal{U}$  is uniformly exponentially stable.

Firstly we prove the following Lemma which is essentially contained in [1, Theorem 2.1].

**Lemma 1.** Let  $\mathcal{U}$  be a strongly continuous and exponentially bounded evolution family of operators on  $X$  such that

$$(6) \quad \sup_{s \geq 0} \int_s^\infty \varphi(\|U(t, s)x\|) dt = M_\varphi(x) < \infty, \quad \text{for all } x \in X$$

Then  $\mathcal{U}$  is uniformly bounded, that is,

$$\sup_{t \geq \xi \geq 0} \|U(t, \xi)\| = C < \infty.$$

*Proof of Lemma 1.* Let  $x \in X$  and  $N(x)$  be a positive integer such that  $M_\varphi(x) < N(x)$  and let  $s \geq 0$ ,  $t \geq s + N$ . For each  $\tau \in [t - N, t]$ , we have

$$(7) \quad e^{-\omega N} 1_{[t-N, t]}(u) \|U(t, s)x\| \leq e^{-\omega(t-\tau)} 1_{[t-N, t]}(u) \|U(t, \tau)U(\tau, s)x\| \leq K_\omega \|U(u, s)x\|,$$

for all  $u \geq s$ . Here  $K_\omega$  and  $\omega$  are as in (2).

From (6) follows that  $\varphi(0) = 0$ . Then from (7) we obtain

$$(8) \quad N(x) \varphi\left(\frac{\|U(t, s)x\|}{K_\omega e^{\omega N}}\right) = \int_s^\infty \varphi\left(\frac{1_{[t-N, t]}(u) \|U(t, s)x\|}{K_\omega e^{\omega N}}\right) du \leq \int_s^\infty \varphi(\|U(u, s)x\|) du = M_\varphi(x).$$

We may assume that  $\varphi(1) = 1$  (if not, we replace  $\varphi$  by some multiple of itself). Moreover, we may assume that  $\varphi$  is a strictly increasing map. Indeed if  $\varphi(1) = 1$  and  $a := \int_0^1 \varphi(t) dt$ , then the function given by

$$\bar{\varphi}(t) = \begin{cases} \int_0^t \varphi(u) du, & \text{if } 0 \leq t \leq 1 \\ \frac{at}{at + 1 - a}, & \text{if } t > 1 \end{cases}$$

is strictly increasing and  $\bar{\varphi} \leq \varphi$ . Now  $\varphi$  can be replaced by some multiple of  $\bar{\varphi}$ . From (8) it follows that

$$\|U(t, s)\| \leq K_\omega e^{\omega N(x)}, \quad \text{for all } x \in X.$$

Now, it is easy to see that

$$(9) \quad \sup_{t \geq \xi \geq 0} \|U(t, \xi)x\| \leq 2K_\omega e^{\omega N(x)} := C(x) < \infty, \quad \text{for all } x \in X.$$

The assertion of Lemma 1 follows from (9) and the Uniform Boundedness Theorem. ■

It is clear that (3) follows by (6), but isn't clear if they are equivalent.

In the proof of Theorem 1 we also use the following variant of Jensen inequality, see e.g. [10, Theorem 3.1].

**Lemma 2.** *Let  $\Phi : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  be a convex function and  $w : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  be a locally integrable function such that  $0 < \int_0^\infty w(t) dt < \infty$ . If  $w\Phi \in L^1(\mathbf{R}_+)$  and  $f : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  such that the map  $t \mapsto \Phi(f(t))$  belongs to  $L^1(\mathbf{R}_+)$  then*

$$(10) \quad \Phi\left(\frac{\int_0^\infty w(t)f(t) dt}{\int_0^\infty w(t) dt}\right) \leq \frac{\int_0^\infty w(t)\Phi(f(t)) dt}{\int_0^\infty w(t) dt}.$$

*Proof of Theorem 1.* Let  $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\}$  be an evolution family of bounded linear operators on  $X$ . We consider the evolution semigroup associated to  $\mathcal{U}$  on  $L^1(\mathbf{R}_+, X)$ . This semigroup is defined by

$$(11) \quad (\mathfrak{T}(t)f)(s) := \begin{cases} U(s, s-t)f(s-t), & \text{if } s \geq t \\ 0, & \text{if } 0 \leq s \leq t \end{cases}, \quad t \geq 0$$

for all  $f \in L^1(\mathbf{R}_+, X)$ . Firstly we will prove that  $\mathfrak{T}(t)$  acts on  $L^1(\mathbf{R}_+, X)$  for each  $t \geq 0$ . Indeed, if  $f_n$  are simple functions and  $f_n$  converges punctually almost everywhere to  $f$  on  $\mathbf{R}_+$  when  $n \rightarrow \infty$ , then using the measurability of the functions given in (1) it follows that for all  $n \in \mathbf{N}$  the maps  $\mathfrak{T}(t)f_n$  are measurable. From (4) and Lemma 1 it follows that

$$\|(\mathfrak{T}(t)f_n)(s) - (\mathfrak{T}f)(s)\| \leq K\|f_n(s-t) - f(s-t)\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

almost everywhere for  $s \in [t, \infty)$ , that is, the function in (11) is measurable.

On the other hand

$$\int_0^\infty \|(\mathfrak{T}f)(s)\| ds = \int_t^\infty \|U(s, s-t)f(s-t)\| dt \leq C\|f\|_{L^1(\mathbf{R}_+, X)} < \infty,$$

i.e. the function  $\mathfrak{T}(t)f$  belongs to  $L^1(\mathbf{R}_+, X)$ . The functions defined in (5) are measurable for each  $f \in L^1(\mathbf{R}_+, X)$ , hence the maps

$$(12) \quad t \mapsto \mathfrak{T}(t)f : \mathbf{R}_+ \rightarrow L^1(\mathbf{R}_+, X)$$

are also measurable. We do not know at this stage if the measurability of the function in (12) can be obtained using only the measurability of functions in (1). We may suppose that  $\varphi(1) = 1$  (if not, we replace  $\varphi$  by some multiple of itself). The function

$$t \mapsto \Phi(t) := \int_0^t \varphi(u) du : \mathbf{R}_+ \rightarrow \mathbf{R}_+$$

is convex, continuous on  $(0, \infty)$  and strictly increasing. Moreover

$$\Phi(t) \leq \varphi(t) \quad \text{for all } t \in [0, 1].$$

Without loss of generality we may assume that

$$\sup_{t \geq 0} \|\mathfrak{T}(t)\| \leq 1.$$

Let  $f \in C_c((0, \infty), X)$ , the space of all continuous,  $X$ -valued functions defined on  $\mathbf{R}_+$  with compact support in  $(0, \infty)$ , such that

$$\|f\|_\infty := \sup\{\|f(t)\| : t \in (0, \infty)\} \leq 1.$$

Using Lemma 2 (in inequality (10) we replace  $w(\cdot)$  by  $\exp_{-1}$ ) and the Fubini Theorem it follows that

$$\begin{aligned}
 \int_0^\infty \Phi \left( \|\mathfrak{T}(t) \exp_{-1} \cdot f\|_{L^1(\mathbf{R}_+, X)} \right) dt &= \int_0^\infty \Phi \left( \int_t^\infty e^{-(s-t)} \|U(s, s-t) f(s-t)\| ds \right) dt \\
 &= \int_0^\infty \Phi \left( \int_0^\infty e^{-\xi} \|U(t+\xi, \xi) f(\xi)\| d\xi \right) dt \\
 &\leq \int_0^\infty \left( \int_0^\infty e^{-\xi} \Phi(\|U(t+\xi, \xi) f(\xi)\|) dt \right) d\xi \\
 &\leq \int_0^\infty \left( \int_0^\infty e^{-\xi} \varphi(\|U(t+\xi, \xi) f(\xi)\|) dt \right) d\xi \\
 &\leq M_\varphi \int_0^\infty e^{-\xi} \|f(\xi)\| d\xi \\
 &\leq M_\varphi < \infty.
 \end{aligned}$$

Let  $g \in L^1(\mathbf{R}_+, X)$  with  $\|g\|_{L^1(\mathbf{R}_+, X)} \leq 1$  and  $f_n \in C_c((0, \infty), X)$  such that  $\|f_n\|_\infty \leq 1$  and  $\exp_{-1} \cdot f_n \rightarrow g$  in  $L^1(\mathbf{R}_+, X)$ . Let  $(f_{n_k})$  be a subsequence of  $(f_n)$  such that  $\exp_{-1} \cdot f_{n_k}$  converges at  $g$ , punctually almost everywhere for  $t \in \mathbf{R}_+$ , when  $k \rightarrow \infty$ . Using the above estimates with  $f$  replaced by  $f_{n_k}$ , and the Dominated Convergence Theorem, it follows that

$$\int_0^\infty \Phi(\|\mathfrak{T}(t)g\|_{L^1(\mathbf{R}_+, X)}) dt \leq M_\varphi < \infty.$$

The assertion of Theorem 1, follows now, using a variant of Neerven's result (see the beginning of our note). We recall that  $\mathcal{U}$  is uniformly exponentially stable if and only if  $\omega_0(\mathfrak{T})$  is negative [15, Theorem 2.2].

■

See also [14], [13] and the references therein for more details about the evolution semigroups on half line and their connections with asymptotic behaviour of evolution families of bounded linear operators acting on Banach spaces.

#### REFERENCES

- [1] C. BUŞE AND S. S. DRAGOMIR, A Theorem of Rolewicz's type in Solid Function Spaces, *Glasgow Mathematical Journal*, to appear.
- [2] R. DATKO, Extending a theorem of A.M. Liapanov to Hilbert space, *J. Math. Anal. Appl.*, **32** (1970), 610-616.
- [3] R. NAGEL (ed), *One Parameter Semigroups of Positive Operators*, Lecture Notes in Math., no. **1184**, Springer-Verlag, Berlin, 1984.
- [4] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer Verlag, 1983.
- [5] S. ROLEWICZ, On uniform  $N$ -equistability, *J. Math. Anal. Appl.*, **115** (1886) 434-441.
- [6] S. ROLEWICZ, *Functional Analysis and Control Theory*, D. Riedel and PWN-Polish Scientific Publishers, Dordrecht-Warszawa, 1985.
- [7] Q. ZHENG, The exponential stability and the perturbation problem of linear evolution systems in Banach spaces, *J. Sichuan Univ.*, **25** (1988), 401-411.

- [8] W. LITTMAN, A generalisation of a theorem of Datko and Pazy, *Lect. Notes in Control and Inform. Sci.*, **130**, Springer Verlag (1989), 318-323.
- [9] J.M.A.M. van NEERVEN, *The Asymptotic Behaviour of Semigroups of Linear Operators*, Birkhäuser Verlag Basel (1996).
- [10] J. E. PECÁRIĆ, F. PROSCHAN, Y. L. TONG, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press (1992).
- [11] K.M. PRZYŁUSKI, On a discrete time version of a problem of A.J. Pritchard and J. Zabczyk, *Proc. Roy. Soc. Edinburgh, Sect. A*, **101** (1985), 159-161.
- [12] A. ZABCZYK, Remarks on the control of discrete-time distributed parameter systems, *SIAM J. Control*, **12** (1974), 731-735.
- [13] N.V. MINH, F. RÄBINGER and R. SCHNAUBELT, Exponential stability, exponential expansiveness, and exponential dichotomy of evolution equations on the half-line, *Integral Eqns. Oper. Theory*, **3R** (1998), 332-353.
- [14] C. CHICONE and Y. LATUSHKIN, *Evolution Semigroups in Dynamical Systems and Differential Equations*, Mathematical Surveys and Monographs, Vol. **70**, Amer. Math. Soc., Providence, RI, 1999.
- [15] S. CLARK, Y. LATUSHKIN, S. MONTGOMERY-SMITH and T. RANDOLPH, Stability radius and internal versus external stability in Banach spaces: An evolution semigroup approach, *SIAM Journal of Control and Optim.*, **38**(6) (2000), 1757-1793.
- [16] G. WEISS, Weakly  $l^p$ -stable linear operators are power stable, *Int. J. Systems Sci.*, **20**(1989), 2323-2328.

DEPARTMENT OF MATHEMATICS, WEST UNIVERSITY OF TIMIŞOARA, Bd. V. PARVAN 4, 1900 TIMIŞOARA, ROMÂNIA.

*E-mail address:* `buse@tim1.math.uvt.ro`

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MELBURNE CITY MC 8001., VICTORIA, AUSTRALIA.

*E-mail address:* `sever@matilda.vu.edu.au`

*URL:* <http://rgmia.vu.edu.au/SSDragomirWeb.html>