

SOME REFINEMENTS OF HILBERT-PACHPATTE TYPE INTEGRAL INEQUALITIES

SEVER S. DRAGOMIR AND YOUNG-HO KIM

ABSTRACT. In this paper, we obtain an extension of a multivariable integral inequality of Hilbert-Pachpatte type. By specializing the upper estimate functions in the hypothesis and the parameters, we obtain many special cases.

1. INTRODUCTION

Hilbert's double series theorem [3, p.226] was proved first by Hilbert in his lectures on integral equations. The determination of the constant, the integral analogue, the extension, other proofs of the whole or of parts of the theorems and generalizations in different directions have been given by several authors (cf. [3, Chap. 9]). In particular, in [10-14] the present author has established some new inequalities similar to Hilbert's double-series inequality and its integral analogue which we believe will serve as a model for further investigation. Recently, G. D. Handley, J. J. Koliha and J. E. Pečarić [2] established a new class of related integral inequalities from which results of Pachpatte [12-14] are obtained by specializing the parameters and the functions Φ_i . A representative sample is the following.

Theorem 1.1 (Handley, Koliha and Pečarić [2, Theorem 3.1]). *Let $u_i \in C^{m_i}([0, x_i])$ for $i \in I$. If*

$$|u_i^{(k_i)}(s_i)| \leq \int_0^{(s_i)} (s_i - \tau_i)^{m_i - k_i - 1} \Phi_i(\tau_i) d\tau_i, \quad s_i \in [0, x_i], i \in I,$$

1991 *Mathematics Subject Classification.* 26 D 15.

Key words and phrases. Hilbert's inequality, Hilbert-Pachpatte type Inequality, Hölder's inequality, Jensen inequality.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - TEX

then

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)/(q_i \omega_i)}} ds_1 \cdots ds_n \\ & \leq U \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} \Phi_i(s_i)^{p_i} ds_i \right)^{1/p_i}, \end{aligned}$$

where

$$U = \frac{1}{\prod_{i=1}^n [(\alpha_i + 1)^{1/q_i} (\beta_i + 1)^{1/p_i]}.$$

The purpose of the present paper is to derive an extension of the inequality given in Theorem 1.1. In addition, we obtain some new inequalities as Hilbert-Pachpatte type inequalities, these inequalities improve the results obtained by Handley, Koliha and Pečarić [2].

2. MAIN RESULTS

In what follows we denote by R the set of real numbers; R_+ denotes the interval $[0, \infty)$. The symbols N, Z have their usual meaning. The following notation and hypotheses will be used throughout the paper:

$$\begin{aligned} I = \{1, \dots, n\} & \quad n \in N \\ m_i, i \in I & \quad m_i \in N \\ k_i, i \in I & \quad k_i \in \{0, 1, \dots, m_i - 1\} \\ x_i, i \in I & \quad x_i \in R, x_i > 0 \\ p_i, q_i, i \in I & \quad p_i, q_i \in R_+, 1/p_i + 1/q_i = 1 \\ p, q & \quad 1/p = \sum_{i=1}^n (1/p_i), 1/q = \sum_{i=1}^n (1/q_i) \\ a_i, b_i, i \in I & \quad a_i, b_i \in R_+, a_i + b_i = 1 \\ \omega_i, i \in I & \quad \omega_i \in R, \omega_i > 0, \sum_{i=1}^n \omega_i = \Omega_n \\ \alpha_i, i \in I & \quad \alpha_i = (a_i + b_i q_i)(m_i - k_i - 1) \\ \beta_i, i \in I & \quad \beta_i = a_i(m_i - k_i - 1) \\ u_i, i \in I & \quad u_i \in C^{m'_i}([0, x_i]) \quad \text{for some } m'_i \geq m_i \\ \Phi_i, i \in I & \quad \Phi_i \in C^1([0, x_i]), \Phi_i \geq m_i \end{aligned}$$

Here the u_i are given functions of sufficient smoothness, and the Φ_i are subject to choice. The coefficients p_i, q_i are conjugate Hölder exponents to be used in applications of Hölder's inequality, and the coefficients a_i, b_i will be used in exponents to factorize integrands. The coefficients ω_i will act as weights in applications of the geometric-arithmetic mean inequality. The coefficients α_i and β_i arise naturally in the derivation of the inequalities.

Our main results are given in the following theorems.

Theorem 2.1. *Let $u_i \in C^{m_i}([0, x_i])$ for $i \in I$. If*

$$(2.1) \quad |u_i^{(k_i)}(s_i)| \leq \int_0^{(s_i)} (s_i - \tau_i)^{m_i - k_i - 1} \Phi_i(\tau_i) d\tau_i, \quad s_i \in [0, x_i], i \in I,$$

then

$$(2.2) \quad \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i + 1)\Omega_n / (q_i \omega_i)}} ds_1 \cdots ds_n \\ \leq V \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i + 1} \Phi_i(s_i)^{p_i} ds_i \right)^{1/p_i},$$

where

$$(2.3) \quad V = \left(\prod_{i=1}^n [(\alpha_i + 1)^{1/q_i} (\beta_i + 1)^{1/p_i}] \right)^{-1}.$$

Proof. Factorize the integration on the right side of (2.1) as

$$(s_i - \tau_i)^{(a_i/q_i + b_i)(m_i - k_i - 1)} \times (s_i - \tau_i)^{(a_i/p_i)(m_i - k_i - 1)} \Phi_i(\tau_i)$$

and apply Hölder's inequality [9, p.106]. Then

$$|u_i^{(k_i)}(s_i)| \leq \left(\int_0^{(s_i)} (s_i - \tau_i)^{(a_i + b_i q_i)(m_i - k_i - 1)} d\tau_i \right)^{1/q_i} \\ \times \left(\int_0^{(s_i)} (s_i - \tau_i)^{a_i(m_i - k_i - 1)} \Phi_i(\tau_i)^{p_i} d\tau_i \right)^{1/p_i} \\ = \frac{s_i^{(\alpha_i + 1)/q_i}}{(\alpha_i + 1)^{1/q_i}} \left(\int_0^{(s_i)} (s_i - \tau_i)^{\beta_i} \Phi_i(\tau_i)^{p_i} d\tau_i \right)^{1/p_i}.$$

Using the inequality of means [9, p. 15]

$$\left(\prod_{i=1}^n s_i^{w_i} \right)^{1/\Omega_n} \leq \left(\frac{1}{\Omega_n} \sum_{i=1}^n w_i s_i^r \right)^{1/r}$$

for $r > 0$, we have

$$\begin{aligned} & \prod_{i=1}^n |u_i^{(k_i)}(s_i)| \\ & \leq W \frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)\Omega_n/(q_i\omega_i)} \prod_{i=1}^n \left(\int_0^{(s_i)} (s_i - \tau_i)^{\beta_i} \Phi_i(\tau_i)^{p_i} d\tau_i \right)^{1/p_i}, \end{aligned}$$

where

$$W = \left(\prod_{i=1}^n (\alpha_i + 1)^{q_i} \right)^{-1}.$$

In the following estimate we apply Hölder's inequality and, at the end, change the order of integration:

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)\Omega_n/(q_i\omega_i)}} ds_1 \cdots ds_n \\ & \leq W \prod_{i=1}^n \left[\int_0^{(x_i)} \left(\int_0^{(s_i)} (s_i - \tau_i)^{\beta_i} \Phi_i(\tau_i)^{p_i} d\tau_i \right)^{1/p_i} ds_i \right] \\ & \leq W \prod_{i=1}^n x_i^{1/q_i} \left[\int_0^{(x_i)} \left(\int_0^{(s_i)} (s_i - \tau_i)^{\beta_i} \Phi_i(\tau_i)^{p_i} d\tau_i \right) ds_i \right]^{1/p_i} \\ & = \frac{W}{\prod_{i=1}^n (\beta_i + 1)^{1/p_i}} \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} \Phi_i(s_i)^{p_i} ds_i \right)^{1/p_i}. \end{aligned}$$

This proves the theorem.

Remark. In Theorem 2.1, setting $\Omega_n = 1$, we have Theorem 1.1.

Corollary 2.2. *Under the assumptions of Theorem 2.1, if $r > 0$, we have*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)\Omega_n/(q_i\omega_i)}} ds_1 \cdots ds_n \\ & \leq p^{1/r \cdot p} V \prod_{i=1}^n x_i^{1/q_i} \left(\sum_{i=1}^n \frac{1}{p_i} \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} \Phi_i(s_i)^{p_i} ds_i \right)^r \right)^{1/r \cdot p}, \end{aligned}$$

Proof. By the inequality of means, for any $A_i \geq 0$ and $r > 0$, we obtain

$$\prod_{i=1}^n A_i^{1/p_i} \leq \left(p \sum_{i=1}^n \frac{1}{p_i} A_i^r \right)^{1/r \cdot p}.$$

The corollary then follows from the preceding theorem.

Lemma 2.3. *Let $\gamma_1 > 0$ and $\gamma_2 < -1$. Let $0 \leq \omega_i, \sum_{i=1}^n \omega_i = \Omega_n$ and let $s_i > 0, i = 1, \dots, n$ be real numbers. Then*

$$\prod_{i=1}^n s_i^{\omega_i \gamma_1 \gamma_2} \geq \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-\gamma_1 \Omega_n}.$$

Proof. By the inequality of means, for any $\gamma_1 > 0$ and $\gamma_2 < -1$,

$$\prod_{i=1}^n s_i^{\omega_i \gamma_1 \gamma_2} \geq \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i \right)^{\gamma_1 \gamma_2 \Omega_n}.$$

Using the fact that x^{-1/γ_2} is concave and using the Jensen inequality and we have that

$$\begin{aligned} \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i \right)^{\gamma_1 \gamma_2 \Omega_n} &= \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i f(s_i^{-\gamma_2}) \right)^{\gamma_1 \gamma_2 \Omega_n} \\ &\geq \left(f \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right) \right)^{\gamma_1 \gamma_2 \Omega_n} \\ &= \left(\left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-1/\gamma_2} \right)^{\gamma_1 \gamma_2 \Omega_n} \\ &= \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-\gamma_1 \Omega_n} \end{aligned}$$

The proof of Lemma is complete.

Theorem 2.4. *Under the assumptions of Theorem 2.1, if $\gamma_2 < -1$, then*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2}\right)^{-(\alpha_i+1)\Omega_n/\gamma_2 q_i \omega_i}} ds_1 \cdots ds_n \\ & \leq V \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} \Phi_i(s_i)^{p_i} ds_i \right)^{1/p_i}, \end{aligned}$$

where V is given by (2.3).

Proof. By the inequality of Lemma 2.3, for any $\gamma_1 > 0$ and $\gamma_2 < -1$,

$$\prod_{i=1}^n s_i^{\omega_i \gamma_1} \leq \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-\gamma_1 \Omega_n / \gamma_2}$$

we get

$$\begin{aligned} & \prod_{i=1}^n |u_i^{(k_i)}(s_i)| \\ & \leq W \left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-W_1} \prod_{i=1}^n \left(\int_0^{(s_i)} (s_i - \tau_i)^{\beta_i} \Phi_i(\tau_i)^{p_i} d\tau_i \right)^{1/p_i}, \end{aligned}$$

where

$$W = \left(\prod_{i=1}^n (\alpha_i + 1)^{q_i} \right)^{-1}, \quad W_1 = \frac{(\alpha_i + 1)\Omega_n}{\gamma_2 q_i \omega_i}.$$

The Theorem then follows from the preceding Theorem 2.1.

Corollary 2.5. *Under the assumptions of Theorem 2.4, if $r > 0$, we have*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2}\right)^{-(\alpha_i+1)\Omega_n/\gamma_2 q_i \omega_i}} ds_1 \cdots ds_n \\ & \leq p^{1/r \cdot p} V \prod_{i=1}^n x_i^{1/q_i} \left(\sum_{i=1}^n \frac{1}{p_i} \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} \Phi_i(s_i)^{p_i} ds_i \right)^r \right)^{1/r \cdot p}, \end{aligned}$$

where V is given by (2.3).

Proof. By the inequality of means, for any $A_i \geq 0$ and $r > 0$, we obtain

$$\prod_{i=1}^n A_i^{1/p_i} \leq \left(p \sum_{i=1}^n \frac{1}{p_i} A_i^r \right)^{1/r \cdot p}.$$

The corollary then follows from the preceding Theorem 2.4.

In the following section we discuss some choice of the functions Φ_i .

3. THE VARIOUS INEQUALITIES

Theorem 3.1. Let $u_i \in C^{m_i}([0, x_i])$ be such that $u_i^{(j)}(0) = 0$ for $j \in \{0, \dots, m_i - 1\}$, $i \in I$. Then

$$(3.1) \quad \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)\Omega_n/(q_i \omega_i)}} ds_1 \cdots ds_n \\ \leq V_1 \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^{1/p_i},$$

where

$$(3.2) \quad V_1 = \left(\prod_{i=1}^n \left((m_i - k_i - 1)! (\alpha_i + 1)^{1/q_i} (\beta_i + 1)^{1/p_i} \right) \right)^{-1}.$$

Proof. By [12, Eq. (7)],

$$u_i^{(k_i)}(s_i) = \frac{1}{(m_i - k_i - 1)!} \int_0^{s_i} (s_i - \tau_i)^{m_i - k_i - 1} u_i^{(m_i)}(\tau_i) d\tau_i.$$

Inequality (3.1) is proved when we set

$$\Phi_i(s_i) = \frac{|u_i^{(m_i)}(s_i)|}{(m_i - k_i - 1)!}$$

in Theorem 2.1.

Corollary 3.2. *Under the assumptions of Theorem 3.1, if $r > 0$, we have*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(\alpha_i+1)\Omega_n/(q_i\omega_i)}} ds_1 \cdots ds_n \\ & \leq p^{1/r \cdot p} V_1 \prod_{i=1}^n x_i^{1/q_i} \left(\sum_{i=1}^n \frac{1}{p_i} \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^r \right)^{1/r \cdot p}, \end{aligned}$$

where V_1 is given by (3.2).

Theorem 3.3. *Under the assumptions of Theorem 3.1, if $\gamma_2 < -1$, then*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-(\alpha_i+1)\Omega_n/\gamma_2 q_i \omega_i}} ds_1 \cdots ds_n \\ (3.3) \quad & \leq V_1 \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^{1/p_i}, \end{aligned}$$

where V_1 is given by (3.2).

Proof. Inequality (3.3) is proved when we set

$$\Phi_i(s_i) = \frac{|u_i^{(m_i)}(s_i)|}{(m_i - k_i - 1)!}$$

in Theorem 2.4.

Corollary 3.4. *Under the assumptions of Theorem 3.3, if $r > 0$, we have*

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\left(\frac{1}{\Omega_n} \sum_{i=1}^n \omega_i s_i^{-\gamma_2} \right)^{-(\alpha_i+1)\Omega_n/\gamma_2 q_i \omega_i}} ds_1 \cdots ds_n \\ & \leq p^{1/r \cdot p} V_1 \prod_{i=1}^n x_i^{1/q_i} \left(\sum_{i=1}^n \frac{1}{p_i} \left(\int_0^{x_i} (x_i - s_i)^{\beta_i+1} |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^r \right)^{1/r \cdot p}. \end{aligned}$$

We discuss a number of special case of Theorem 3.1. Similar examples apply also to Corollary 3.2, Theorem 3.3 and Corollary 3.4.

Example 3.1. If $a_i = 0$ and $b_i = 1$ for $i \in I$, then Theorem 3.1 becomes

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(q_i m_i - q_i k_i - q_i + 1) \Omega_n / (q_i \omega_i)}} ds_1 \cdots ds_n \\ & \leq V_2 \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i) |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^{1/p_i}, \end{aligned}$$

where

$$V_2 = \left(\prod_{i=1}^n \left((m_i - k_i - 1)! (q_i m_i - q_i k_i - q_i + 1)^{1/q_i} \right) \right)^{-1}.$$

Example 3.2. If $a_i = 0$, $b_i = 1$, $q_i = n$, $p_i = n/(n-1)$, $m_i = m$ and $k_i = k$ for $i \in I$, then

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(nm - nk - n + 1) \Omega_n / (m \omega_i)}} ds_1 \cdots ds_n \\ & \leq \frac{\sqrt[n]{x_1 \cdots x_n}}{\left((m - k - 1)! \right)^n (nm - nk - n + 1)} \\ & \quad \times \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i) |u_i^{(m)}(s_i)|^{n/(n-1)} ds_i \right)^{(n-1)/n}. \end{aligned}$$

For $q = p = n = 2$ and $\omega_i = 1/n$ this is [12, Theorem 1]. Setting $q = p = 2$, $k = 0$, $n = 1$ and $\omega_i = 1/n$, we recover the result of [14].

Example 3.3. If $a_i = 0$ and $b_i = 1$ for $i \in I$, then Theorem 3.1 becomes

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(m_i - k_i) \Omega_n / (q_i \omega_i)}} ds_1 \cdots ds_n \\ & \leq V_3 \prod_{i=1}^n x_i^{1/q_i} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{m_i - k_i} |u_i^{(m_i)}(s_i)|^{p_i} ds_i \right)^{1/p_i}, \end{aligned}$$

where

$$V_3 = \left(\prod_{i=1}^n \left((m_i - k_i)! \right) \right)^{-1}.$$

Example 3.4. If $a_i = 1$, $b_i = 0$, $q_i = n$, $p_i = n/(n-1)$, $m_i = m$ and $k_i = k$ for $i \in I$. Then (3.1) becomes

$$\begin{aligned} & \int_0^{x_1} \cdots \int_0^{x_n} \frac{\Omega_n \prod_{i=1}^n |u_i^{(k_i)}(s_i)|}{\sum_{i=1}^n \omega_i s_i^{(m-k)\Omega_n/(n\omega_i)}} ds_1 \cdots ds_n \\ & \leq \frac{\sqrt[n]{x_1 \cdots x_n}}{((m-k)!)^n} \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i)^{m-k} |u_i^{(m)}(s_i)|^{n/(n-1)} ds_i \right)^{(n-1)/n}. \end{aligned}$$

Example 3.5. Let $p_1, p_2 \in R_+$. If we set $n = 2$, $\omega_1 = 1/p_1$, $\omega_2 = 1/p_2$, $m_i = 1$ and $k_i = 0$ for $i = 1, 2$ in Theorem 3.1, then by our assumptions $q_1 = (p_1 - 1)/p_1$, $q_2 = (p_2 - 1)/p_2$, and we obtain

$$\begin{aligned} & \int_0^{x_1} \int_0^{x_2} \frac{\Omega_2 |u_1(s_1)| |u_2(s_2)|}{p_2 s_1^{\Omega_2(p_1-1)} + p_1 s_2^{\Omega_2(p_2-1)}} ds_1 ds_2 \\ & \leq \frac{x_1^{(p_1-1)/p_1} x_2^{(p_2-1)/p_2}}{p_1 p_2} \left(\int_0^{x_1} (x_1 - s_1) |u_1'(s_1)|^{p_1} ds_1 \right)^{1/p_1} \\ & \quad \times \left(\int_0^{x_2} (x_2 - s_2) |u_2'(s_2)|^{p_2} ds_2 \right)^{1/p_2}. \end{aligned}$$

If we set $\omega_1 + \omega_2 = 1$ in Example 3.5, then we have [13, Theorem 2]. (The values of a_i and b_i are irrelevant.)

REFERENCES

- [1] Yang Bicheng, *On Hilbert's integral inequality*, J. Math. Anal. Appl. **220** (1988), 778-785.
- [2] G. D. Handley, J. J. Koliha and J. E. Pečarić, *New Hilbert-Pachpatte type integral inequalities*, J. Math. Anal. Appl. **257** (2001), 238-250.
- [3] G. H. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*, Cambridge Univ. Press, London, 1952.
- [4] Young-Ho Kim, *Refinements and Extensions of an inequality*, J. Math. Anal. Appl. **245** (2000), 628-632.
- [5] V. Levin, *On the two-parameter extension and analogue of Hilbert's inequality*, J. London Math. Soc. **11** (1936), 119-124.
- [6] G. Mingze, *On Hilbert's inequality and its applications*, J. Math. Anal. Appl. **212** (1997), 316-323.
- [7] D. S. Mitrinović, *Analytic inequalities*, Springer-Verlag, Berlin, New York, 1970.
- [8] D. S. Mitrinović and J. E. Pečarić, *On inequalities of Hilbert and Widder*, Proc. Edinburgh Math. Soc. **34** (1991), 411-414.

- [9] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic, Dordrecht, 1993.
- [10] B. G. Pachpatte, *A note on Hilbert type inequality*, Tamkang J. Math. **29** (1998), 293-298.
- [11] B. G. Pachpatte, *On some new inequalities similar to Hilbert's inequality*, J. Math. Anal. Appl. **226** (1998), 166-179.
- [12] B. G. Pachpatte, *Inequalities similar to the integral analogue of Hilbert's Inequality*, Tamkang J. Math. **30** (1999), 139-146.
- [13] B. G. Pachpatte, *Inequalities similar to certain extensions of Hilbert's inequality*, J. Math. Anal. Appl. **243** (2000), 217-227.
- [14] B. G. Pachpatte, *A note on inequality of Hilbert type*, Demonstratio Math., in press.
- [15] D. V. Widder, *An inequality related to one of Hilbert's*, J. London Math. Soc. **4** (1929), 194-198.

SCHOOL OF COMMUNICATIONS AND INFORMATICS VICTORIA UNIVERSITY OF TECHNOLOGY
PO Box 14428, MCMC MELBOURNE, VICTORIA 8001, AUSTRALIA

E-mail address: sever.dragomir@vu.edu.au

DEPARTMENT OF APPLIED MATHEMATICS BRAIN KOREA 21 PROJECT CORPS CHANGWON
NATIONAL UNIVERSITY, CHANGWON 641-773, KOREA

E-mail address: yhkim@sarim.changwon.ac.kr