

SOME NEW IYENGAR TYPE INEQUALITIES

FENG QI, PIETRO CERONE, AND SEVER S. DRAGOMIR

ABSTRACT. Some new Iyengar type inequalities for an integral are obtained by using the generalised Taylor formula with integral remainder.

1. INTRODUCTION

Let $f(x)$ be a differentiable function on a closed interval $[a, b]$ such that $|f'(x)| \leq M$, then

$$\left| \int_a^b f(x) dx - \frac{(b-a)[f(a) + f(b)]}{2} \right| \leq \frac{(b-a)^2 M}{4} - \frac{[f(b) - f(a)]^2}{4M}. \quad (1)$$

In 1938, K. S. K. Iyengar [15] established inequality (1) by using a geometric approach. So, we call (1) the Iyengar inequality.

Using the Rolle and Lagrange mean value theorems, the following inequalities were obtained naturally and simply in [24], producing a refinement of the Iyengar inequality (1).

Theorem A. *Let $f(x)$ be continuous on the closed interval $[a, b]$ and differentiable in the open interval (a, b) , and $m \leq f'(x) \leq M$ for $x \in (a, b)$. If $f(x)$ is not a constant, then we have*

$$\begin{aligned} \frac{mM(b-a)^2 + 2(b-a)(Mf(a) - mf(b)) + (f(a) - f(b))^2}{2(M-m)} &\leq \int_a^b f(x) dx \\ &\leq -\frac{mM(b-a)^2 + 2(b-a)(mf(a) - Mf(b)) + (f(a) - f(b))^2}{2(M-m)}. \end{aligned} \quad (2)$$

For $f'(x)$ integrable on $[a, b]$, the inequalities in (2) were obtained independently by R.P. Agarwal and S.S. Dragomir in [2] using the Hayashi inequality. They

Date: December 10, 2001.

1991 Mathematics Subject Classification. Primary 26D15; Secondary 41A55.

Key words and phrases. Iyengar type inequality, generalized Taylor formula, Appell condition, harmonic sequence of polynomials, Bernoulli polynomials, Euler polynomials.

The first author was supported in part by NNSF (#10001016) of China, SF for the Prominent Youth of Henan Province, SF of Henan Innovation Talents at Universities, NSF of Henan Province (#004051800), SF for Pure Research of Natural Science of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, China.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

expressed the result (2) as

$$\left| \int_a^b f(t) dt - (b-a) \frac{f(a) + f(b)}{2} \right| \leq \frac{[f(b) - f(a) - m(b-a)][M(b-a) - f(b) + f(a)]}{2(M-m)}. \quad (3)$$

In [3], the inequalities in (2) and in (3) were derived in a different manner and rearranged as

$$\left| \int_a^b f(x) dx - \frac{b-a}{2}[f(a) + f(b)] \right| \leq \frac{(b-a)^2}{2(M-m)}(S-m)(M-S) \quad (4)$$

$$\leq \frac{M-m}{2} \left(\frac{b-a}{2} \right)^2, \quad (5)$$

where $S = \frac{f(b)-f(a)}{b-a}$.

In the papers [25, 26], using the Taylor formula for functions with a single variable or several variables, the Iyengar inequality (1) was generalized as follows.

Theorem B. *Let $f(x)$ be a differentiable function of $C^n([a, b])$ satisfying $N \leq f^{(n+1)}(x) \leq M$ for $x \in (a, b)$. Denote*

$$S_n(u, v, w) = \sum_{k=1}^{n-1} \frac{(-1)^k}{k!} u^k f^{(k-1)}(v) + (-1)^n \frac{w}{n!} u^n \quad (6)$$

$$\text{and } \frac{\partial^k S_n}{\partial u^k} = S_n^{(k)}(u, v, w), \quad (7)$$

then, for any $t \in [a, b]$, we have when n is an odd,

$$\begin{aligned} \sum_{i=0}^{n+2} \frac{(-1)^i}{i!} \left(S_{n+2}^{(i)}(a, a, N) - S_{n+2}^{(i)}(b, b, N) \right) t^i &\leq \int_a^b f(x) x \\ &\leq \sum_{i=0}^{n+2} \frac{(-1)^i}{i!} \left(S_{n+2}^{(i)}(a, a, M) - S_{n+2}^{(i)}(b, b, M) \right) t^i; \end{aligned} \quad (8)$$

and when n is even

$$\begin{aligned} \sum_{i=0}^{n+2} \frac{(-1)^i}{i!} \left(S_{n+2}^{(i)}(a, a, N) - S_{n+2}^{(i)}(b, b, M) \right) t^i &\leq \int_a^b f(x) x \\ &\leq \sum_{i=0}^{n+2} \frac{(-1)^i}{i!} \left(S_{n+2}^{(i)}(a, a, M) - S_{n+2}^{(i)}(b, b, N) \right) t^i. \end{aligned} \quad (9)$$

We may deduce both Theorem A and the following result as specialisations of Theorem B.

Theorem C. Let $f(x)$ be differentiable on the closed interval $[a, b]$, and $N \leq f''(x) \leq M$ for $x \in (a, b)$, then

$$\begin{aligned} & \frac{N(b^3 - a^3)}{6} + \frac{\left\{ f(a) - f(b) + bf'(b) - af'(a) + \frac{N(a^2 - b^2)}{2} \right\}^2}{2[(a-b)N - f'(a) + f'(b)]} \\ & \leq \int_a^b f(x) x - bf(b) + af(a) + \frac{b^2 f'(b) - a^2 f'(a)}{2} \quad (10) \\ & \leq \frac{M(b^3 - a^3)}{6} + \frac{\left\{ f(a) - f(b) + bf'(b) - af'(a) + \frac{M(a^2 - b^2)}{2} \right\}^2}{2[(a-b)M - f'(a) + f'(b)]}. \end{aligned}$$

In recent years, R.P. Agarwal, P. Cerone, S.S. Dragomir and S. Wang, utilizing Hayashi inequality or Steffensen inequality, generalized Iyengar' inequality to those involving higher derivatives, weighted integral, and so on. See [2]–[6], and [10].

Meanwhile, F. Qi, B.-N. Guo, Y.-J. Zhang and L.-H. Cui, proceeded in an alternate direction. Using the mean value theorems for functions with a single variable or several variables, generalized Iyengar inequality to involve bounds of higher order derivatives, with norm bounds, for weighted multiple integral, and the like. See [8, 12, 14] and [24]–[29].

The Iyengar inequality (1) has been researched by many mathematicians, and there is much literature devoted to it. Please refer to references in this paper. Before elaborating on the contributions of the current article, it is useful to present some background notation and definitions.

Definition 1. A sequence of polynomials $\{P_i(t, x)\}_{i=0}^\infty$ is called harmonic if it satisfies the following Appell condition

$$P'_i(t) \triangleq \frac{\partial P_i(t, x)}{\partial t} = P_{i-1}(t, x) \triangleq P_{i-1}(t) \quad (11)$$

and $P_0(t, x) = 1$ for all defined (t, x) and $i \in \mathbb{N}$.

It is well-known that Bernoulli polynomials $B_i(t)$ can be defined by the following expansion

$$\frac{xe^{tx}}{e^x - 1} = \sum_{i=0}^{\infty} \frac{B_i(t)}{i!} x^i, \quad |x| < 2\pi, \quad t \in \mathbb{R}, \quad (12)$$

and are uniquely determined by the following formulae

$$B'_i(t) = iB_{i-1}(t), \quad B_0(t) = 1; \quad (13)$$

$$\text{and } B_i(t+1) - B_i(t) = it^{i-1}. \quad (14)$$

Similarly, Euler polynomials can be defined by

$$\frac{2e^{tx}}{e^x + 1} = \sum_{i=0}^{\infty} \frac{E_i(t)}{i!} x^i, \quad |x| < \pi, \quad t \in \mathbb{R}, \quad (15)$$

and are uniquely determined by the following properties

$$E'_i(t) = iE_{i-1}(t), \quad E_0(t) = 1; \quad (16)$$

$$E_i(t+1) + E_i(t) = 2t^i. \quad (17)$$

For further details about Bernoulli polynomials and Euler polynomials, please refer to [1, 23.1.5 and 23.1.6]. Moreover, some new generalizations of Bernoulli numbers and polynomials can be found in [13, 17, 27].

There are many examples of harmonic sequences of polynomials. For instance ([3, 7, 18]), for i a nonnegative integer, $t, \tau, \theta \in \mathbb{R}$ and $\tau \neq \theta$,

$$P_{i,\lambda}(t) \triangleq P_{i,\lambda}(t; \tau; \theta) = \frac{[t - (\lambda\theta + (1-\lambda)\tau)]^i}{i!}, \quad (18)$$

$$P_{i,B}(t) \triangleq P_{i,B}(t; \tau; \theta) = \frac{(\tau - \theta)^i}{i!} B_i \left(\frac{t - \theta}{\tau - \theta} \right), \quad (19)$$

$$P_{i,E}(t) \triangleq P_{i,E}(t; \tau; \theta) = \frac{(\tau - \theta)^i}{i!} E_i \left(\frac{t - \theta}{\tau - \theta} \right). \quad (20)$$

As usual, let $B_i = B_i(0)$, $i \in \mathbb{N}$, denote Bernoulli numbers. From properties (13) and (14), (16) and (17) of Bernoulli and Euler polynomials respectively, we can obtain easily that, for $i \geq 1$,

$$B_{i+1}(0) = B_{i+1}(1) = B_{i+1}, \quad B_1(0) = -B_1(1) = -\frac{1}{2}, \quad (21)$$

and, for $j \in \mathbb{N}$,

$$E_j(0) = -E_j(1) = -\frac{2}{j+1}(2^{j+1} - 1)B_{j+1}. \quad (22)$$

It is also a well known fact that $B_{2i+1} = 0$ for all $i \in \mathbb{N}$.

In [18], the following generalized Taylor formula was established.

Theorem D. *Let $\{P_i(x)\}_{i=0}^{\infty}$ be a harmonic sequence of polynomials. Further, let $I \subset \mathbb{R}$ be a closed interval and $a \in I$. If $f : I \rightarrow \mathbb{R}$ is any function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, then, for any $x \in I$, we have*

$$f(x) = f(a) + \sum_{k=1}^n (-1)^{k+1} [P_k(x)f^{(k)}(x) - P_k(a)f^{(k)}(a)] + R_n(f; a, x), \quad (23)$$

where

$$R_n(f; a, x) = (-1)^n \int_a^x P_n(t) f^{(n+1)}(t) dt. \quad (24)$$

If we set in (23) and (24) $P_n(t) = \frac{(t-x)^n}{n!}$, then we obtain directly the classical Taylor formula with remainder of integral form. Generalised Taylor formulae from product Appel polynomials was considered in [4] which involves the product of polynomials satisfying the Appel condition (1).

In this article, using the generalized Taylor formula (23), we will derive some new Iyengar type inequalities for an integral of functions with single variable, which generalize some related known results obtained in [8, 12, 14, 15, 24, 25, 28], for example.

2. SOME INTEGRAL IDENTITIES

In this section, we establish two identities involving integrals which form the basis for the procurement of our main results.

Theorem 1. Let $\{P_i(x)\}_{i=0}^{\infty}$ and $\{Q_i(x)\}_{i=0}^{\infty}$ be harmonic sequences of polynomials. If $f : [a, b] \rightarrow \mathbb{R}$ is a function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, then we have the following generalized Taylor identity

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= b \sum_{k=0}^n q(k, k; b) - a \sum_{k=0}^n p(k, k; a) \\ &\quad + \sum_{k=1}^n \sum_{i=1}^k [p(i, i-1, a) - q(i, i-1; b)] \\ &\quad + t \sum_{k=0}^n [p(k, k; a) - q(k, k; b)] + \sum_{k=1}^n \sum_{i=1}^k [q(i, i-1, t) - p(i, i-1; t)] \\ &\quad + \int_a^t (t-s)p(n, n+1; s) ds + \int_t^b (t-s)q(n, n+1; s) ds, \end{aligned} \quad (25)$$

where $t \in [a, b]$ and

$$p(\ell, m; x) = (-1)^\ell P_\ell(x) f^{(m)}(x), \quad (26)$$

$$q(\ell, m; x) = (-1)^\ell Q_\ell(x) f^{(m)}(x) \quad (27)$$

for any nonnegative integers $0 \leq \ell \leq n$ and $0 \leq m \leq n+1$ and $x \in [a, b]$.

Proof. Let t be a parameter such that $a \leq t \leq b$, then

$$\int_a^b f(x) dx = \int_a^t f(x) dx + \int_t^b f(x) dx. \quad (28)$$

From the generalized Taylor formula (23), for $x \in [a, b]$, it follows that

$$f(x) = f(a) + \sum_{k=1}^n (-1)^{k+1} [P_k(x) f^{(k)}(x) - P_k(a) f^{(k)}(a)] + R_{n,P}(f; a, x), \quad (29)$$

$$f(x) = f(b) + \sum_{k=1}^n (-1)^{k+1} [Q_k(x) f^{(k)}(x) - Q_k(b) f^{(k)}(b)] + R_{n,Q}(f; b, x). \quad (30)$$

By integration by parts we have

$$\begin{aligned} &\int_a^t P_k(x) f^{(k)}(x) dx \\ &= [P_k(t) f^{(k-1)}(t) - P_k(a) f^{(k-1)}(a)] - \int_a^t P_{k-1}(x) f^{(k-1)}(x) dx. \end{aligned}$$

Clearly, we can apply the same procedure to the term $\int_a^t P_{k-1}(x) f^{(k-1)}(x) dx$. So, by successive integration by parts we obtain

$$\begin{aligned} &(-1)^k \int_a^t P_k(x) f^{(k)}(x) dx \\ &= \sum_{i=1}^k (-1)^i [P_i(t) f^{(i-1)}(t) - P_i(a) f^{(i-1)}(a)] + \int_a^t f(x) dx. \end{aligned} \quad (31)$$

Similarly, we have

$$\begin{aligned} & (-1)^k \int_t^b Q_k(x) f^{(k)}(x) dx \\ &= \sum_{i=1}^k (-1)^i [Q_i(b) f^{(i-1)}(b) - Q_i(t) f^{(i-1)}(t)] + \int_t^b f(x) dx. \end{aligned} \quad (32)$$

Integrating both sides of formula (29) over the interval $[a, t]$ and utilizing identity (31) yields

$$\begin{aligned} (n+1) \int_a^t f(x) dx &= (t-a) \sum_{k=0}^n (-1)^k P_k(a) f^{(k)}(a) \\ &+ \sum_{k=1}^n \sum_{i=1}^k (-1)^{i+1} [P_i(t) f^{(i-1)}(t) - P_i(a) f^{(i-1)}(a)] \\ &+ (-1)^n \int_a^t \int_a^x P_n(s) f^{(n+1)}(s) ds dx. \end{aligned} \quad (33)$$

Similarly, integrating both sides of the result (30) over the interval $[t, b]$ and using the identity (32) gives

$$\begin{aligned} (n+1) \int_t^b f(x) dx &= (b-t) \sum_{k=0}^n (-1)^k Q_k(b) f^{(k)}(b) \\ &+ \sum_{k=1}^n \sum_{i=1}^k (-1)^{i+1} [Q_i(b) f^{(i-1)}(b) - Q_i(t) f^{(i-1)}(t)] \\ &+ (-1)^n \int_t^b \int_b^x Q_n(s) f^{(n+1)}(s) ds dx. \end{aligned} \quad (34)$$

Now, combining (33) and (34) and utilising (28) produces

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= b \sum_{k=0}^n (-1)^k Q_k(b) f^{(k)}(b) - a \sum_{k=0}^n (-1)^k P_k(a) f^{(k)}(a) \\ &+ \sum_{k=1}^n \sum_{i=1}^k (-1)^{i+1} [Q_i(b) f^{(i-1)}(b) - P_i(a) f^{(i-1)}(a)] \\ &+ t \sum_{k=0}^n (-1)^{k+1} [Q_k(b) f^{(k)}(b) - P_k(a) f^{(k)}(a)] \\ &+ \sum_{k=1}^n \sum_{i=1}^k (-1)^i [Q_i(t) - P_i(t)] f^{(i-1)}(t) \\ &+ (-1)^n \left[\int_a^t \int_a^x P_n(s) f^{(n+1)}(s) ds dx + \int_t^b \int_b^x Q_n(s) f^{(n+1)}(s) ds dx \right]. \end{aligned} \quad (35)$$

Further, interchanging the order of integration for the last two terms in (35) leads to

$$\int_a^t \int_a^x P_n(s) f^{(n+1)}(s) ds dx = \int_a^t (t-s) P_n(s) f^{(n+1)}(s) ds, \quad (36)$$

$$\int_t^b \int_b^x Q_n(s) f^{(n+1)}(s) ds dx = \int_t^b (t-s) Q_n(s) f^{(n+1)}(s) ds. \quad (37)$$

Substituting (36) and (37) into (35), rearranging and introducing the notation in (26) and (27) produces the identity (25). The proof is complete. ■

Remark 1. If we set

$$P_i(s) = Q_i(s) = \frac{(s-x)^i}{i!} \quad (38)$$

for i a nonnegative natural number in (29) and (30) from the very start, then following the same procedure as in the proof of Theorem 1, we would obtain

$$\int_a^t f(x) dx = \sum_{i=0}^n \frac{f^{(i)}(a)}{(i+1)!} (t-a)^{i+1} + \frac{1}{(n+1)!} \int_a^t (t-s)^{n+1} f^{(n+1)}(s) ds, \quad (39)$$

$$\int_t^b f(x) dx = -\sum_{i=0}^n \frac{f^{(i)}(b)}{(i+1)!} (t-b)^{i+1} + \frac{1}{(n+1)!} \int_t^b (t-s)^{n+1} f^{(n+1)}(s) ds, \quad (40)$$

the combination of which, produces

$$\begin{aligned} \int_a^b f(x) dx &= \sum_{i=0}^n \frac{f^{(i)}(a)}{(i+1)!} (t-a)^{i+1} - \sum_{i=0}^n \frac{f^{(i)}(b)}{(i+1)!} (t-b)^{i+1} \\ &\quad + \frac{1}{(n+1)!} \int_a^b (t-s)^{n+1} f^{(n+1)}(s) ds. \end{aligned} \quad (41)$$

The following result is a particularisation of Theorem 1, but because of its intrinsic importance is denoted as a theorem in its own right, rather than a corollary.

Theorem 2. *Let $\{P_i(x)\}_{i=0}^{\infty}$ be a harmonic sequence of polynomials. If $f : [a, b] \rightarrow \mathbb{R}$ is a function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, then we have the following identity*

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= b \sum_{k=0}^n p(k, k; b) - a \sum_{k=0}^n p(k, k; a) \\ &\quad - \sum_{k=1}^n \sum_{i=1}^k [p(i, i-1; b) - p(i, i-1; a)] - t \sum_{k=0}^n [p(k, k; b) - p(k, k; a)] \\ &\quad + \int_a^b (t-s) p(n, n+1; s) ds, \end{aligned} \quad (42)$$

where $t \in [a, b]$ and $p(\ell, m; x) = (-1)^\ell P_\ell(x) f^{(m)}(x)$, $x \in [a, b]$, for any nonnegative integers $0 \leq \ell \leq n$ and $0 \leq m \leq n+1$.

Proof. This follows from taking $P_i(s) = Q_i(s)$ for $0 \leq i \leq \infty$ and $s \in [a, b]$ in Theorem 1. ■

Remark 2. If we set

$$P_i(s) = \frac{[s - (\lambda b + (1 - \lambda)a)]^i}{i!} \quad \text{and} \quad Q_i(s) = \frac{[s - (\mu b + (1 - \mu)a)]^i}{i!} \quad (43)$$

in identity (25), then

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= \sum_{k=0}^n \frac{(b-a)^k}{k!} [b(\mu-1)^k f^{(k)}(b) - a\lambda^k f^{(k)}(a)] \\ &\quad - \sum_{k=1}^n \sum_{i=1}^k \frac{(b-a)^i}{i!} [(\mu-1)^i f^{(i-1)}(b) - \lambda^i f^{(i-1)}(a)] \\ &\quad - t \sum_{k=0}^n \frac{(b-a)^k}{k!} [(\mu-1)^k f^{(k)}(b) - \lambda^k f^{(k)}(a)] \\ &\quad + \sum_{k=1}^n \sum_{i=1}^k \frac{(-1)^i}{i!} \{ [t - (\mu b + (1 - \mu)a)]^i - [t - (\lambda b + (1 - \lambda)a)]^i \} f^{(i-1)}(t) \\ &\quad + \frac{(-1)^n}{n!} \left(\int_a^t (t-s) [s - (\lambda b + (1 - \lambda)a)]^n f^{(n+1)}(s) ds \right. \\ &\quad \left. + \int_t^b (t-s) [s - (\mu b + (1 - \mu)a)]^n f^{(n+1)}(s) ds dx \right). \quad (44) \end{aligned}$$

If we set

$$P_i(s) = Q_i(s) = \frac{[s - (\lambda b + (1 - \lambda)a)]^i}{i!} \quad (45)$$

in formula (42), then

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= \sum_{k=0}^n \frac{(b-a)^k}{k!} [b(\lambda-1)^k f^{(k)}(b) - a\lambda^k f^{(k)}(a)] \\ &\quad - \sum_{k=1}^n \sum_{i=1}^k \frac{(b-a)^i}{i!} [(\lambda-1)^i f^{(i-1)}(b) - \lambda^i f^{(i-1)}(a)] \\ &\quad - t \sum_{k=0}^n \frac{(b-a)^k}{k!} [(\lambda-1)^k f^{(k)}(b) - \lambda^k f^{(k)}(a)] \\ &\quad + \frac{(-1)^n}{n!} \int_a^b (t-s) [s - (\lambda b + (1 - \lambda)a)]^n f^{(n+1)}(s) ds. \quad (46) \end{aligned}$$

Taking $\lambda = 0$ or $\lambda = 1$ in (46) yields expansions about b and a respectively. Namely,

$$\begin{aligned} (n+1) \int_a^b f(x) dx &= b \sum_{k=0}^n \frac{(a-b)^k}{k!} f^{(k)}(b) - \sum_{k=1}^n \sum_{i=1}^k \frac{(a-b)^i}{i!} f^{(i-1)}(b) \\ &\quad - t \sum_{k=0}^n \frac{(a-b)^k}{k!} f^{(k)}(b) + \frac{1}{n!} \int_a^b (t-s)(a-s)^n f^{(n+1)}(s) ds. \quad (47) \end{aligned}$$

or

$$(n+1) \int_a^b f(x) dx = \sum_{k=1}^n \sum_{i=1}^k \frac{(b-a)^i}{i!} f^{(i-1)}(a) - a \sum_{k=0}^n \frac{(b-a)^k}{k!} f^{(k)}(a) \\ + t \sum_{k=0}^n \frac{(b-a)^k}{k!} f^{(k)}(a) + \frac{1}{n!} \int_a^b (t-s)(b-s)^n f^{(n+1)}(s) ds. \quad (48)$$

Remark 3. If the assumption is made that,

$$P_i(s) = P_{i,B}(s; b; a) = \frac{(b-a)^i}{i!} B_i \left(\frac{s-a}{b-a} \right) \quad (49)$$

for nonnegative integer i , then from Theorem 2, straightforward calculations and utilising the properties (21), (22) together with the fact that $B_{2i+1} = 0$ for $i \in \mathbb{N}$, produces

$$(n+1) \int_a^b f(x) dx = [bf(b) - af(a)] - \frac{1}{2}(b-a)[af'(a) + bf'(b)] \\ + \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor} \frac{(b-a)^{2k}}{(2k)!} B_{2k} [bf^{(2k)}(b) - af^{(2k)}(a)] \\ + \frac{n}{2}(b-a)[f(a) + f(b)] - \sum_{k=2}^n \sum_{i=1}^{\lfloor \frac{k}{2} \rfloor} \frac{(b-a)^{2i}}{(2i)!} B_{2i} [f^{(2i-1)}(b) - f^{(2i-1)}(a)] \\ - t \left\{ [f(b) - f(a)] - \frac{1}{2}(b-a)[f'(a) + f'(b)] + \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor} \frac{(b-a)^{2k}}{(2k)!} B_{2k} [f^{(2k)}(b) - f^{(2k)}(a)] \right\} \\ + \frac{(a-b)^n}{n!} \int_a^b (t-s) B_n \left(\frac{s-a}{b-a} \right) f^{(n+1)}(s) ds, \quad (50)$$

where $[x]$ denotes the Gauss function, whose value is the largest integer not exceeding x .

Remark 4. Suppose that in Theorem 2

$$P_i(s) = P_{i,E}(s; a; b) = \frac{(a-b)^i}{i!} E_i \left(\frac{s-b}{a-b} \right) \quad (51)$$

for nonnegative integer i , then, by direct calculation and utilising (21), (22) and $B_{2i+1} = 0$ for $i \in \mathbb{N}$, we obtain

$$(n+1) \int_a^b f(x) dx = af(a) + bf(b) \\ + \sum_{k=1}^{\lfloor \frac{n+1}{2} \rfloor} \frac{2(1-4^k)(b-a)^{2k-1}}{(2k)!} B_{2k} [af^{(2k-1)}(a) + bf^{(2k-1)}(b)] \\ + \sum_{k=1}^n \sum_{i=1}^{\lfloor \frac{k+1}{2} \rfloor} \frac{2(4^i-1)(b-a)^{2i-1}}{(2i)!} B_{2i} [f^{(2(i-1))}(a) + f^{(2(i-1))}(b)]$$

$$\begin{aligned}
& -t \left\{ af(a) + bf(b) + \sum_{k=1}^{\lfloor \frac{n+1}{2} \rfloor} \frac{2(4^k - 1)(b-a)^{2k-1}}{(2k)!} B_{2k} [f^{(2k-1)}(a) + f^{(2k-1)}(b)] \right\} \\
& \quad + \frac{(b-a)^n}{n!} \int_a^b (t-s) E_n \left(\frac{s-b}{a-b} \right) f^{(n+1)}(s) ds, \quad (52)
\end{aligned}$$

where $[x]$ also denotes the Gauss function as in Remark 3.

3. SOME NEW IYENGAR TYPE INEQUALITIES

In the work that follows, we adopt the notation:

$$\begin{aligned}
\Gamma_P &= \sup_{x \in [a,b]} p(n, n+1; x) = \sup_{x \in [a,b]} \{(-1)^n P_n(x) f^{(n+1)}(x)\}, \\
\gamma_P &= \inf_{x \in [a,b]} p(n, n+1; x) = \inf_{x \in [a,b]} \{(-1)^n P_n(x) f^{(n+1)}(x)\}, \\
\Gamma_Q &= \sup_{x \in [a,b]} q(n, n+1; x) = \sup_{x \in [a,b]} \{(-1)^n Q_n(x) f^{(n+1)}(x)\}, \\
\gamma_Q &= \inf_{x \in [a,b]} q(n, n+1; x) = \inf_{x \in [a,b]} \{(-1)^n Q_n(x) f^{(n+1)}(x)\}, \\
\Gamma &= \sup_{x \in [a,b]} f^{(n+1)}(x), \quad \gamma = \inf_{x \in [a,b]} f^{(n+1)}(x),
\end{aligned} \quad (53)$$

where the definitions (26) and (27) have been used.

Suppose $f(x)$ is a n -times differentiable function on the closed interval $[a, b]$ and $f^{(n+1)}(x)$ exists on (a, b) . Let

$$M = \Gamma = \sup_{s \in (a,b)} f^{(n+1)}(s), \quad N = \gamma = \inf_{s \in (a,b)} f^{(n+1)}(s), \quad (54)$$

then, from (41), we can deduce Theorem B and Theorem C, the results obtained in [25] and [26].

Bounds may also be obtained in terms of the traditional Lebesgue norms for $f^{(n+1)} \in L^p([a, b])$, $1 \leq p < \infty$. Specifically, if $f^{(n)}(x)$ is absolutely continuous on the closed interval $[a, b]$, then, for any $t \in [a, b]$, it follows from (41), that

$$\begin{aligned}
& \left| \int_a^b f(x) dx - \sum_{i=0}^n \frac{f^{(i)}(a)}{(i+1)!} (t-a)^{i+1} + \sum_{i=0}^n \frac{f^{(i)}(b)}{(i+1)!} (t-b)^{i+1} \right| \\
& \leq \frac{(t-a)^{n+2} + (b-t)^{n+2}}{(n+2)!} \int_a^b |f^{(n+1)}(s)| ds. \quad (55)
\end{aligned}$$

Also, as a simple consequence of Hölder inequality, if $f^{(n+1)} \in L^p([a, b])$ for a positive number $p > 1$, then, for any $t \in [a, b]$, we have

$$\begin{aligned}
& \left| \int_a^b f(x) dx - \sum_{i=0}^n \frac{f^{(i)}(a)}{(i+1)!} (t-a)^{i+1} + \sum_{i=0}^n \frac{f^{(i)}(b)}{(i+1)!} (t-b)^{i+1} \right| \\
& \leq \frac{(t-a)^{n+1+\frac{1}{q}} + (b-t)^{n+1+\frac{1}{q}}}{(n+1)! \sqrt[q]{nq+q+1}} \left(\int_a^b |f^{(n+1)}(s)|^p ds \right)^{\frac{1}{p}}, \quad (56)
\end{aligned}$$

where q satisfies $\frac{1}{p} + \frac{1}{q} = 1$.

Note that inequality (56) has been proved in [12].

Let $\{P_i(x)\}_{i=0}^{\infty}$ and $\{Q_i(x)\}_{i=0}^{\infty}$ be two harmonic sequences of polynomials. If $f : [a, b] \rightarrow \mathbb{R}$ is a function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, then, from identity (25), it is easy to obtain the expression

$$\begin{aligned}
& \frac{1}{2} [(a-t)^2 \gamma_Q - (b-t)^2 \Gamma_P] \\
& \leq (n+1) \int_a^b f(x) dx + a \sum_{k=0}^n p(k, k; a) - b \sum_{k=0}^n q(k, k; b) \\
& \quad - \sum_{k=1}^n \sum_{i=1}^k [p(i, i-1, a) - q(i, i-1; b)] \\
& \quad - t \sum_{k=0}^n [p(k, k; a) - q(k, k; b)] - \sum_{k=1}^n \sum_{i=1}^k [q(i, i-1, t) - p(i, i-1; t)] \\
& \leq \frac{1}{2} [(a-t)^2 \Gamma_P - (b-t)^2 \gamma_Q]. \quad (57)
\end{aligned}$$

Let $\{P_i(x)\}_{i=0}^{\infty}$ be a harmonic sequence of polynomials. If $f : [a, b] \rightarrow \mathbb{R}$ is a function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, then we have the following inequalities

$$\begin{aligned}
& \frac{1}{2} [a^2 \gamma_P - b^2 \Gamma_P] - \frac{\{b\Gamma_P - a\gamma_P - \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{2(\gamma_P - \Gamma_P)} \\
& \leq (n+1) \int_a^b f(x) dx - b \sum_{k=0}^n p(k, k; b) + a \sum_{k=0}^n p(k, k; a) \\
& \quad + \sum_{k=1}^n \sum_{i=1}^k [p(i, i-1; b) - p(i, i-1; a)] \\
& \leq \frac{1}{2} [a^2 \Gamma_P - b^2 \gamma_P] - \frac{\{b\gamma_P - a\Gamma_P - \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{2(\Gamma_P - \gamma_P)}. \quad (58)
\end{aligned}$$

In fact, (58) follows from the identity (42) and maximizing or minimizing a quadratic polynomial in t .

Let $\{P_i(x)\}_{i=0}^{\infty}$ be a harmonic sequence of polynomials. If $f : [a, b] \rightarrow \mathbb{R}$ is a function such that $f^{(n)}(x)$ is absolutely continuous for some $n \in \mathbb{N}$, and defining

$$\mathcal{U} := \max_{s \in [a, b]} |P_n(s)|, \quad \mathcal{V} := \max_{s \in [a, b]} |f^{(n+1)}(s)|, \quad \text{and} \quad \mathcal{W} := \max_{s \in [a, b]} |P_n(s) f^{(n+1)}(s)|,$$

then, from (42), maximizing or minimizing a quadratic polynomial in t yields

$$\begin{aligned}
& \frac{\{(a+b)\mathcal{UV} - \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{4\mathcal{UV}} - \frac{a^2 + b^2}{2}\mathcal{UV} \\
& \leq \frac{\{(a+b)\mathcal{W} - \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{4\mathcal{W}} - \frac{a^2 + b^2}{2}\mathcal{W} \\
& \leq (n+1) \int_a^b f(x) dx - b \sum_{k=0}^n p(k, k; b) + a \sum_{k=0}^n p(k, k; a) \\
& \quad + \sum_{k=1}^n \sum_{i=1}^k [p(i, i-1; b) - p(i, i-1; a)] \\
& \leq \frac{a^2 + b^2}{2}\mathcal{W} - \frac{\{(a+b)\mathcal{W} + \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{4\mathcal{W}} \\
& \leq \frac{a^2 + b^2}{2}\mathcal{UV} - \frac{\{(a+b)\mathcal{UV} + \sum_{k=0}^n [p(k, k; b) - p(k, k; a)]\}^2}{4\mathcal{UV}}. \quad (59)
\end{aligned}$$

Remark 5. Now it is clear that, if estimating the following general integral remainders

$$\int_a^t (t-s)p(n, n+1; s) ds + \int_t^b (t-s)q(n, n+1; s) ds, \quad (60)$$

$$\text{and } \int_a^b (t-s)p(n, n+1; s) ds \quad (61)$$

that appear in (25) and (42), then using better and more appropriate techniques than previously gives rise to more accurate Iyengar type inequalities.

Acknowledgements. This paper was completed during the first author's visit to the RGMIA between November 1, 2001 and January 31, 2002, as a Visiting Professor with grants from the Victoria University of Technology and Jiaozuo Institute of Technology.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series 55, 4th printing, Washington, 1965.
- [2] R. P. Agarwal and S. S. Dragomir, *An application of Hayashi inequality for differentiable functions*, *Comput. Math. Appl.* **32** (1996), no. 6, 95–99.
- [3] P. Cerone, *Generalised trapezoidal rules with error involving bounds of the n -th derivative*, *Math. Ineq. & Appl.*, (accepted).
- [4] P. Cerone, *Generalised Taylor's formula with estimates of the remainder*, *Inequality Theory and Applications*, 2 (2002), Y.J. Cho, J.K. Kim and S.S. Dragomir (Ed.), Nova Science Publishers, (accepted).
- [5] P. Cerone and S. S. Dragomir, *Lobatto type quadrature rules for functions with bounded derivative*, *Math. Inequal. Appl.* **3** (2000), no. 2, 197–209. RGMIA Res. Rep. Coll. **2** (1999), no. 2, Art. 1, 133–146. Available online at <http://rgmia.vu.edu.au/v2n2.html>.
- [6] P. Cerone and S. S. Dragomir, *On a weighted generalization of Iyengar type inequalities involving the bounded first derivative*, *Math. Inequal. Appl.* **3** (2000), no. 1, 35–44. RGMIA Res. Rep. Coll. **2** (1999), no. 2, Art. 2, 147–157. Available online at <http://rgmia.vu.edu.au/v2n2.html>.
- [7] P. Cerone and S. S. Dragomir, *New bounds for a perturbed generalized Taylor's formula*, *East Asian Math. J.* **17** (2001), no. 2, 197–215.

- [8] L.-H. Cui and B.-N. Guo, *On proofs of an integral inequality and its generalizations*, J. Zhengzhou Grain College, **17** (1996), Supplement, 152–154 and 158. (Chinese)
- [9] S. S. Dragomir, *Better bounds in some Ostrowski-Grüss type inequalities*, Tamkang J. Math. **32** (2001), no. 3, 211–216.
- [10] S. S. Dragomir and S. Wang, *Applications of Iyengar type inequalities to the estimation of error bounds for the trapezoidal quadrature rule*, Tamkang J. Math. **29** (1998), no. 1, 55–58.
- [11] A. M. Fink, *Bounds on the deviation of a function from its averages*, Czechoslovak Math. J. **42** (1992), no. 117, 289–310.
- [12] B.-N. Guo and F. Qi, *Estimates for an integral in L^p norm of the $(n + 1)$ -th derivative of its integrand*, RGMIA Res. Rep. Coll. **3** (2000), no. 3, Art. 2. Available online at <http://rgmia.vu.edu.au/v3n3.html>.
- [13] B.-N. Guo and F. Qi, *Generalisation of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. (2001), in the press.
- [14] B.-N. Guo and F. Qi, *Proofs of an integral inequality*, Mathematics and Informatics Quarterly **7** (1997), no. 4, 182–184.
- [15] K. S. K. Iyengar, *Note on an inequality*, Math. Student **6** (1938), 75–76.
- [16] J.-Ch. Kuang, *Chángyòng Bùdēngshì (Applied Inequalities)*, 2nd edition, Hunan Education Press, Changsha City, China, 1993. (Chinese)
- [17] Q.-M. Luo, B.-N. Guo, and F. Qi, *Generalizations of Bernoulli numbers and polynomials*, submitted.
- [18] M. Matić, J. Pečarić and N. Ujević, *On new estimation of the remainder in generalized Taylor formula*, Math. Inequal. Appl. **2** (1999), no. 3, 343–361.
- [19] G. V. Milovanović and J. E. Pečarić, *Some considerations on Iyengar inequality and some related applications*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No. **544–576** (1976), 166–170.
- [20] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993.
- [21] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, 1991.
- [22] C. E. M. Pearce, J. E. Pečarić, N. Ujević and S. Varonsaneč, *Generalisation of some inequalities of Ostrowski-Grüss type*, Math. Ineq. & Appl. **3** (2000), 25–34.
- [23] C. E. M. Pearce and J. E. Pečarić, *On Anastassiou's generalisation of the Ostrowski's inequality and related results*, J. Comp. Anal. & Appl. **2** (2000), 215–235.
- [24] F. Qi, *Inequalities for an integral*, Math. Gaz. **80** (1996), no. 488, 376–377.
- [25] F. Qi, *Further generalizations of inequalities for an integral*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. **8** (1997), 79–83.
- [26] F. Qi, *Inequalities for a multiple integral*, Acta Math. Hungar. **84** (1999), no. 1-2, 19–26.
- [27] F. Qi and B.-N. Guo, *Generalized Bernoulli polynomials*, RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 10. Available online at <http://rgmia.vu.edu.au/v4n4.html>.
- [28] F. Qi and Y.-J. Zhang, *Inequalities for a weighted integral*, Adv. Stud. Contemp. Math. **4** (2002), no. 2, in press. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 2, 967–975. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [29] F. Qi, *Inequalities for a weighted multiple integral*, J. Math. Anal. Appl. **253** (2001), no. 2, 381–388. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 4, 991–997. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [30] P. M. Vasić and G. V. Milonović, *On an inequality of Iyengar*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No. **544–576** (1976), 18–24.

(Qi) DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY,
HENAN 454000, CHINA

E-mail address: `qifeng@jzit.edu.cn`, `qifeng618@hotmail.com`

URL: `http://rgmia.vu.edu.au/qi.html`

(Cerone) SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECH-
NOLOGY, P. O. BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA

E-mail address: `pc@matilda.vu.edu.au`

URL: `http://rgmia.vu.edu.au/cerone`

(Dragomir) SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECH-
NOLOGY, P. O. BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA

E-mail address: `sever.dragomir@vu.edu.au`

URL: `http://rgmia.vu.edu.au/SSDragomirWeb.html`