

**NEW INEQUALITIES FOR THE
ČEBYŠEV FUNCTIONAL INVOLVING
TWO n -TUPLES OF REAL NUMBERS
AND APPLICATIONS**

P. CERONE AND S.S. DRAGOMIR

ABSTRACT. New upper and lower bounds for the unweighted Čebyšev functional involving two n -tuples of real numbers are developed and applications for guessing mappings are given.

1. INTRODUCTION

For two n -tuples of real numbers, consider the Čebyšev's functional

$$(1.1) \quad D_n(\bar{x}, \bar{y}) := \frac{1}{n} \sum_{i=1}^n x_i y_i - \frac{1}{n} \sum_{i=1}^n x_i \cdot \frac{1}{n} \sum_{i=1}^n y_i,$$

where $\bar{x} := (x_1, \dots, x_n)$, $\bar{y} := (y_1, \dots, y_n) \in \mathbb{R}^n$.

If \bar{x}, \bar{y} are synchronous (asynchronous), such that

$$(1.2) \quad (x_i - x_j)(y_i - y_j) \geq (\leq) 0$$

for each $i, j \in \{1, \dots, n\}$

then the well known Čebyšev's inequality

$$(1.3) \quad D_n(\bar{x}, \bar{y}) \geq (\leq) 0$$

holds.

In [9], the following refinement of Čebyšev's inequality (1.3) has been obtained. Namely,

$$(1.4) \quad D_n(\bar{x}, \bar{y}) \geq \max \{ |D_n(|\bar{x}|, \bar{y})|, |D_n(\bar{x}, |\bar{y}|)|, |D_n(|\bar{x}|, |\bar{y}|)| \},$$

provided \bar{x}, \bar{y} are synchronous and

$$|\bar{x}| := (|x_1|, \dots, |x_n|).$$

If $x \leq x_i \leq X$, $y \leq y_i \leq Y$ for each $i \in \{1, \dots, n\}$, then the magnitude of the difference $D_n(\bar{x}, \bar{y})$ may be evaluated by the use of Biernacki, Pidek and Ryll-Nardzewski's inequality [1]

$$(1.5) \quad |D_n(\bar{x}, \bar{y})| \leq \frac{1}{n} \left[\frac{n}{2} \right] \left(1 - \frac{1}{n} \left[\frac{n}{2} \right] \right) (X - x)(Y - y)$$

Date: April 12, 2002.

1991 *Mathematics Subject Classification*. Primary 26D15, 26D10; Secondary 94A05.

Key words and phrases. Čebyšev inequality, Guessing mapping.

$$= \frac{1}{n^2} \left[\frac{n^2}{4} \right] (X - x)(Y - y) \leq \frac{1}{4} (X - x)(Y - y).$$

The following results similar to that in (1.5) are also known

$$(1.6) \quad |D_n(\bar{x}, \bar{y})| \leq \begin{cases} \frac{n^2-1}{12} \cdot \max_{j=1, n-1} |\Delta x_j| \max_{j=1, n-1} |\Delta y_j|, [3] \\ \frac{1}{2} \left(1 - \frac{1}{n} \right) \sum_{i=1}^{n-1} |\Delta x_i| \sum_{i=1}^{n-1} |\Delta y_i|, [5] \\ \frac{n^2-1}{6n} \left(\sum_{j=1}^{n-1} |\Delta x_j|^p \right)^{\frac{1}{p}} \left(\sum_{j=1}^{n-1} |\Delta y_j|^q \right)^{\frac{1}{q}}, \\ \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1; [4]. \end{cases}$$

The constants $\frac{1}{12}, \frac{1}{2}$ and $\frac{1}{6}$ in (1.6) respectively are sharp in the sense that they can not be replaced by smaller constants.

The main aim of this paper is both to point out other sufficient conditions for the positivity of the Čebyšev functional and to determine upper bounds for the magnitude of $D_n(\cdot, \cdot)$. Some applications for the moments of guessing mapping are also mentioned.

2. SOME UPPER BOUNDS

In dealing with the magnitude of the difference, $D_n(\bar{x}, \bar{y})$ as defined in (1.1), a natural approach is embodied in the following theorem.

Theorem 1. *Let \bar{c} be the constant n -tuple with all of its elements equal to $c \in \mathbb{R}$. For any two n -tuples $\bar{x} := (x_1, \dots, x_n)$, $\bar{y} := (y_1, \dots, y_n)$ of real numbers one has the inequalities*

$$(2.1) \quad 0 \leq |D_n(\bar{x}, \bar{y})| \leq \begin{cases} \frac{1}{n} \|\bar{y} - \bar{y}_M\|_1 \inf_{c \in \mathbb{R}} \|\bar{x} - \bar{c}\|_\infty; \\ \frac{1}{n} \|\bar{y} - \bar{y}_M\|_q \inf_{c \in \mathbb{R}} \|\bar{x} - \bar{c}\|_p, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \|\bar{y} - \bar{y}_M\|_\infty \inf_{c \in \mathbb{R}} \|\bar{x} - \bar{c}\|_1; \end{cases}$$

$$\leq \begin{cases} \frac{1}{n} \|\bar{y} - \bar{y}_M\|_1 \min \{ \|\bar{x}\|_\infty, \|\bar{x} - \bar{x}_M\|_\infty \}; \\ \frac{1}{n} \|\bar{y} - \bar{y}_M\|_q \min \left\{ \|\bar{x}\|_p, \|\bar{x} - \bar{x}_M\|_p \right\}, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \|\bar{y} - \bar{y}_M\|_\infty \min \{ \|\bar{x}\|_1, \|\bar{x} - \bar{x}_M\|_1 \}; \end{cases}$$

where

$$x_M := \frac{1}{n} \sum_{k=1}^n x_k, \quad y_M := \frac{1}{n} \sum_{k=1}^n y_k,$$

and $\bar{\mathbf{x}}_M, \bar{\mathbf{y}}_M$ the vectors with all components equal to x_M, y_M . Here, $\|\cdot\|_p$ ($p \in [1, \infty]$) are the usual p -norms on \mathbb{R}^n , namely

$$\|\bar{\mathbf{x}}\|_\infty := \max_{i=1, \dots, n} |x_i|,$$

$$\|\bar{\mathbf{x}}\|_p := \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}, \quad 1 \leq p < \infty.$$

Proof. Firstly, let us observe that for any $c \in \mathbb{R}$ one has the identity

$$(2.2) \quad D_n(\bar{\mathbf{x}}, \bar{\mathbf{y}}) = D_n(\bar{\mathbf{x}} - \bar{\mathbf{c}}, \bar{\mathbf{y}} - \bar{\mathbf{y}}_M)$$

$$= \frac{1}{n} \sum_{i=1}^n (x_i - c)(y_i - y_M).$$

Taking the modulus and using Hölder's inequality, we have

$$(2.3) \quad |D_n(\bar{\mathbf{x}}, \bar{\mathbf{y}})|$$

$$\leq \frac{1}{n} \sum_{i=1}^n |x_i - c| |y_i - y_M|$$

$$\leq \begin{cases} \frac{1}{n} \max_{i=1, \dots, n} |x_i - c| \sum_{i=1}^n |y_i - y_M|; \\ \frac{1}{n} \left(\sum_{i=1}^n |x_i - c|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |y_i - y_M|^q \right)^{\frac{1}{q}}, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \sum_{i=1}^n |x_i - c| \max_{i=1, \dots, n} |y_i - y_M|; \end{cases}$$

$$= \begin{cases} \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_\infty \|\bar{\mathbf{y}} - \bar{\mathbf{y}}_M\|_1; \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_p \|\bar{\mathbf{y}} - \bar{\mathbf{y}}_M\|_q, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_1 \|\bar{\mathbf{y}} - \bar{\mathbf{y}}_M\|_\infty. \end{cases}$$

Taking the inf over $c \in \mathbb{R}$ in (2.3), we deduce the second inequality in (2.1).

Since

$$\inf_{c \in \mathbb{R}} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_p \leq \begin{cases} \|\bar{\mathbf{x}}\|_p \\ \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_p \end{cases} \quad \text{for any } p \in [1, \infty],$$

the final part of (2.1) is also proved. ■

Corollary 1. For any $\bar{\mathbf{x}}$ an n -tuple of real numbers one has

$$(2.4) \quad 0 \leq D_n(\bar{\mathbf{x}}, \bar{\mathbf{x}}) = \frac{1}{n} \sum_{i=1}^n x_i^2 - \left(\frac{1}{n} \sum_{i=1}^n x_i \right)^2$$

$$\leq \begin{cases} \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_1 \inf_{c \in \mathbb{R}} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_\infty, \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_q \inf_{c \in \mathbb{R}} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_p, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_\infty \inf_{c \in \mathbb{R}} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_1 \end{cases}$$

$$\leq \begin{cases} \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_1 \min \{ \|\bar{\mathbf{x}}\|_\infty, \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_\infty \}, \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_q \min \left\{ \|\bar{\mathbf{x}}\|_p, \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_p \right\}, \\ \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{n} \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_\infty \min \{ \|\bar{\mathbf{x}}\|_1, \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_1 \}, \end{cases}$$

Remark 1. For $p = q = 2$, we know that

$$(2.5) \quad \inf_{c \in \mathbb{R}} \|\bar{\mathbf{x}} - \bar{\mathbf{c}}\|_2 = \|\bar{\mathbf{x}} - \bar{\mathbf{x}}_M\|_2$$

$$= \frac{1}{n} \left[n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 \right]^{\frac{1}{2}}$$

$$= D_n^{\frac{1}{2}}(\bar{\mathbf{x}}, \bar{\mathbf{x}})$$

to produce the known inequality:

$$(2.6) \quad [D_n(\bar{\mathbf{x}}, \bar{\mathbf{y}})]^2 \leq D_n(\bar{\mathbf{x}}, \bar{\mathbf{x}}) D_n(\bar{\mathbf{y}}, \bar{\mathbf{y}}).$$

3. SOME POSITIVITY RESULTS

To study the positivity of $D_n(\cdot, \cdot)$, we introduce the following class of real numbers associated with two given n -tuples $\bar{\mathbf{x}} = (x_1, \dots, x_n)$, and $\bar{\mathbf{y}} = (y_1, \dots, y_n)$, namely,

$$(3.1) \quad \mathfrak{C}_n(\bar{\mathbf{x}}, \bar{\mathbf{y}}) := \{c \in \mathbb{R} \mid (x_i - c)(y_i - y_M) \geq 0$$

$$\text{for each } i \in \{1, \dots, n\}\}.$$

For $n = 2$ and if we assume that $y_1 < y_2$, then the condition $(x_i - c)(y_i - y_M) \geq 0$, $i \in \{1, 2\}$ is equivalent to

$$\begin{cases} (x_1 - c)(y_1 - y_2) \geq 0 \\ (x_2 - c)(y_2 - y_1) \geq 0 \end{cases}$$

or to $x_1 \leq c \leq x_2$.

So, $\mathfrak{C}_2(\bar{\mathbf{x}}, \bar{\mathbf{y}})$ is nonempty iff $x_1 \leq x_2$.

We will say that the n -tuples $(\bar{\mathbf{x}}, \bar{\mathbf{y}})$ (in this particular order) are *positively correlated*, if $\mathfrak{C}_n(\bar{\mathbf{x}}, \bar{\mathbf{y}})$ is nonempty.

For instance, for any $\bar{\mathbf{x}} \in \mathbb{R}^n$ we have $(\bar{\mathbf{x}}, \bar{\mathbf{x}})$ are positively correlated as $c = x_M \in \mathfrak{C}_n(\bar{\mathbf{x}}, \bar{\mathbf{x}})$.

The following result providing a refinement of Čebyšev's inequality holds.

Theorem 2. Assume that the n -tuples $(\bar{\mathbf{x}}, \bar{\mathbf{y}})$ are positively correlated. Then one has the inequality:

$$(3.2) \quad D_n(\bar{\mathbf{x}}, \bar{\mathbf{y}}) \geq \max \left\{ |A_n|, \sup_{c \in \mathfrak{C}_n(\bar{\mathbf{x}}, \bar{\mathbf{y}})} |B_n(c)|, \right. \\ \left. \sup_{c \in \mathfrak{C}_n(\bar{\mathbf{x}}, \bar{\mathbf{y}})} |C_n(c)| \right\} \geq 0,$$

where

$$A_n := \frac{1}{n} \sum_{i=1}^n |x_i| y_i - \frac{1}{n} \sum_{i=1}^n |x_i| \cdot \frac{1}{n} \sum_{i=1}^n y_i$$

$$B_n(c) := \frac{1}{n} \sum_{i=1}^n |x_i y_i| - |c| \cdot \frac{1}{n} \sum_{i=1}^n |y_i| \\ - \frac{1}{n} \sum_{i=1}^n |x_i| \left| \frac{1}{n} \sum_{i=1}^n y_i \right| + |c| \cdot \left| \frac{1}{n} \sum_{i=1}^n y_i \right|$$

and

$$C_n(c) := \frac{1}{n} \sum_{i=1}^n x_i |y_i| - c \cdot \frac{1}{n} \sum_{i=1}^n |y_i| \\ - \frac{1}{n} \sum_{i=1}^n x_i \left| \frac{1}{n} \sum_{i=1}^n y_i \right| + c \left| \frac{1}{n} \sum_{i=1}^n y_i \right|.$$

Proof. Let $c \in \mathfrak{C}_n(\bar{x}, \bar{y})$, then

$$(3.3) \quad (x_i - c)(y_i - y_M) \\ = |(x_i - c)(y_i - y_M)| \\ \geq \begin{cases} |(|x_i| - |c|)(y_i - y_M)| \\ |(|x_i| - |c|)(|y_i| - |y_M|)| \\ |(x_i - c)(|y_i| - |y_M|)| \end{cases}$$

for each $i \in \{1, \dots, n\}$.

Summing over i from 1 to n in (3.3) and using the generalised triangle inequality, we get

$$(3.4) \quad \frac{1}{n} \sum_{i=1}^n (x_i - c)(y_i - y_M) \\ \geq \frac{1}{n} \begin{cases} \left| \sum_{i=1}^n (|x_i| - |c|)(y_i - y_M) \right|, \\ \left| \sum_{i=1}^n (|x_i| - |c|)(|y_i| - |y_M|) \right|, \\ \left| \sum_{i=1}^n (x_i - c)(|y_i| - |y_M|) \right|. \end{cases}$$

Since

$$\sum_{i=1}^n (|x_i| - |c|)(y_i - y_M) \\ = \sum_{i=1}^n |x_i| y_i - |c| \sum_{i=1}^n y_i - \sum_{i=1}^n |x_i| \cdot y_M + n|c| y_M \\ = \sum_{i=1}^n |x_i| y_i - \sum_{i=1}^n |x_i| \cdot y_M = nA_n,$$

$$\sum_{i=1}^n (|x_i| - |c|)(|y_i| - |y_M|) \\ = \sum_{i=1}^n |x_i y_i| - |c| \sum_{i=1}^n |y_i| - |y_M| \sum_{i=1}^n |x_i| + n|c| |y_M| \\ = nB_n(c)$$

and

$$\sum_{i=1}^n (x_i - c)(|y_i| - |y_M|) \\ = \sum_{i=1}^n x_i |y_i| - c \sum_{i=1}^n |y_i| - |y_M| \sum_{i=1}^n x_i + n c |y_M| \\ = nC_n(c),$$

then by the identity (2.2) and the inequality (3.4) we deduce

$$(3.5) \quad D_n(\bar{x}, \bar{y}) \geq \begin{cases} |A_n| \\ |B_n(c)| \\ |C_n(c)| \end{cases} \quad \text{for any } c \in \mathfrak{C}_n(\bar{x}, \bar{y}).$$

Taking the sup in (3.5) for $c \in \mathfrak{C}_n(\bar{x}, \bar{y})$, produces (3.2). ■

The following corollaries are natural.

Corollary 2. Assume that \bar{x}, \bar{y} are such that

$$(3.6) \quad x_i (y_i - y_M) \geq 0 \quad \text{for each } i \in \{1, \dots, n\}.$$

Then, one has the inequality

$$(3.7) \quad D_n(\bar{x}, \bar{y}) \geq \max \left\{ |A_n|, |B_n^{(1)}|, |C_n^{(1)}| \right\} \geq 0,$$

where A_n was defined in Theorem 2 and

$$B_n^{(1)} := \frac{1}{n} \sum_{i=1}^n |x_i y_i| - \frac{1}{n} \sum_{i=1}^n |x_i| \left| \frac{1}{n} \sum_{i=1}^n y_i \right|, \\ C_n^{(1)} := \frac{1}{n} \sum_{i=1}^n x_i |y_i| - \frac{1}{n} \sum_{i=1}^n x_i \left| \frac{1}{n} \sum_{i=1}^n y_i \right|.$$

Corollary 3. Assume that \bar{x}, \bar{y} are such that

$$(3.8) \quad (x_i - x_M)(y_i - y_M) \geq 0 \\ \text{for each } i \in \{1, \dots, n\}.$$

Then, one has the inequality

$$(3.9) \quad D_n(\bar{x}, \bar{y}) \geq \max \left\{ |A_n|, |B_n^{(2)}|, |C_n^{(2)}| \right\} \geq 0,$$

where A_n is as in Theorem 2 and

$$B_n^{(2)} := \frac{1}{n} \sum_{i=1}^n |x_i y_i| - \left| \frac{1}{n} \sum_{i=1}^n x_i \right| \cdot \frac{1}{n} \sum_{i=1}^n y_i \\ - \left| \frac{1}{n} \sum_{i=1}^n y_i \right| \left| \frac{1}{n} \sum_{i=1}^n |x_i| \right| + \left| \frac{1}{n} \sum_{i=1}^n x_i \right| \left| \frac{1}{n} \sum_{i=1}^n y_i \right|,$$

$$C_n^{(2)} := \frac{1}{n} \sum_{i=1}^n x_i |y_i| - \frac{1}{n} \sum_{i=1}^n x_i \cdot \frac{1}{n} \sum_{i=1}^n |y_i|.$$

Remark 2. We shall show now that there are positively correlated sequences (\bar{x}, \bar{y}) that are not synchronous.

Let $a < b$ and consider $y_1 = a$, $y_2 = \frac{a+b}{2}$, $y_3 = b$. The sequence (x_1, x_2, x_3) is positively correlated with (y_1, y_2, y_3) iff there is a $c \in \mathbb{R}$ such that

$$(x_i - c) \left(y_i - \frac{y_1 + y_2 + y_3}{3} \right) \geq 0$$

which is equivalent to:

$$(3.10) \quad (x_i - c) \left(y_i - \frac{a+b}{2} \right) \geq 0, \quad i = 1, 2, 3.$$

The assumption (3.10) is equivalent to:

$$\begin{cases} (x_1 - c)(a - b) \geq 0 \\ (x_2 - c) \left(\frac{a+b}{2} - \frac{a+b}{2} \right) \geq 0 \\ (x_3 - c)(a - b) \geq 0 \end{cases}$$

so that

$$\begin{cases} x_1 \leq c, \\ x_2 \in \mathbb{R}, \quad c \in \mathbb{R}. \\ c \leq x_3, \end{cases}$$

So, if we assume that $x_1 < x_3$ and $x_2 \in \mathbb{R}$, then (x_1, x_2, x_3) is positively correlated to (y_1, y_2, y_3) .

If we choose $x_2 < x_1$, then $(y_2 - y_1)(x_2 - x_1) < 0$ while $(y_3 - y_1)(x_3 - x_1) \geq 0$ showing that (x_1, x_2, x_3) and (y_1, y_2, y_3) are not synchronous.

Remark 3. It remains an open question if there are synchronous sequences that are not positively correlated.

4. SOME APPLICATIONS FOR MOMENTS OF GUESSING MAPPINGS

In 1994, J.L. Massey [13] considered the problem of guessing the value taken on by a discrete random variable X in one trial of a random experiment by asking questions of the form ‘‘Did X take on its i^{th} possible value?’’ until the answer is in the affirmative.

This problem arises for instance when a cryptologist must try different possible secret keys one at a time *after* minimising the possibilities by some cryptanalysis.

Consider a random variable X with finite range $X = \{x_1, \dots, x_n\}$ and distribution $P_X(x_k) = p_k$ for $k = 1, 2, \dots, n$.

A one-to-one function $G : \chi \rightarrow \{1, \dots, n\}$ is a guessing function for X . Thus

$$(4.1) \quad E(G^m) := \sum_{k=1}^n k^m p_k$$

is the m^{th} moment of this function, provided we renumber the x_i such that x_k is always the k^{th} guess.

In [13], Massey observed that, $E(G)$, the average number of guesses, is minimised by a guessing strategy that guesses the possible values of X in decreasing order of probability.

In the same paper [13], Massey proved that for an optimal guessing strategy

$$(4.2) \quad E(G) \geq \frac{1}{4} 2^{H(X)} + 1$$

provided $H(X) \geq 2$ bits,

where $H(X)$ is the Shannon entropy

$$(4.3) \quad H(X) = - \sum_{i=1}^n p_i \log_2(p_i).$$

He also showed that $E(G)$ may be arbitrarily large when $H(X)$ is an arbitrarily small positive number so that there is no interesting upper bound on $E(G)$ in terms of $H(X)$.

In 1996, Arikan [14] proved that any guessing algorithm for X obeys the lower bound

$$(4.4) \quad E(G^\rho) \geq \frac{\left[\sum_{k=1}^n p_k^{\frac{1}{1+\rho}} \right]^{1+\rho}}{[1 + \ln n]^\rho}, \quad \rho \geq 0$$

where as an optimal guessing algorithm for X satisfies

$$(4.5) \quad E(G^\rho) \leq \left[\sum_{k=1}^n p_k^{\frac{1}{1+\rho}} \right]^{1+\rho}, \quad \rho \geq 0.$$

In 1997, Boztaş [15] proved that for $m \geq 1$, and integer

$$(4.6) \quad E(G^m) \leq \frac{1}{m+1} \left[\sum_{k=1}^n p_k^{\frac{1}{1+m}} \right]^{1+m} + \frac{1}{m+1} \left\{ \binom{m+1}{2} E(G^{m-1}) - \binom{m+1}{3} E(G^{m-2}) + \dots + (-1)^{m+1} \right\}$$

provided the guessing strategy satisfies the relation:

$$(4.7) \quad p_{k+1}^{\frac{1}{1+m}} \leq \frac{1}{k} \left(p_1^{\frac{1}{1+m}} + \dots + p_k^{\frac{1}{1+m}} \right),$$

$k = 1, \dots, n-1.$

In 1997, Dragomir and Boztaş [16] obtained, for any guessing sequence, the following bounds for the

expectation:

$$(4.8) \quad \left| E(G) - \frac{n+1}{2} \right| \leq \frac{(n-1)(n+1)}{6} \max_{1 \leq i < j \leq n} |p_i - p_j|,$$

$$(4.9) \quad \left| E(G) - \frac{n+1}{2} \right| \leq \sqrt{\frac{(n-1)(n+1)(n\|p\|_2^2 - 1)}{12}},$$

where $\|p\|_2^2 = \sum_{i=1}^n p_i^2$ and

$$(4.10) \quad \left| E(G) - \frac{n+1}{2} \right| \leq \left[\frac{n+1}{2} \right] \left(n - \left[\frac{n+1}{2} \right] \right) \max_{1 \leq k \leq n} \left| p_k - \frac{1}{n} \right|,$$

with $[x]$ representing the integer part of x .

For other results on $E(G^p)$, $p > 0$ see also [17]. We highlight only the following result which uses the Grüss inequality, giving for $p, q > 0$ that

$$(4.11) \quad |E(G^{p+q}) - E(G^p)E(G^q)| \leq \frac{1}{4}(n^q - 1)(n^p - 1).$$

The result (4.11) may be complemented in the following way (see for example [10]).

Theorem 3. *With the above assumptions, we have the inequality*

$$(4.12) \quad \left| E(G^{p+q}) - \frac{1+n^q}{2}E(G^p) - \frac{1+n^p}{2}E(G^q) + \frac{1+n^q}{2} \cdot \frac{1+n^p}{2} \right| \leq \frac{1}{4}(n^q - 1)(n^p - 1).$$

for any $p, q > 0$.

Applications for different particular instances of $p, q > 0$ may be provided, but we omit the details.

To obtain other inequalities for the moments of guessing mappings, we use the following Čebyšev type inequality

$$(4.13) \quad D_n(\bar{x}, \bar{y}) \geq (\leq) 0$$

provided

$$(x_i - x_M)(y_i - y_M) \geq (\leq) 0 \quad \text{for each } i \in \{1, \dots, n\}.$$

with a subscript M denoting the arithmetic mean.

The following result holds.

Theorem 4. *Assume that $S_n(p)$, $p > 0$ denotes the sum of p^{th} -power of the first n natural numbers, that is*

$$S_n(p) := \sum_{k=1}^n k^p.$$

If

$$p_i \begin{cases} \leq (\geq) \frac{1}{n}, & \text{for } i \leq \left[\frac{S_n(p)}{n} \right]^{1/p} \\ \geq (\leq) \frac{1}{n}, & \text{otherwise} \end{cases}$$

where $[x]$ represents the integer part of x , then we have the inequality

$$E(G^p) \geq (\leq) \frac{1}{n} S_n(p).$$

The proof follows by the inequality (4.13) on choosing $x_i = p_i$ and $y_i = i^p$, but we omit the details.

For particular values of p , one may produce some interesting particular inequalities.

If $p = 1$, then we have the inequality

$$E(G) \geq (\leq) \frac{n+1}{2}$$

provided

$$p_i \begin{cases} \leq (\geq) \frac{1}{n}, & i \leq \left[\frac{n+1}{2} \right] \\ \geq (\leq) \frac{1}{n}, & \text{otherwise} \end{cases}.$$

For $p = 2$, then

$$E(G) \geq (\leq) \frac{1}{6}(n+1)(2n+1)$$

provided

$$p_i \begin{cases} \leq (\geq) \frac{1}{n}, & i \leq \left[\frac{1}{6}(n+1)(2n+1) \right]^{1/2} \\ \geq (\leq) \frac{1}{n}, & \text{otherwise} \end{cases}.$$

REFERENCES

- [1] Biernacki, M., Pidek, H. and Ryll-Nardzewski, C. (1950), Sur une inégalité entre des intégrales définies, *Ann. Univ. Mariae Curie-Skolodowska*, **A4**, 1-4.
- [2] Andrica, D. and Badea, C. (1988), Grüss' inequality for positive linear functionals, *Periodica Math. Hungarica*, **19**(2), 155-167.
- [3] Dragomir, S.S. and Booth, G.L. (2000), On a Grüss-Lupaş type inequality and its application for the estimation of p -moments of guessing mappings, *Math. Comm.*, **5**, 117-126.
- [4] Dragomir, S.S. (2002), Another Grüss type inequality for sequences of vectors in normed linear spaces and applications, *J. Comp. Analysis & Appl.*, **4**(2), 157-172.
- [5] Dragomir, S.S. (2002), A Grüss type inequality for sequences of vectors in normed linear spaces, (Preprint) *RGMIA Res. Rep. Coll.*, **5**(2), Article 9. (ONLINE: <http://rgmia.vu.edu.au/v5n2.html>)
- [6] Dragomir, S. S. (2001), Integral Grüss inequality for mappings with values in Hilbert spaces and applications. *J. Korean Math. Soc.* **38**, no. 6, 1261-1273.

- [7] Dragomir, S. S. (1999), A generalization of Grüss's inequality in inner product spaces and applications. *J. Math. Anal. Appl.* **237**, no. 1, 74–82
- [8] Cerone, P. and Dragomir, S.S. (2002), A refinement of Grüss' inequality and applications, *RGMIA Res. Rep. Coll.*, **5**(2002), No.2, Article 15. (ONLINE: <http://rgmia.vu.edu.au/v5n2.html>)
- [9] Dragomir, S.S. and Pečarić, J.(1989), Refinements of some inequalities for isotonic functionals, *Anal. Num. Theor. Approx.*, **18**, 61-65.
- [10] Dragomir, S.S. (2002), A companion of the Grüss inequality and applications, *RGMIA Res. Rep. Coll.*, **5**, Supplement, Article 13. (ONLINE: [http://rgmia.vu.edu.au/v5\(E\).html](http://rgmia.vu.edu.au/v5(E).html))
- [11] Fink, A. M. (1999), A treatise on Grüss' inequality. *Analytic and geometric inequalities and applications, 93–113, Math. Appl.*, 478, Kluwer Acad. Publ., Dordrecht,
- [12] Pečarić, J. (1980), On some inequalities analogous to Grüss inequality. *Mat. Vesnik* **4(17)(32)**, no. 2, 197–202.
- [13] Massey J.L. (1994), Guessing and entropy, *Proc. 1994 IEEE Int. Symp. on Inf. Th.*, (Trondheim, Norway, 1994), p. 204.
- [14] Arikan E. (1996), An inequality on guessing and its application to sequential decoding, *IEEE Tran. Inf. Th.*, **42**(1), 99-105.
- [15] Boztaş S. (1997), Comments on “An Inequality of Guessing and Its Applications to Sequential Decoding”, *IEEE Tran. Inf. Th.*, **43**(6), 2062-2063.
- [16] Dragomir S.S. and Boztaş S. (1997), Some estimates of the average number of guesses to determine a random variable, *Proc. 1997 IEEE Int. Symp. on Inf. Th.*, (Ulm, Germany, 1997), p. 159.
- [17] Dragomir S.S. and Boztaş S. (1998), Estimation of arithmetic means and their applications in guessing theory, *Math. Comput. Modelling*, **28**(10) (1998), 31-43.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MCMC 8001, VICTORIA, AUSTRALIA

E-mail address: pc@matilda.vu.edu.au

URL: <http://rgmia.vu.edu.au/cerone>

E-mail address: sever@matilda.vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>