

Hermite-Hadamard type inequalities for increasing radiant functions

E.V. Sharikov

Tver State University, Tver, Russia

Abstract

We study Hermite-Hadamard type inequalities for increasing radiant functions and give some simple examples of such inequalities.

Keywords: Increasing radiant functions; Abstract convexity; Hermite-Hadamard type inequalities.

1 Introduction

In this paper we consider one generalization of Hermite-Hadamard inequalities for the class InR of increasing radiant functions defined on the cone $\mathbb{R}_{++}^n = \{x \in \mathbb{R}^n : x_i > 0 \ (i = 1, \dots, n)\}$.

Recall that for a function $f : [a, b] \rightarrow \mathbb{R}$, which is convex on $[a, b]$, we have the following:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{1}{2}(f(a) + f(b)). \quad (1)$$

These inequalities are well known as the Hermite-Hadamard inequalities. There are many generalizations of these inequalities for classes of nonconvex functions. For more information see ([2], Section 6.5), [1] and references therein. In this paper we consider generalizations of the inequalities from the both sides of (1). Some technique and notions, which are used here, can be found in [1].

In Section 2 of this paper we give definition if InR functions and recall some results related to these functions. In Section 3 we consider Hermite-Hadamard type inequalities for the class InR. Some examples of such inequalities for functions defined on \mathbb{R}_{++} and \mathbb{R}_{++}^2 are given in Section 4.

2 Preliminaries

We assume that the cone \mathbb{R}_{++}^n is equipped with coordinate-wise order relation.

Recall that a function $f : \mathbb{R}_{++}^n \rightarrow \bar{\mathbb{R}}_+ = [0, +\infty]$ is called increasing radiant (InR) if:

1. f is increasing: $x \geq y \implies f(x) \geq f(y)$;
2. f is radiant: $f(\lambda x) \leq \lambda f(x)$ for all $\lambda \in (0, 1)$ and $x \in \mathbb{R}_{++}^n$.

For example, any function f of the following form belongs to the class InR:

$$f(x) = \sum_{|k| \geq 1} c_k x_1^{k_1} \cdots x_n^{k_n},$$

where $k = (k_1, \dots, k_n)$, $|k| = k_1 + \dots + k_n$, $k_i \geq 0$, $c_k \geq 0$.

For each $f \in \text{InR}$ its conjugate function ([4])

$$f^*(x) = \frac{1}{f(1/x)},$$

where $1/x = (1/x_1, \dots, 1/x_n)$, is also increasing and radiant. Hence any function

$$f(x) = \frac{1}{\sum_{|k| \geq 1} c_k x_1^{-k_1} \cdots x_n^{-k_n}}$$

is InR. In more general case we have the following InR functions:

$$f(x) = \left(\frac{\sum_{|k| \geq u} c_k x_1^{k_1} \cdots x_n^{k_n}}{\sum_{|k| \geq v} d_k x_1^{-k_1} \cdots x_n^{-k_n}} \right)^t,$$

where $u, v > 0$, $t \geq 1/(u+v)$. Indeed, these functions are increasing and for any $\lambda \in (0, 1)$

$$\begin{aligned} f(\lambda x) &= \left(\frac{\sum_{|k| \geq u} \lambda^{|k|} c_k x_1^{k_1} \cdots x_n^{k_n}}{\sum_{|k| \geq v} \lambda^{-|k|} d_k x_1^{-k_1} \cdots x_n^{-k_n}} \right)^t \leq \\ &= \left(\frac{\lambda^u \sum_{|k| \geq u} c_k x_1^{k_1} \cdots x_n^{k_n}}{\lambda^{-v} \sum_{|k| \geq v} d_k x_1^{-k_1} \cdots x_n^{-k_n}} \right)^t = \lambda^{(u+v)t} f(x) \leq \lambda f(x). \end{aligned}$$

Consider the coupling function φ defined on $\mathbb{R}_{++}^n \times \mathbb{R}_{++}^n$:

$$\varphi(h, x) = \begin{cases} 0, & \text{if } \langle h, x \rangle < 1, \\ \langle h, x \rangle, & \text{if } \langle h, x \rangle \geq 1, \end{cases} \quad (2)$$

where

$$\langle h, x \rangle = \min\{h_i x_i : i = 1, \dots, n\}$$

is the so-called min-type function.

Denote by φ_h the function defined on \mathbb{R}_{++}^n by the formula: $\varphi_h(x) = \varphi(h, x)$.

It is known (see [4]) that the set

$$H = \left\{ \frac{1}{c} \varphi_h : h \in \mathbb{R}_{++}^n, c \in (0, +\infty] \right\}$$

is the supremal generator of the class InR of all increasing radiant functions defined on \mathbb{R}_{++}^n .

It is known also that for any InR function f

$$f(h)\varphi\left(\frac{1}{h}, x\right) \leq f(x) \quad \text{for all } x, h \in \mathbb{R}_{++}^n. \quad (3)$$

Note that for $c = +\infty$ we set $c\varphi_h(x) = \sup_{l>0} (l\varphi_h(x))$.

Formula (3) implies the following statement.

Proposition 2.1 *Let f be an InR function defined on \mathbb{R}_{++}^n and $\Delta \subset \mathbb{R}_{++}^n$. Then the function*

$$f_\Delta(x) = \sup_{h \in \Delta} f(h)\varphi\left(\frac{1}{h}, x\right)$$

is InR, and it possesses the properties:

- 1.) $f_\Delta(x) \leq f(x)$ for all $x \in \mathbb{R}_{++}^n$,
- 2.) $f_\Delta(x) = f(x)$ for all $x \in \Delta$.

3 Hermite-Hadamard type inequalities

Let $D \subset \mathbb{R}_{++}^n$ be a closed domain (in topology of \mathbb{R}_{++}^n), i.e. D is bounded set such that $\text{cl int } D = D$. Denote by $Q(D)$ the set of all points $\bar{x} \in D$ such that

$$\frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1, \quad (4)$$

where $A(D) = \int_D dx$, $dx = dx_1 \cdots dx_n$.

Proposition 3.1 *Let f be an InR function defined on \mathbb{R}_{++}^n . If the set $Q(D)$ is nonempty and f is integrable on D then*

$$\sup_{\bar{x} \in Q(D)} f(\bar{x}) \leq \frac{1}{A(D)} \int_D f(x) dx. \quad (5)$$

Proof: First, let $\bar{x} \in Q(D)$ and $f(\bar{x}) < +\infty$. Then $f(\bar{x})\varphi(1/\bar{x}, x) \leq f(x)$ for all $x \in D \subset \mathbb{R}_{++}^n$ (see (3)). By (4), we get

$$f(\bar{x}) = f(\bar{x}) \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \frac{1}{A(D)} \int_D f(\bar{x})\varphi\left(\frac{1}{\bar{x}}, x\right) dx \leq \frac{1}{A(D)} \int_D f(x) dx.$$

Now, suppose that $f(\bar{x}) = +\infty$. Then for all $l > 0$ function $l\varphi_{1/\bar{x}}(x)$ is minorant of f . Hence $l \leq (1/A(D)) \int_D f(x) dx \quad \forall l > 0$, that implies that function f is not integrable on D . This contradiction shows that $f(\bar{x}) < +\infty$ for any $\bar{x} \in Q(D)$. \square

As it was done in [1], we may introduce the set $Q_m(D)$ of all maximal elements of $Q(D)$. It means that a point $\bar{x} \in Q(D)$ belongs to $Q_m(D)$ if and only if for any $\bar{y} \in Q(D)$: $(\bar{y} \geq \bar{x}) \implies (\bar{y} = \bar{x})$. Suppose that the set $Q(D)$ is nonempty. It is easy to see that $Q(D)$ is closed set in topology of \mathbb{R}_{++}^n . Hence, using Zorn Lemma we conclude that $Q_m(D)$ is nonempty closed set and for any $\bar{x} \in Q(D)$ there exists $\bar{y} \in Q_m(D)$, for which $\bar{x} \leq \bar{y}$.

So, in assumptions of Proposition 3.1 we have the following estimate:

$$\sup_{\bar{x} \in Q_m(D)} f(\bar{x}) \leq \frac{1}{A(D)} \int_D f(x) dx. \quad (6)$$

Since f is increasing function then this inequality implies inequality (5).

Remark 3.1 Let $D \subset \mathbb{R}_{++}^n$ be a closed domain and the set $Q(D)$ is nonempty. Then for every $\bar{x} \in Q(D)$ inequality

$$f(\bar{x}) \leq \frac{1}{A(D)} \int_D f(x) dx$$

is sharp. For example, if we set $f = \varphi_{1/\bar{x}}$ then (see (4))

$$f(\bar{x}) = \varphi\left(\frac{1}{\bar{x}}, \bar{x}\right) = 1 = \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \frac{1}{A(D)} \int_D f(x) dx.$$

Note that here we used only the values of function f on a set D . Therefore we need the following definition.

Definition 3.1 Let $D \subset \mathbb{R}_{++}^n$. A function $f : D \rightarrow [0, +\infty]$ is called increasing radiant on D if there exists an InR function F defined on \mathbb{R}_{++}^n such that $F|_D = f$, that is $F(x) = f(x)$ for all $x \in D$.

We assume here, as above, that for $c = +\infty$: $c\varphi_h(x) = \sup_{l>0}(l\varphi_h(x))$.

Proposition 3.2 Let $f : D \rightarrow [0, +\infty]$ be a function defined on $D \subset \mathbb{R}_{++}^n$. Then the following assertions are equivalent:

- 1.) f is increasing radiant on D ,
- 2.) $f(h)\varphi(1/h, x) \leq f(x)$ for all $h, x \in D$,
- 3.) f is abstract convex with respect to the set of functions $(1/c)\varphi_{(1/h)} : D \rightarrow [0, +\infty]$ with $h \in D, c \in (0, +\infty]$.

Proof: 1.) \implies 2.) By Definition 3.1, there exists an InR function $F : \mathbb{R}_{++}^n \rightarrow [0, +\infty]$ such that $F(x) = f(x)$ for all $x \in D$. Then Proposition 2.1 implies that the function

$$F_D(x) = \sup_{h \in D} F(h)\varphi\left(\frac{1}{h}, x\right)$$

interpolates F in all points $x \in D$. Hence

$$\sup_{h \in D} f(h)\varphi\left(\frac{1}{h}, x\right) = f(x) \text{ for all } x \in D,$$

that implies the assertion 2.)

2.) \implies 3.) Consider the function f_D defined on D

$$f_D(x) = \sup_{h \in D} f(h)\varphi\left(\frac{1}{h}, x\right).$$

First, it is clear that f_D is abstract convex with respect to the set of functions defined on D : $\{(1/c)\varphi(1/h) : h \in D, c \in (0, +\infty)\}$. Further, using 2.) we get for all $x \in D$

$$f_D(x) \leq f(x) = f(x)\varphi\left(\frac{1}{x}, x\right) \leq \sup_{h \in D} f(h)\varphi\left(\frac{1}{h}, x\right) = f_D(x).$$

So, $f_D(x) = f(x)$ for all $x \in D$ and we have the desired statement 3.)

3.) \implies 1.) It is obvious since any function $(1/c)\varphi_h$ defined on D can be considered as elementary function $(1/c)\varphi_h \in H$ defined on \mathbb{R}_{++}^n . \square

Remark 3.2 We may require in Proposition 3.1, formula (6) and Remark 3.1 only that function f is increasing radiant and integrable on D .

Remark 3.3 We may consider more general case of Hermite-Hadamard type inequalities for InR functions. Let f be an increasing radiant function on D . Then Proposition 3.2 implies that $f(h)\varphi(1/h, x) \leq f(x)$ for all $h, x \in D$. If $f(\bar{x}) < +\infty$ and f is integrable on D then

$$f(\bar{x}) \int_D \varphi(1/\bar{x}, x) dx \leq \int_D f(x) dx. \quad (7)$$

This inequality is sharp for any $\bar{x} \in D$ since we have the equality in (7) for $f = \varphi(1/\bar{x})$.

Proposition 3.2 implies also that the class InR is broad enough.

Proposition 3.3 Let $S \subset \mathbb{R}_{++}^n$ be a set such that every point $x \in S$ is maximal in S . Then for any function $f : S \rightarrow [0, +\infty]$ there exists an increasing radiant function $F : \mathbb{R}_{++}^n \rightarrow [0, +\infty]$, for which $F|_S = f$.

Proof: It is sufficiently to check only that $f(h)\varphi(1/h, x) \leq f(x)$ for all $h, x \in S$. If $h = x$ then $\varphi(1/h, x) = 1$, $f(h) = f(x)$. If $h \neq x$ then $\langle 1/h, x \rangle = \min_i x_i/h_i < 1$ since h is maximal point in S , hence $\varphi(1/h, x) = 0$ and $f(h)\varphi(1/h, x) = 0 \leq f(x)$. \square

In particular, Proposition 3.3 holds if $S = \{x \in \mathbb{R}_{++}^n : (x_1)^p + \dots + (x_n)^p = 1\}$, where $p > 0$.

Now we present two assertions supported by definition of function φ . Recall that a set $\Omega \subset \mathbb{R}_{++}^n$ is called normal if for each $x \in \Omega$ we have $(y \in \Omega \text{ for all } y \leq x)$. Normal hull $N(\Omega)$ of a set Ω is defined as follows: $N(\Omega) = \{x \in \mathbb{R}_{++}^n : (\exists y \in \Omega) x \leq y\}$ (see, for example, [3]).

Proposition 3.4 *Let $D, \Omega \subset \mathbb{R}_{++}^n$ be a closed domains and $D \subset \Omega$. If the set $Q(\Omega)$ is nonempty and*

$$(\Omega \setminus D) \subset N(Q(\Omega)) \quad (8)$$

then the set $Q(D)$ consists of all points $\bar{x} \in \Omega$ such that

$$\frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1.$$

Proof: If $D = \Omega$ then the assertion is clear. Assume that $D \neq \Omega$. Since D, Ω are closed domains and $D \subset \Omega$ then

$$A(D) < A(\Omega). \quad (9)$$

Let $\bar{x} \in \Omega$ and

$$\frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1. \quad (10)$$

We show that $\varphi(1/\bar{x}, x) = 0$ for all $x \in \Omega \setminus D$. If $x \in \Omega \setminus D$ then, by (8), there exists a point $\bar{y} \in Q(\Omega)$: $\bar{y} \geq x$; hence $\langle 1/\bar{x}, x \rangle \leq \langle 1/\bar{x}, \bar{y} \rangle$. Suppose that $\langle 1/\bar{x}, \bar{y} \rangle \geq 1$. Then $\bar{y} \geq \bar{x} \implies 1/\bar{y} \leq 1/\bar{x}$. Since $\bar{y} \in Q(\Omega)$ then, by (9) and (10)

$$1 = \frac{1}{A(\Omega)} \int_{\Omega} \varphi\left(\frac{1}{\bar{y}}, x\right) dx < \frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{y}}, x\right) dx \leq \frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1.$$

So, we have the inequalities: $\langle 1/\bar{x}, x \rangle \leq \langle 1/\bar{x}, \bar{y} \rangle < 1$. Therefore $\varphi(1/\bar{x}, x) = 0$ for all $x \in \Omega \setminus D \implies$

$$1 = \frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx.$$

The equality ($\varphi(1/\bar{x}, \cdot) = 0$ on $\Omega \setminus D$) implies also that $\bar{x} \neq x$ for all $x \in \Omega \setminus D$, hence $\bar{x} \notin \Omega \setminus D \implies \bar{x} \in D$. Thus, we have the established result: $\bar{x} \in Q(D)$.

Conversely, let $\bar{x} \in Q(D)$. For any $x \in \Omega \setminus D$ there exists $\bar{y} \in Q(\Omega)$ such that $\bar{y} \geq x \implies \langle 1/\bar{x}, x \rangle \leq \langle 1/\bar{x}, \bar{y} \rangle$. Moreover, we may assume that \bar{y} is maximal point in $Q(\Omega)$, i.e. $\bar{y} \in Q_m(\Omega)$. First, we check that

$$\left\langle \frac{1}{\bar{y}}, x \right\rangle \leq 1 \text{ for all } x \in \Omega \setminus D, \bar{y} \in Q_m(\Omega). \quad (11)$$

Indeed, if $x \in \Omega \setminus D$ then for some $\bar{z} \in Q_m(\Omega)$: $x \leq \bar{z} \implies \langle 1/\bar{y}, x \rangle \leq \langle 1/\bar{y}, \bar{z} \rangle$. But $\langle 1/\bar{y}, \bar{z} \rangle \leq 1$ since $\bar{y}, \bar{z} \in Q_m(\Omega)$ (otherwise, if $\langle 1/\bar{y}, \bar{z} \rangle > 1$ then $\bar{z} > \bar{y} \implies \bar{y} \notin Q_m(\Omega)$). Now we verify that $\langle 1/\bar{x}, x \rangle < 1$ for all $x \in \Omega \setminus D$. If $x \in \Omega \setminus D$ then for some $\bar{y} \in Q_m(\Omega)$: $\langle 1/\bar{x}, x \rangle \leq \langle 1/\bar{x}, \bar{y} \rangle$. Suppose that $\langle 1/\bar{x}, \bar{y} \rangle \geq 1$. Then $\bar{y} \geq \bar{x}$ and therefore, using inclusion $\bar{x} \in Q(D)$, we get

$$1 = \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx > \frac{1}{A(\Omega)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx \geq \frac{1}{A(\Omega)} \int_D \varphi\left(\frac{1}{\bar{y}}, x\right) dx. \quad (12)$$

Let $D_1 = \{x \in \Omega \setminus D : \langle 1/\bar{y}, x \rangle < 1\}$, $D_2 = \{x \in \Omega \setminus D : \langle 1/\bar{y}, x \rangle = 1\}$. It follows from (11) that $\Omega \setminus D = D_1 \cup D_2$ ($D_1 \cap D_2 = \emptyset$), hence

$$\int_{\Omega \setminus D} \varphi\left(\frac{1}{\bar{y}}, x\right) dx = \int_{D_1} \varphi\left(\frac{1}{\bar{y}}, x\right) dx + \int_{D_2} \varphi\left(\frac{1}{\bar{y}}, x\right) dx = \int_{D_2} \varphi\left(\frac{1}{\bar{y}}, x\right) dx = \int_{D_2} dx.$$

But the last integral $\int_{D_2} dx$ is also equal to zero, since the set D_2 has no interior points. Thus, by (12)

$$1 > \frac{1}{A(\Omega)} \int_D \varphi\left(\frac{1}{\bar{y}}, x\right) dx = \frac{1}{A(\Omega)} \int_{\Omega} \varphi\left(\frac{1}{\bar{y}}, x\right) dx.$$

This inequality contradicts to the inclusion $\bar{y} \in Q_m(\Omega)$. So, we conclude that the inequality $\langle 1/\bar{x}, \bar{y} \rangle \geq 1$ is impossible. Hence $\langle 1/\bar{x}, x \rangle \leq \langle 1/\bar{x}, \bar{y} \rangle < 1$ for all $x \in \Omega \setminus D$ and $\bar{y} = \bar{y}(x) \in Q_m(\Omega)$, that implies required equality:

$$1 = \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx.$$

□

Corollary 3.1 Let $D_1, D_2 \subset \mathbb{R}_{++}^n$ be a closed domains such that

$$A(D_1) = A(D_2).$$

If there exists a closed domain $\Omega \subset \mathbb{R}_{++}^n$, for which the set $Q(\Omega)$ is nonempty and

$$D_i \subset \Omega, \quad (\Omega \setminus D_i) \subset N(Q(\Omega)) \quad (i = 1, 2),$$

then

$$Q(D_1) = Q(D_2).$$

Proposition 3.5 Let $D, \Omega \subset \mathbb{R}_{++}^n$ be a closed domains and $D \subset \Omega$. If

$$N(\Omega \setminus D) \cap D = \emptyset \tag{13}$$

then the set $Q(D)$ consists of all points $\bar{x} \in D$ such that

$$\frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1.$$

Proof: Formula (13) implies that if $\bar{x} \in D$ then $\bar{x} \notin N(\Omega \setminus D)$. It means that for all $x \in \Omega \setminus D$: $x < \bar{x} \implies \langle 1/\bar{x}, x \rangle < 1 \implies \varphi(1/\bar{x}, x) = 0$.

Thus, for any $\bar{x} \in D$

$$\frac{1}{A(D)} \int_{\Omega} \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1 \iff \frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1 \iff \bar{x} \in Q(D).$$

□

Now consider the generalization of the inequality from the right-hand side of (1). Let f be an increasing radiant function defined on a closed domain $D \subset \mathbb{R}_{++}^n$, and f is integrable on D . Then $f(h)\varphi(1/h, x) \leq f(x)$ for all $h, x \in D$. In particular, $f(h)\langle 1/h, x \rangle \leq f(x)$ if $\langle 1/h, x \rangle \geq 1$. Hence for all $x \geq h$

$$f(h) \leq \frac{f(x)}{\langle 1/h, x \rangle} = \langle h, 1/x \rangle^+ f(x),$$

where $h(y) = \langle h, y \rangle^+ = \max_i h_i y_i$ is the so-called max-type function. So, if $\bar{x} \in D$ and $\bar{x} \geq x$ for all $x \in D$, then $f(x) \leq \langle x, 1/\bar{x} \rangle^+ f(\bar{x})$ for any $\bar{x} \in D$. This reduces to the following assertion.

Proposition 3.6 *Let function f be an increasing radiant and integrable on D . If $\bar{x} \in D$ and $\bar{x} \geq x$ for all $x \in D$, then*

$$\int_D f(x) dx \leq f(\bar{x}) \int_D \langle x, 1/\bar{x} \rangle^+ dx. \quad (14)$$

Inequality (14) is sharp since we get equality for $f(x) = \langle x, 1/\bar{x} \rangle^+$.

In more general case we have the following inequalities:

$$f(x) \leq \langle x, 1/\bar{x} \rangle^+ \sup_{y \in D} f(y) \quad \text{for all } \bar{x} \geq x.$$

Hence

$$f(x) \leq \sup_{y \in D} f(y) \inf\{\langle x, 1/\bar{x} \rangle^+ : \bar{x} \geq x, \bar{x} \in D\} \quad \text{for all } x \in D$$

and therefore

$$\int_D f(x) dx \leq \sup_{y \in D} f(y) \int_D \inf\{\langle x, 1/\bar{x} \rangle^+ : \bar{x} \geq x, \bar{x} \in D\} dx. \quad (15)$$

4 Examples

Here we describe the set $Q(D)$ for some special domains D of the cones \mathbb{R}_{++} and \mathbb{R}_{++}^2 .

Let $a, b \in \mathbb{R}$ be a numbers such that $0 \leq a < b$. We denote by $[a, b]$ the segment $\{x \in \mathbb{R}_{++} : a \leq x \leq b\}$.

Example 4.1 Let $D = [a, b] \subset \mathbb{R}_{++}$, where $0 \leq a < b$. According to definition, the set $Q(D)$ consists of all points $\bar{x} \in D$, for which

$$\frac{1}{A(D)} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \frac{1}{b-a} \int_a^b \varphi\left(\frac{1}{\bar{x}}, x\right) dx = 1.$$

We have:

$$\varphi\left(\frac{1}{\bar{x}}, x\right) = \begin{cases} 0, & \text{if } x < \bar{x}, \\ x/\bar{x}, & \text{if } x \geq \bar{x}. \end{cases}$$

Hence, if $\bar{x} \in D = [a, b]$ then

$$\int_a^b \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \int_{\bar{x}}^b \frac{x}{\bar{x}} dx = \frac{1}{2\bar{x}}(b^2 - \bar{x}^2). \quad (16)$$

So, a point $\bar{x} \in [a, b]$ belongs to $Q(D)$ if and only if

$$\frac{1}{2(b-a)\bar{x}}(b^2 - \bar{x}^2) = 1 \iff \bar{x}^2 + 2(b-a)\bar{x} - b^2 = 0.$$

We get

$$\bar{x} = \sqrt{(b-a)^2 + b^2} - (b-a). \quad (17)$$

Show that for the point (17)

$$a < \bar{x} < \frac{a+b}{2}. \quad (18)$$

Since $b > a \geq 0$ then $\bar{x} = \sqrt{(b-a)^2 + b^2} - (b-a) > \sqrt{b^2} - (b-a) = a$. Further,

$$\begin{aligned} \bar{x} < \frac{a+b}{2} &\iff \sqrt{(b-a)^2 + b^2} < (b-a) + \frac{a+b}{2} = \frac{3b-a}{2} \iff \\ &4(b-a)^2 + 4b^2 < (3b-a)^2 \iff 0 < b^2 + 2ab - 3a^2. \end{aligned}$$

The last inequality follows from the same conditions $b > a \geq 0$.

Thus, $Q([a, b]) = \{\sqrt{(b-a)^2 + b^2} - (b-a)\}$. Remark 3.1 implies that for every InR function $f \in L_1[a, b]$

$$f\left(\sqrt{(b-a)^2 + b^2} - (b-a)\right) \leq \frac{1}{b-a} \int_a^b f(x) dx$$

and this inequality is sharp. (Compare it with the corresponding estimate for convex functions (1), see also (18)).

Remark 3.3 and formula (16) imply the following inequalities

$$f(u) \leq \frac{2u}{b^2 - u^2} \int_a^b f(x) dx, \quad (19)$$

which are sharp in the class of all InR functions $f \in L_1[a, b]$ and hold for any $u \in [a, b]$. In particular, we get for $u = (a+b)/2$

$$f\left(\frac{a+b}{2}\right) \leq \frac{4(a+b)}{(a+3b)(b-a)} \int_a^b f(x) dx.$$

Note that here

$$\frac{4(a+b)}{(a+3b)(b-a)} > \frac{1}{b-a}.$$

Further, Proposition 3.6 implies that

$$\int_a^b f(x) dx \leq f(b) \int_a^b \frac{x}{b} dx = \frac{b^2 - a^2}{2b} f(b),$$

hence

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{a+b}{2b} f(b)$$

for every InR function $f \in L_1[a, b]$.

Let $D \subset \mathbb{R}_{++}^2$, $\bar{x} = (\bar{x}_1, \bar{x}_2) \in D$. We denote by $D(\bar{x})$ the set $\{x \in D : x_1 \geq \bar{x}_1, x_2 \geq \bar{x}_2\}$. It is clear that

$$\int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx = \int_{D(\bar{x})} \left\langle \frac{1}{\bar{x}}, x \right\rangle dx = \int_{D(\bar{x})} \min\left(\frac{x_1}{\bar{x}_1}, \frac{x_2}{\bar{x}_2}\right) dx_1 dx_2.$$

In order to calculate such integral we represent the set $D(\bar{x})$ as union $D_1(\bar{x}) \cup D_2(\bar{x})$, where

$$D_1(\bar{x}) = \left\{x \in D(\bar{x}) : \frac{x_2}{\bar{x}_2} \leq \frac{x_1}{\bar{x}_1}\right\}, \quad D_2(\bar{x}) = \left\{x \in D(\bar{x}) : \frac{x_1}{\bar{x}_1} \leq \frac{x_2}{\bar{x}_2}\right\}.$$

Then

$$\begin{aligned} \int_D \varphi\left(\frac{1}{\bar{x}}, x\right) dx &= \int_{D_1(\bar{x})} \langle 1/\bar{x}, x \rangle dx + \int_{D_2(\bar{x})} \langle 1/\bar{x}, x \rangle dx = \\ &= \frac{1}{\bar{x}_2} \int_{D_1(\bar{x})} x_2 dx_1 dx_2 + \frac{1}{\bar{x}_1} \int_{D_2(\bar{x})} x_1 dx_1 dx_2. \end{aligned}$$

In the next examples we will use the number k , which possesses properties:

$$2k^3 - 3k^2 - 3k + 1 = 0, \quad 0 < k < 1. \quad (20)$$

Let $g(k) = 2k^3 - 3k^2 - 3k + 1$. We have: $g(0) > 0$, $g(1) < 0$, $g'(k) = 6k^2 - 6k - 3 < 6k - 6k - 3 < 0$ for all $k \in (0, 1)$. So, there exists unique solution of the equation (20), which belongs to interval $(0, 1)$. We denote this solution by the same symbol k .

Example 4.2 Let $D \subset \mathbb{R}_{++}^2$ be the triangle with vertices $(0, 0)$, $(a, 0)$ and $(0, b)$, that is

$$D = \left\{x \in \mathbb{R}_{++}^2 : \frac{x_1}{a} + \frac{x_2}{b} \leq 1\right\}.$$

If $\bar{x} \in D$ then we get

$$\begin{aligned} D_1(\bar{x}) &= \left\{x \in \mathbb{R}_{++}^2 : \bar{x}_2 \leq x_2 \leq \frac{ab\bar{x}_2}{a\bar{x}_2 + b\bar{x}_1}, \frac{\bar{x}_1}{\bar{x}_2} x_2 \leq x_1 \leq a - \frac{a}{b} x_2\right\}, \\ D_2(\bar{x}) &= \left\{x \in \mathbb{R}_{++}^2 : \bar{x}_1 \leq x_1 \leq \frac{ab\bar{x}_1}{a\bar{x}_2 + b\bar{x}_1}, \frac{\bar{x}_2}{\bar{x}_1} x_1 \leq x_2 \leq b - \frac{b}{a} x_1\right\}. \end{aligned}$$

Therefore

$$\int_{D_1(\bar{x})} \langle 1/\bar{x}, x \rangle dx = \frac{1}{\bar{x}_2} \int_{\bar{x}_2}^{(ab\bar{x}_2)/(a\bar{x}_2+b\bar{x}_1)} dx_2 \int_{(\bar{x}_1/\bar{x}_2)x_2}^{a-(a/b)x_2} x_2 dx_1.$$

This reduces to

$$\int_{D_1(\bar{x})} \langle 1/\bar{x}, x \rangle dx = \frac{ab}{6} \frac{\bar{x}_2/b}{(\bar{x}_1/a + \bar{x}_2/b)^2} - \frac{ab \bar{x}_2}{2b} + \frac{ab \bar{x}_2}{3b} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right).$$

By analogy,

$$\int_{D_2(\bar{x})} \langle 1/\bar{x}, x \rangle dx = \frac{ab}{6} \frac{\bar{x}_1/a}{(\bar{x}_1/a + \bar{x}_2/b)^2} - \frac{ab \bar{x}_1}{2a} + \frac{ab \bar{x}_1}{3a} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right).$$

Thus, the sum of these quantities is

$$\int_D \varphi \left(\frac{1}{\bar{x}}, x \right) dx = \frac{ab}{6} \frac{1}{(\bar{x}_1/a + \bar{x}_2/b)} - \frac{ab}{2} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + \frac{ab}{3} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2. \quad (21)$$

Since $A(D) = (ab)/2$ then for $\bar{x} \in D$

$$\begin{aligned} \bar{x} \in Q(D) &\iff \frac{1}{3} \frac{1}{(\bar{x}_1/a + \bar{x}_2/b)} - \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + \frac{2}{3} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 = 1 \\ &\iff 2 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^3 - 3 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 - 3 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + 1 = 0. \end{aligned}$$

Using inequalities $0 < (\bar{x}_1/a + \bar{x}_2/b) \leq 1$ for $\bar{x} \in D$ we get

$$Q(D) = \left\{ \bar{x} \in \mathbb{R}_{++}^2 : \frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} = k \right\},$$

where k is the solution of (20).

In more general case we have inequality (see (7) and (21))

$$f(\bar{x}_1, \bar{x}_2) \leq \frac{6u}{ab(1 - 3u^2 + 2u^3)} \int_D f(x) dx,$$

where $u = u(\bar{x}_1, \bar{x}_2) = \bar{x}_1/a + \bar{x}_2/b < 1$, function f is increasing radiant and integrable on D .

Consider now inequality (15) for our triangle D . We show that $\inf\{\langle x, 1/\bar{x} \rangle^+ : \bar{x} \geq x, \bar{x} \in D\} = (x_1/a + x_2/b)$. Let $\bar{x} = (\bar{x}_1, \bar{x}_2) = (x_1/(x_1/a + x_2/b), x_2/(x_1/a + x_2/b))$. Then $\bar{x} \geq x$ and $\bar{x} \in D$ since $\bar{x}_1/a + \bar{x}_2/b = 1$. Hence

$$\inf\{\langle x, 1/\bar{x} \rangle^+ : \bar{x} \geq x, \bar{x} \in D\} \leq \max \left\{ x_1 \frac{(x_1/a + x_2/b)}{x_1}, x_2 \frac{(x_1/a + x_2/b)}{x_2} \right\} = \frac{x_1}{a} + \frac{x_2}{b}.$$

Suppose that the converse inequality does not hold, then $\langle x, 1/\bar{x} \rangle^+ < x_1/a + x_2/b$ for some $\bar{x} \geq x, \bar{x} \in D$, hence $x/(x_1/a + x_2/b) < \bar{x}$. But this implies that $\bar{x} \notin D$.

So, it follows from (15) that

$$\int_D f(x) dx \leq \sup_{y \in D} f(y) \int_D \left(\frac{x_1}{a} + \frac{x_2}{b} \right) dx.$$

Calculation gives the quantity

$$\int_D \left(\frac{x_1}{a} + \frac{x_2}{b} \right) dx = \frac{ab}{3}.$$

Since $A(D) = ab/2$ then the final result is

$$\frac{1}{A(D)} \int_D f(x) dx \leq \frac{2}{3} \sup_{y \in D} f(y).$$

Example 4.3 Let now Ω be the triangle from the Example 4.2:

$$\Omega = \left\{ x \in \mathbb{R}_{++}^2 : \frac{x_1}{a} + \frac{x_2}{b} \leq 1 \right\}.$$

Denote by D the subset of Ω such that

$$\Omega \setminus D = \left\{ x \in \Omega : \frac{k}{3} < \frac{x_1}{a}, \frac{k}{3} < \frac{x_2}{b}, \frac{x_1}{a} + \frac{x_2}{b} < k \right\}.$$

Then $(\Omega \setminus D) \subset N(Q(\Omega)) = \{x \in \mathbb{R}_{++}^2 : x_1/a + x_2/b \leq k\}$. Note that $A(\Omega \setminus D) = (1/18)k^2ab$, hence $A(D) = (ab)/2 - (1/18)k^2ab = ab(1/2 - k^2/18)$. It follows from Proposition 3.4 and formula (21) (with Ω instead of D) that a point $\bar{x} \in \Omega$ belongs to $Q(D)$ if and only if

$$\begin{aligned} \frac{1}{ab(1/2 - k^2/18)} \left[\frac{ab}{6} \frac{1}{(\bar{x}_1/a + \bar{x}_2/b)} - \frac{ab}{2} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + \frac{ab}{3} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 \right] = 1 &\iff \\ 2 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^3 - 3 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 - \left(3 - \frac{k^2}{3} \right) \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + 1 = 0. \end{aligned}$$

It is easy to check that there exists unique solution s of the equation:

$$2s^3 - 3s^2 - (3 - k^2/3)s + 1 = 0, \quad 0 < s \leq 1.$$

Hence

$$Q(D) = \left\{ \bar{x} \in \mathbb{R}_{++}^2 : \frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} = s \right\}.$$

We may establish also that $s > k$.

Remark 4.1 For any other closed domain D' such that $(\Omega \setminus D') \subset N(Q(\Omega)) = \{x \in \mathbb{R}_{++}^2 : x_1/a + x_2/b \leq k\}$ the set $Q(D')$ has the same form, i.e. it is intersection of \mathbb{R}_{++}^2 and a line $(\bar{x}_1/a + \bar{x}_2/b) = s'$ with some s' : $k < s' < 1$.

Example 4.4 Let Ω be the same triangle: $\Omega = \{x \in \mathbb{R}_{++}^2 : (x_1/a + x_2/b) \leq 1\}$. Let $D \subset \Omega$ and

$$\Omega \setminus D = \{x \in \Omega : x_1 < a/2, x_2 < b/2\}.$$

Then $\Omega \setminus D$ is the normal set, hence $N(\Omega \setminus D) \cap D = (\Omega \setminus D) \cap D$ is the empty set. Since $A(\Omega \setminus D) = ab/4$ then $A(D) = ab/2 - ab/4 = ab/4$. By Proposition 3.5, we have for $\bar{x} \in D$

$$\begin{aligned} \bar{x} \in Q(D) &\iff \frac{1}{ab/4} \left[\frac{ab}{6} \frac{1}{(\bar{x}_1/a + \bar{x}_2/b)} - \frac{ab}{2} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + \frac{ab}{3} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 \right] = 1 \iff \\ &2 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^3 - 3 \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right)^2 - \frac{3}{2} \left(\frac{\bar{x}_1}{a} + \frac{\bar{x}_2}{b} \right) + 1 = 0. \end{aligned}$$

So,

$$Q(D) = D \cap \{ \bar{x} \in \mathbb{R}_{++}^2 : \bar{x}_1/a + \bar{x}_2/b = p \} =$$

$$\{ \bar{x} \in \mathbb{R}_{++}^2 : \bar{x}_1 \geq a/2, \bar{x}_1/a + \bar{x}_2/b = p \} \cup \{ \bar{x} \in \mathbb{R}_{++}^2 : \bar{x}_2 \geq b/2, \bar{x}_1/a + \bar{x}_2/b = p \},$$

where $2p^3 - 3p^2 - (3/2)p + 1 = 0$, $0 < p \leq 1$.

The following two examples were considered in [1] for ICAR functions defined on \mathbb{R}_+^2 . Note that the coefficient k plays here the same role as the number $(1/3)$ in [1].

Example 4.5 Consider the triangle D with vertices $(0,0)$, $(a,0)$ and (a,va) :

$$D = \{ x \in \mathbb{R}_{++}^2 : x_1 \leq a, x_2 \leq vx_1 \}.$$

If $\bar{x} \in D$ then

$$D_1(\bar{x}) = \{ x \in \mathbb{R}_{++}^2 : \bar{x}_1 \leq x_1 \leq a, \bar{x}_2 \leq x_2 \leq (\bar{x}_2/\bar{x}_1)x_1 \},$$

$$D_2(\bar{x}) = \{ x \in \mathbb{R}_{++}^2 : \bar{x}_1 \leq x_1 \leq a, (\bar{x}_2/\bar{x}_1)x_1 \leq x_2 \leq vx_1 \}.$$

Calculation gives the following quantities

$$\frac{1}{\bar{x}_2} \int_{D_1(\bar{x})} x_2 dx_1 dx_2 = \frac{1}{\bar{x}_2} \int_{\bar{x}_1}^a dx_1 \int_{\bar{x}_2}^{(\bar{x}_2/\bar{x}_1)x_1} x_2 dx_2 = \bar{x}_2 \left(\frac{a^3}{6\bar{x}_1^2} - \frac{a}{2} + \frac{\bar{x}_1}{3} \right),$$

$$\frac{1}{\bar{x}_1} \int_{D_2(\bar{x})} x_1 dx_1 dx_2 = \frac{1}{\bar{x}_1} \int_{\bar{x}_1}^a dx_1 \int_{(\bar{x}_2/\bar{x}_1)x_1}^{vx_1} x_1 dx_2 = \left(\frac{va^3}{3\bar{x}_1} - \frac{v\bar{x}_1^2}{3} \right) - \bar{x}_2 \left(\frac{a^3}{3\bar{x}_1^2} - \frac{\bar{x}_1}{3} \right).$$

Further,

$$\int_D \varphi \left(\frac{1}{\bar{x}}, x \right) dx = \left(\frac{va^3}{3\bar{x}_1} - \frac{v\bar{x}_1^2}{3} \right) + \bar{x}_2 \left(\frac{2\bar{x}_1}{3} - \frac{a}{2} - \frac{a^3}{6\bar{x}_1^2} \right).$$

Since $A(D) = va^2/2$ then a point $\bar{x} \in D$ belongs to $Q(D)$ if and only if

$$\left(\frac{2}{3} \frac{a}{\bar{x}_1} - \frac{2}{3} \frac{\bar{x}_1^2}{a^2} \right) + \frac{\bar{x}_2}{va} \left(\frac{4}{3} \frac{\bar{x}_1}{a} - 1 - \frac{1}{3} \frac{a^2}{\bar{x}_1^2} \right) = 1 \iff$$

$$\bar{x}_2 \left(1 + 3 \frac{\bar{x}_1^2}{a^2} - 4 \frac{\bar{x}_1^3}{a^3} \right) = v\bar{x}_1 \left(2 - 3 \frac{\bar{x}_1}{a} - 2 \frac{\bar{x}_1^3}{a^3} \right).$$

In particular, if $\bar{x}_2 = v\bar{x}_1$ then we get the equation $2(\bar{x}_1/a)^3 - 3(\bar{x}_1/a)^2 - 3(\bar{x}_1/a) + 1 = 0$, hence $(\bar{x}_1/a) = k$. So, the point (ka, vka) belongs to $Q(D)$. This implies that for each InR function f , which is integrable on D :

$$f(ka, vka) \leq \frac{1}{A(D)} \int_D f(x) dx.$$

If $\bar{x}_2 = v\bar{x}_1/2$ then equation has the form $(\bar{x}_1/a)^2 + 2(\bar{x}_1/a) - 1 = 0$. This shows that $(\bar{x}_1/a) = \sqrt{2} - 1$, therefore $((\sqrt{2} - 1)a, v(\sqrt{2} - 1)a/2) \in Q(D)$.

Further, we may set in (14) $\bar{x} = (a, va)$:

$$\begin{aligned} \int_D f(x) dx &\leq f(a, va) \int_D \max \left\{ \frac{x_1}{a}, \frac{x_2}{va} \right\} dx_1 dx_2 = f(a, va) \int_D \frac{x_1}{a} dx_1 dx_2 \\ &= \frac{f(a, va)}{a} \int_0^a dx_1 \int_0^{vx_1} x_1 dx_2 = \frac{va^2}{3} f(a, va). \end{aligned}$$

Thus,

$$\frac{1}{A(D)} \int_D f(x) dx \leq \frac{2}{3} f(a, va).$$

Example 4.6 Let D be the square:

$$D = \{x \in \mathbb{R}_{++}^2 : x_1 \leq 1, x_2 \leq 1\}.$$

We consider two possible cases for $\bar{x} \in D$: $(\bar{x}_2/\bar{x}_1) \leq 1$ and $(\bar{x}_2/\bar{x}_1) \geq 1$.

a.) If $(\bar{x}_2/\bar{x}_1) \leq 1$ then we have

$$\begin{aligned} \frac{1}{\bar{x}_2} \int_{D_1(\bar{x})} x_2 dx_1 dx_2 &= \frac{1}{\bar{x}_2} \int_{\bar{x}_1}^1 dx_1 \int_{\bar{x}_2}^{(\bar{x}_2/\bar{x}_1)x_1} x_2 dx_2 = \frac{\bar{x}_2}{2} \left(\frac{1}{3\bar{x}_1^2} - 1 + \frac{2\bar{x}_1}{3} \right), \\ \frac{1}{\bar{x}_1} \int_{D_2(\bar{x})} x_1 dx_1 dx_2 &= \frac{1}{\bar{x}_1} \int_{\bar{x}_1}^1 dx_1 \int_{(\bar{x}_2/\bar{x}_1)x_1}^1 x_1 dx_2 = \frac{1}{2} \left(\frac{1}{\bar{x}_1} - \bar{x}_1 \right) + \frac{\bar{x}_2}{3} \left(\bar{x}_1 - \frac{1}{\bar{x}_1^2} \right). \end{aligned}$$

Hence

$$\int_D \varphi \left(\frac{1}{\bar{x}}, x \right) dx = \frac{1}{2} \left(\frac{1}{\bar{x}_1} - \bar{x}_1 \right) + \frac{\bar{x}_2}{6} \left(4\bar{x}_1 - 3 - \frac{1}{\bar{x}_1^2} \right).$$

Since $A(D) = 1$ then we get the equation for $\bar{x} \in Q(D)$

$$\begin{aligned} \frac{1}{2} \left(\frac{1}{\bar{x}_1} - \bar{x}_1 \right) + \frac{\bar{x}_2}{6} \left(4\bar{x}_1 - 3 - \frac{1}{\bar{x}_1^2} \right) &= 1 \iff \\ \bar{x}_2 \left(1 + 3\bar{x}_1^2 - 4\bar{x}_1^3 \right) &= 3\bar{x}_1 \left(1 - 2\bar{x}_1 - \bar{x}_1^2 \right). \end{aligned}$$

b.) If $(\bar{x}_2/\bar{x}_1) \geq 1$ then we get the symmetric equation

$$\bar{x}_1 \left(1 + 3\bar{x}_2^2 - 4\bar{x}_2^3 \right) = 3\bar{x}_2 \left(1 - 2\bar{x}_2 - \bar{x}_2^2 \right).$$

So, the set $Q(D)$ can be represented as the union of two sets:

$$\left\{ \bar{x} \in \mathbb{R}_{++}^2 : \bar{x}_2 \leq \bar{x}_1 \leq 1, \bar{x}_2 \left(1 + 3\bar{x}_1^2 - 4\bar{x}_1^3 \right) = 3\bar{x}_1 \left(1 - 2\bar{x}_1 - \bar{x}_1^2 \right) \right\}$$

and

$$\left\{ \bar{x} \in \mathbb{R}_{++}^2 : \bar{x}_1 \leq \bar{x}_2 \leq 1, \bar{x}_1 \left(1 + 3\bar{x}_2^2 - 4\bar{x}_2^3 \right) = 3\bar{x}_2 \left(1 - 2\bar{x}_2 - \bar{x}_2^2 \right) \right\}.$$

In particular, if $\bar{x}_1 = \bar{x}_2$ then

$$\begin{aligned} \bar{x} \in Q(D) &\iff \left(0 < \bar{x}_1 \leq 1, \left(1 + 3\bar{x}_1^2 - 4\bar{x}_1^3 \right) = 3 \left(1 - 2\bar{x}_1 - \bar{x}_1^2 \right) \right) \\ &\iff \left(0 < \bar{x}_1 \leq 1, 2\bar{x}_1^3 - 3\bar{x}_1^2 - 3\bar{x}_1 + 1 = 0 \right). \end{aligned}$$

This implies that $(k, k) \in Q(D)$.

At last we investigate inequality (14) with $\bar{x} = (1, 1)$ for the square D :

$$\int_D f(x) dx \leq f(1, 1) \int_D \max\{x_1, x_2\} dx_1 dx_2.$$

Since $A(D) = 1$ and

$$\begin{aligned} \int_D \max\{x_1, x_2\} dx_1 dx_2 &= \int_0^1 dx_1 \int_0^{x_1} x_1 dx_2 + \int_0^1 dx_1 \int_{x_1}^1 x_2 dx_2 \\ &= \frac{1}{3} + \int_0^1 \frac{(1-x_1^2)}{2} dx_1 = \frac{1}{3} + \frac{1}{2} - \frac{1}{6} = \frac{2}{3} \end{aligned}$$

then

$$\frac{1}{A(D)} \int_D f(x) dx \leq \frac{2}{3} f(1, 1),$$

and this estimate holds for every increasing radiant and integrable on D function f .

Acknowledgement

The author is very grateful to A. M. Rubinov for formulation of the tasks and very useful discussions.

References

- [1] S.S. Dragomir, J. Dutta and A.M. Rubinov. *Hermite-Hadamard-type inequalities for increasing convex-along-rays functions*, (submitted to *Real Analysis Exchange*).
- [2] A.M. Rubinov (2000). *Abstract convexity and global optimization*. Kluwer Academic Publishers, Boston-Dordrecht-London.
- [3] A.M. Rubinov and B.M. Glover (1998). Duality for increasing positively homogeneous functions and normal sets. *RAIRO-Operations Research*, **32**, 105-123.
- [4] E.V. Sharikov. *Increasing radiant functions*, (submitted to *Optimization*).