

**A NECESSARY OPTIMALITY CONDITION FOR  
QUASICONVEX FUNCTIONS ON CONVEX SETS  
DEFINED BY INEQUALITY CONSTRAINTS**

YOUSSEF JABRI AND ABDESSAMAD JADDAR

University Mohamed I, Department of Mathematics, Oujda, Morocco

ABSTRACT. We give a necessary optimality condition for the minima of quasiconvex functions on closed convex sets using subdifferentials and normal cones. We consider both the case of a general convex set and a convex set defined as a constrained set for a quasiconvex l.s.c. function.

INTRODUCTION

Consider the following problem

$$(\mathcal{P}) \quad \begin{cases} \text{minimize } f(x), \\ x \in C \subset X, \end{cases}$$

where  $X$  is a Banach space,  $f: X \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $C$  is closed convex.

When looking in the literature for the nature of the different conditions on the objective function  $f$ , to solve  $(\mathcal{P})$ , we see clearly that convexity and differentiability are among the widely used candidates. Nevertheless, even the case when  $f$  is neither convex nor differentiable has been treated. In [1], Clarke considered the case of locally Lipschitz functions and in [2], Huriart-Urruty the case of directionally stable functions. The two papers may be considered as contributions to the case where  $f$  enjoys some “regularity.”

A natural question is the following: what happens when  $f$  is less regular, but instead possesses some kind of convexity?

In this paper, we consider the case of quasiconvex functions. The case of pseudoconvex functions is treated in the paper [3].

The paper is organized as follows. After recalling basic definitions and properties, we give in the next section a necessary condition for a minimization problem of a quasiconvex function on a closed convex set. We consider both the case of a general convex set and a convex set defined as a constrained set for a quasiconvex l.s.c. function.

As usual,  $X^*$  denotes the dual space to  $X$  and  $\langle \cdot, \cdot \rangle$  the duality pairing. The interval  $[a, b] = \{a + t(b - a); 0 \leq t \leq 1\}$  and  $]a, b[ = [a, b] \setminus \{a, b\}$ . The open ball

---

*Key words and phrases.* minimization, quasiconvexity, normal cone, subdifferential.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

centered at  $x$  with radius  $r$  is denoted by  $B_r(x)$ . We recall that  $f$  is quasiconvex if for any  $x, y \in X$  and any  $z \in [x, y]$ ,

$$f(z) \leq \max\{f(x), f(y)\}.$$

This is equivalent to the convexity of the level sets

$$S_\lambda(f) = \{x \in X; f(x) \leq \lambda\}, \quad \forall \lambda \in \mathbb{R}.$$

We will also use the notations  $\tilde{S}_\lambda(f) = \{x \in X; f(x) < \lambda\}$ ,  $L_f(x_0) = \{x \in X; f(x) = f(x_0)\}$ . The mapping  $f$  is lower semicontinuous (l.s.c.) if  $S_\lambda(f)$  is closed for any  $\lambda \in \mathbb{R}$ . When  $f$  is l.s.c., the Clarke-Rockafellar generalized derivative at  $x$  along the direction  $v$  is defined by

$$f^\nearrow(x; v) = \sup_{\varepsilon > 0} \limsup_{\substack{y \rightarrow_f x \\ t \searrow 0}} \inf_{u \in B_\varepsilon(v)} \frac{f(y + tu) - f(y)}{t},$$

where  $y \rightarrow_f x$  means that  $y \rightarrow x$  and  $f(y) \rightarrow f(x)$ . The Clarke-Rockafellar subdifferential of  $f$  at  $x$  is

$$\partial f(x) = \{x^* \in X^*; \langle x^*, v \rangle \leq f^\nearrow(x; v), \forall v \in X\}$$

with the convention that  $\partial f(x)$  is empty if  $f$  is not finite at  $x$ . And last, the normal cone of  $f$  to the convex set  $C$  at  $x_0$  is defined by

$$N(C; x_0) = \{x^* \in X^*; \langle x^*, x - x_0 \rangle \leq 0, \forall x \in C\}.$$

#### MINIMIZATION OF QUASICONVEX FUNCTIONS

The main result in this note is a necessary optimality condition for  $(\mathcal{P})$  when  $f$  is l.s.c., quasiconvex and  $C$  is any nonempty closed convex set of  $X$ .

**Theorem 1.** *Let  $X$  be a Banach space,  $f: X \rightarrow \mathbb{R} \cup \{+\infty\}$  a l.s.c. quasiconvex function. Consider  $x_0 \in C$  such that*

- (i)  $\tilde{S}_{f(x_0)}(f)$  is nonempty and open in  $X$ .
- (ii)  $\partial f(x_0)$  is nonempty and  $w^*$ -compact in  $X^*$ .
- (iii) There is  $\nu > 0$  such that

$$\forall x \in B_\nu(x_0) \cap L(x_0), \quad 0 \notin \partial f(x). \quad (1)$$

*Then, a necessary condition for  $x_0$  to be a solution of  $(\mathcal{P})$  is that*

$$0 \in \partial f(x_0) + N(C; x_0). \quad (2)$$

We will need in the sequel the following technical result.

**Lemma 1.** *Let  $X$  be a Banach space,  $f: X \rightarrow \mathbb{R} \cup \{+\infty\}$  a l.s.c. quasiconvex function.*

(i) *If  $\partial f(x_0)$  is nonempty and there exists  $r > 0$  such that  $0 \notin \partial f(x)$  for all  $x \in B_r(x_0) \cap L_f(x_0)$ , then*

$$N(S_{f(x_0)}(f); x_0) = Cl(\mathbb{R}^+ \partial f(x_0)).$$

(ii) *If moreover,  $\partial f(x_0)$  is  $w^*$ -compact, then*

$$N(S_{f(x_0)}(f); x_0) = \mathbb{R}^+ \partial f(x_0).$$

PROOF.

The point (i) is [Proposition 2.2, 4].

(ii) Consider a sequence  $(\lambda_n x_n^*)_n \subset \mathbb{R}^+ \partial f(x_n)$  such that  $\lambda_n x_n^* \rightharpoonup y^*$ . We will show that  $y^* = \lambda x^*$  for some  $\lambda \in \mathbb{R}^+$  and  $x^* \in \partial f(x_0)$ .

Since  $x_n^* \in \partial f(x_0)$  which is  $w^*$ -compact, for a subsequence still denoted  $(x_n^*)_n$ ,  $x_n^* \rightharpoonup x^* \in \partial f(x_0)$ .

**Claim 1.** *There is a subsequence of  $(\lambda_n)_n$ , still denoted  $(\lambda_n)_n$ , that is bounded.*

Indeed,  $0 \notin \partial f(x_0)$ . By the Hahn-Banach theorem, there is  $v \in X$  such that

$$\langle z^*, v \rangle > 0, \quad \forall z^* \in \partial f(x_0). \quad (3)$$

But  $\lambda_n x_n^* \rightharpoonup y^*$ , so there is  $M > 0$  such that

$$M \geq \langle \lambda_n x_n^*, v \rangle.$$

If  $(\lambda_n)_n$  was not bounded, for some subsequence, still denoted  $(\lambda_n)_n$ , we would get

$$\frac{M}{\lambda_n} \geq \langle x_n^*, v \rangle > 0.$$

At the limit, we get a contradiction with (3).  $\square$

PROOF OF THEOREM 1.

Suppose that  $x_0$  minimizes  $f$  on  $C$ . Then,  $C \cap \tilde{S}_{f(x_0)}(f) = \emptyset$ . But  $\tilde{S}_{f(x_0)}(f) \cap B_{\nu/4}(x_0) \neq \emptyset$  because otherwise,  $x_0$  would be a local minimum of  $f$  and hence we would get  $0 \in \partial f(x_0)$ , a contradiction with (iii). Moreover,

$$(C \cap Cl(B_{\nu/2}(x_0))) \cap (\tilde{S}_{f(x_0)}(f) \cap B_{\nu/2}(x_0)) = \emptyset.$$

By (i) and using the Hahn-Banach theorem, there is  $u^* \in X^*$  such that  $u^* \neq 0$  and  $\alpha \in \mathbb{R}$  separating our two convex sets:

$$\langle u^*, x \rangle \leq \alpha, \quad \forall x \in \tilde{S}_{f(x_0)}(f) \cap B_{\nu/2}(x_0), \quad (4)$$

$$\langle u^*, x \rangle \geq \alpha, \quad \forall x \in C \cap B_{\nu/2}(x_0), \quad (5)$$

We claim that  $\langle u^*, x_0 \rangle = \alpha$ . Indeed, it is clear that  $\langle u^*, x_0 \rangle \geq \alpha$ . It suffices to check the other sense.

Let us first show the equality

$$Cl(\tilde{S}_{f(x_0)}(f)) \cap B_{\nu/2}(x_0) = S_{f(x_0)}(f) \cap B_{\nu/2}(x_0). \quad (6)$$

Indeed, the sense “ $\subset$ ” is obvious. For the inverse inclusion, suppose by contradiction that there is  $y \in \left( Cl(\tilde{S}_{f(x_0)}(f)) \cap B_{\nu/2}(x_0) \right) \setminus (S_{f(x_0)}(f) \cap B_{\nu/2}(x_0))$ . Then,  $y \in L_f(x_0) \cap B_{\nu/2}(x_0)$  and it is a local minimum of  $f$ . So  $0 \in \partial f(y)$ , a contradiction with (iii).

By (6), there is a sequence  $(x_n)_n \subset \tilde{S}_{f(x_0)}(f) \cap B_{\nu/2}(x_0)$  such that  $x_n \rightarrow x_0$  and hence  $\langle u^*, x_0 \rangle \leq \alpha$ .

Using (4),  $\langle u^*, x_0 \rangle = \alpha$  and (6), we get

$$\begin{aligned} u^* \in N(\tilde{S}_{f(x_0)}(f) \cap B_{\nu/2}(x_0); x_0) &= && \text{(property of normal cones)} \\ N(Cl(\tilde{S}_{f(x_0)}(f)) \cap B_{\nu/2}(x_0); x_0) &= && \text{(by (6))} \\ N(S_{f(x_0)}(f) \cap B_{\nu/2}(x_0); x_0) &= && \text{(property of normal cones)} \\ N(S_{f(x_0)}(f); x_0) \end{aligned}$$

By (ii) of Lemma 1, we have

$$u^* \in \mathbb{R}^+ \partial f(x_0).$$

And by (5),

$$-u^* \in N(C; x_0).$$

Since  $u^* \neq 0$ , we finally get

$$0 \in \partial f(x_0) + N(C; x_0).$$

□

This theorem refines the results of Clarke [1] and Huriart-Urruty [2] when we require the quasiconvexity of the objective function  $f$ .

In the case where the general convex set  $C$ , appearing in the former result, is defined as the constraint set

$$C = \{x \in X; g(x) \leq 0\},$$

where  $g$  is quasiconvex, l.s.c. and satisfies (ii), (iii) of Theorem 1, and  $g(x_0) = 0$ , we obtain the following result where appears some Lagrange multiplier.

**Corollary 1.** *A necessary condition for  $x_0$  to solve (P) is*

$$0 \in \partial f(x_0) + \lambda \partial g(x_0), \quad \text{for some } \lambda > 0.$$

Note that  $g(x_0) = 0$  is not a problem, we could always use  $h(t) = g(x) - g(x_0)$ . For the proof, it suffices to use Theorem 1 and Lemma 1(ii).

The case where  $f$  is pseudoconvex is investigated in an other paper [3].

#### REFERENCES

1. F.H. Clarke, *Optimization and nonsmooth analysis*, Wiley-Interscience, New York, 1983.
2. J.B. Huriart-Urruty, *Tangent cones, generalized gradients and mathematical programming in Banach spaces*, Math. Oper. Res. **4** (1979), 79–97.
3. Y. Jabri and A. Jaddar., *Characterization of minima of pseudoconvex functions on closed convex sets*, Preprint (2002).
4. A. Hassouni and A. Jaddar., *Quasiconvex functions and applications to optimality conditions in nonlinear programming*, Applied Mathematics Letters **14** (2001), 241–244.