

# A NOTE ON OSTROWSKI'S INEQUALITY

CONSTANTIN P. NICULESCU AND AURELIA FLOREA

ABSTRACT. This paper deals with the problem of estimating the deviation of the values of a function from its mean value. We consider the following special cases: i) the case of random variables (attached to arbitrary probability fields); ii) the comparison is performed additively or multiplicatively; iii) the mean value is attached to a multiplicative averaging process.

## 1. INTRODUCTION

The inequality of Ostrowski [10] gives us an estimate for the deviation of the values of a smooth function from its mean value. More precisely, if  $f : [a, b] \rightarrow \mathbb{R}$  is a differentiable function with bounded derivative, then

$$(O) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{(x - (a+b)/2)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for every  $x \in [a, b]$ . Moreover the constant  $1/4$  is the best possible.

The proof is an application of Lagrangian's mean value theorem:

$$\begin{aligned} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| &= \left| \frac{1}{b-a} \int_a^b (f(x) - f(t)) dt \right| \\ &\leq \frac{1}{b-a} \int_a^b |f(x) - f(t)| dt \\ &\leq \frac{\|f'\|_\infty}{b-a} \int_a^b |x-t| dt \\ &= \left[ \frac{(x-a)^2 + (b-x)^2}{2(b-a)} \right] \|f'\|_\infty \\ &= \left[ \frac{1}{4} + \left( \frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty. \end{aligned}$$

The optimality of the constant  $1/4$  is also immediate, checking the inequality for the family of functions  $f_\alpha(t) = |x-t|^\alpha \cdot (b-a)$  ( $t \in [a, b]$ ,  $\alpha > 1$ ) and then passing to the limit as  $\alpha \rightarrow 1+$ .

---

2000 *Mathematics Subject Classification.* Primary 26D15; Secondary 26B30, 26E60.

*Key words and phrases.* Lipschitz mapping, mean value, multiplicative mean.

Partially supported by CNCSIS Grant 3/2002.

Presented to the 6th *Conference on Mathematical Analysis and its Applications*, Cluj-Napoca, November 8-9, 2002.

It is worth to notice that the smoothness condition can be relaxed. In fact, the Lipschitz class suffices as well, by replacing  $\|f'\|_\infty$  with the Lipschitz constant of  $f$ , i.e.,

$$\|f\|_L = \sup_{x \neq y} \left| \frac{f(x) - f(y)}{x - y} \right|.$$

The extension to the context of vector-valued functions, with values in a Banach space, is straightforward.

Since a Lipschitz function on  $[a, b]$  is absolutely continuous, a natural direction of generalization of the Ostrowski inequality was its investigation within this larger class of functions (with refinements for  $f' \in L^p([a, b])$ ,  $1 \leq p < \infty$ ). See Fink [5]. Also, several Ostrowski type inequalities are known within the framework of Hölder functions as well as for functions of bounded variation.

The problem to estimate the deviation of a function from its mean value can be investigated from many other points of view:

- by considering random variables (attached to arbitrary probability fields);
- by changing the algebraic nature of the comparison (e.g., switching to the multiplicative framework);
- by considering other means (for example, the geometric mean);
- by estimating the deviation via other norms (the classical case refers to the sup norm, but  $L^p$ -norms are better motivated in other situations).

The aim of this paper is to present a number of examples giving support to this program.

## 2. OSTROWSKI TYPE INEQUALITIES FOR RANDOM VARIABLES

In what follows  $X$  will denote a locally compact metric space and  $E$  a Banach space.

**Theorem 1.** *The following two assertions are equivalent for  $f : X \rightarrow E$  a continuous mapping:*

- i)  $f$  is Lipschitz i.e.,  $\|f\|_L = \sup_{x \neq y} \frac{\|f(x) - f(y)\|}{d(x, y)} < \infty$ ;*
- ii) For every  $x \in X$  and every Borel probability measure  $\mu$  on  $X$  such that  $f \in \mathcal{L}^1(\mu)$  we have*

$$\left\| f(x) - \int_X f \, d\mu \right\| \leq \|f\|_L \int_X^* d(x, y) \, d\mu.$$

Here  $*$  marks the upper integral.

*Proof.* *i)  $\Rightarrow$  ii).* As  $d(x, \cdot)$  is continuous it is also Borel measurable, so being nonnegative its upper integral is perfectly motivated. Then we can proceed as in the classical case, described in the Introduction.

*ii)  $\Rightarrow$  i).* Consider the particular case of Dirac measure  $\delta_y$  (concentrated at  $y$ ). Then

$$\|f(x) - f(y)\| \leq \|f\|_L d(x, y)$$

which shows that  $f$  must be Lipschitz. ■

If  $X$  is a bounded metric space, the above theorem works for all continuous mappings. In fact, if  $\|f\|_L < \infty$  then  $f$  is necessarily bounded (and thus it belongs to  $\mathcal{L}^\infty(\mu) \subset \mathcal{L}^1(\mu)$ ). Also, the mappings  $d(x, \cdot)$  are  $\mu$ -integrable (which makes  $*$  unnecessary).

The condition  $f \in \mathcal{L}^1(\mu)$  is automatically fulfilled by all continuous bounded functions regardless what Borel probability measure  $\mu$  we consider on  $X$ ; in fact, they are in  $\mathcal{L}^\infty(\mu) \subset \mathcal{L}^1(\mu)$ . In general, not every continuous function  $f$  is  $\mu$ -integrable. For, think at the case where  $X = \mathbb{R}$ ,  $f(x) = x$ , and  $\mu = \frac{1}{\pi(1+x^2)} dx$ .

We shall illustrate Theorem 1 in a number of particular situations. The first one, concerns the case of classical probability fields:

**Corollary 1.** *Let  $E$  be a normed vector space and let  $x_1, \dots, x_n$  be  $n$  vectors in  $E$ . Then*

$$\left\| x_i - \frac{1}{n} \sum_{k=1}^n x_k \right\| \leq \frac{1}{n} \left[ \left( i - \frac{n+1}{2} \right)^2 + \frac{n^2-1}{4} \right] \cdot \sup_{1 \leq k \leq n-1} \|x_{k+1} - x_k\|.$$

*Proof.* We consider the measure space  $(X, \Sigma, \mu)$ , where  $X = \{1, \dots, n\}$ ,  $\Sigma = \mathcal{P}(X)$  and  $\mu(A) = |A|$  for every  $A \subset X$ .

$X$  has a natural structure of metric subspace of  $\mathbb{R}$ . The function

$$f : X \rightarrow E, \quad f(i) = x_i,$$

is Lipschitz, with Lipschitz constant

$$(L) \quad L = \sup_{1 \leq k \leq n-1} \|x_{k+1} - x_k\|.$$

In fact, if  $i < j$ ,

$$\begin{aligned} \|f(i) - f(j)\| &= \|x_i - x_j\| \\ &\leq \|x_i - x_{i+1}\| + \dots + \|x_{j-1} - x_j\| \\ &\leq (j-i) \cdot \sup_{1 \leq k \leq n-1} \|x_{k+1} - x_k\|, \end{aligned}$$

which proves the inequality  $\leq$  in (L). The other inequality is clear.

According to Theorem 1,

$$\left\| f(i) - \frac{1}{\mu(X)} \int_X f(k) d\mu(k) \right\| \leq \frac{L}{\mu(X)} \int_X |i-k| d\mu(k),$$

which can be easily shown to be equivalent to the inequality in the statement of Corollary 1 because

$$\int_X |i-k| d\mu(k) = \sum_{k=1}^n |i-k| = \left( i - \frac{n+1}{2} \right)^2 + \frac{n^2-1}{4}. \quad \blacksquare$$

Notice that the right hand side of the inequality in Corollary 1 is  $\geq \sqrt{\text{var}(f)}$ , where

$$\text{var}(f) = \frac{1}{\mu(X)} \int_X \left| f(x) - \frac{1}{\mu(X)} \int_X f(t) d\mu(t) \right|^2 d\mu(x)$$

represents the variance of  $f$ . According to the classical Chebyshev inequality,

$$\mu \left\{ \left| f - \frac{1}{\mu(X)} \int_X f d\mu \right| \leq \varepsilon \right\} \geq 1 - \frac{\text{var } f}{\varepsilon^2}$$

and the discussion above shows that the range of interest in this inequality is precisely

$$\text{var } f < \varepsilon \leq \frac{1}{n} \left[ \left( i - \frac{n+1}{2} \right)^2 + \frac{n^2-1}{4} \right] \cdot \sup_{1 \leq k \leq n-1} \|x_{k+1} - x_k\|.$$

As well known, convolution by smooth kernels leads to good approximation schemes. Theorem 1 allows us to estimate the speed of convergence. Here is an example:

**Corollary 2.** *Let  $f : \mathbb{R} \rightarrow E$  be a Lipschitz mapping. Then*

$$\left\| f(x) - \frac{n}{(2\pi)^{1/2}} \int_{\mathbb{R}} f(t) e^{-n^2(x-t)^2/2} dt \right\| \leq \frac{2 \|f\|_L}{n(2\pi)^{1/2}}$$

for every  $x \in \mathbb{R}$ . Particularly,  $f$  is the uniform limit of a sequence  $(f_n)_n$  of Lipschitz functions of class  $C^\infty$ , with  $\|f_n\|_L \leq \|f\|_L$  for every  $n$ .

We end this section with the case of functions of several variables:

**Corollary 3.** *Let  $f = f(x, y)$  be a differentiable function defined on a compact 2-dimensional interval  $X = [a, b] \times [c, d]$  such that  $|\partial f / \partial x| \leq L$  and  $|\partial f / \partial y| \leq M$ . Then*

$$\begin{aligned} \left\| f(x, y) - \frac{1}{\text{Area } X} \iint_X f(u, v) du dv \right\| &\leq L \left[ \frac{1}{4} + \left( \frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] \text{Area } X + \\ &+ M \left[ \frac{1}{4} + \left( \frac{y - \frac{c+d}{2}}{d-c} \right)^2 \right] \text{Area } X. \end{aligned}$$

*Proof.* Clearly, we may assume that  $L, M > 0$ . Then  $f$  is a Lipschitz function (with Lipschitz constant 1) provided that  $X$  is endowed with the metric

$$d((x, y), (u, v)) = L|x - u| + M|y - v|.$$

Now the conclusion follows from Theorem 1. ■

### 3. COMPARISON OF ARITHMETIC MEANS

Suppose that  $X$  is a locally compact bounded metric space on which there are given two Borel probability measures  $\mu$  and  $\nu$ . We are interested to estimate the difference

$$\int_X f d\mu - \int_X f d\nu$$

for  $f$  a Lipschitz function on  $X$ , with values in a Banach space  $E$ . For  $\nu = \delta_x$ , this reduces to the classical Ostrowski inequality.

Following the ideas in the preceding section we are led to

$$\left\| \int_X f d\mu - \int_X f d\nu \right\| \leq \|f\|_L \int_X \int_X d(x, y) d\mu(x) d\nu(y)$$

but this is not always the best result.

For example, for  $\mu = dx/(b-a)$  and  $\nu = (\delta_a + \delta_b)/2$  (on  $X = [a, b]$ ) it yields the trapezoid inequality

$$\left\| \frac{1}{b-a} \int_a^b f(x) dx - \frac{f(a) + f(b)}{2} \right\| \leq C \|f\|_L (b-a),$$

with  $C = 1/2$ . This can be improved up to  $C = 1/4$  within Ostrowski theory (by applying (O) to  $f|[a, (a+b)/2]$  and  $f|[(a+b)/2, b]$ , for  $x = a$  and  $x = b$  respectively).

However, the Iyengar inequality gives us a better upper bound:

$$(Iy) \quad \left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{f(a)+f(b)}{2} \right| \leq \frac{\|f\|_L}{4}(b-a) - \frac{(f(b)-f(a))^2}{4(b-a)\|f\|_L}.$$

See [6], [7], or [9] for details. We can combine (O) and (Iy) to get a more general result:

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x) dx - \left[ \lambda f\left(\frac{a+b}{2}\right) + (1-\lambda) \frac{f(a)+f(b)}{2} \right] \right| \\ & \leq \frac{\|f\|_L}{4}(b-a) - (1-\lambda) \frac{(f(b)-f(a))^2}{4(b-a)\|f\|_L}, \end{aligned}$$

for every Lipschitz function  $f$  and every  $\lambda \in [0, 1]$ . This is optimal in the Lipschitz class, but better results are known for smooth functions. See [7].

There is a large activity (motivated by the problems in numerical integration) concerning the approximation of probability measures by convex combinations of Dirac measures. However, sharp general formulas remain to be found.

#### 4. THE MULTIPLICATIVE SETTING

According to [8], the *multiplicative mean value* of a continuous function  $f : [a, b] \rightarrow (0, \infty)$  (where  $0 < a < b$ ) is defined by the formula

$$\begin{aligned} M_*(f) &= \exp\left(\frac{1}{\log b - \log a} \int_{\log a}^{\log b} \log f(e^t) dt\right) \\ &= \exp\left(\frac{1}{\log b - \log a} \int_a^b \log f(t) \frac{dt}{t}\right) \end{aligned}$$

The main properties of the multiplicative mean are listed below:

$$\begin{aligned} M_*(1) &= 1 \\ m \leq f \leq M &\Rightarrow m \leq M_*(f) \leq M \\ M_*(fg) &= M_*(f) M_*(g). \end{aligned}$$

Given a function  $f : I \rightarrow (0, \infty)$  (with  $I \subset (0, \infty)$ ) we shall say that  $f$  is *multiplicatively Lipschitzian* provided there exists a constant  $L > 0$  such that

$$\max\left\{\frac{f(x)}{f(y)}, \frac{f(y)}{f(x)}\right\} \leq \left(\frac{y}{x}\right)^L$$

for all  $x < y$  in  $I$ ; the smallest  $L$  for which the above inequality holds constitutes the *multiplicative Lipschitz constant* of  $f$  and it will be denoted by  $\|f\|_{*Lip}$ .

**Remark 1.** *Though the family of multiplicatively Lipschitz functions is large enough (to deserve attention in its own), we know the exact value of the multiplicative Lipschitz constant only in few cases:*

- i) *If  $f$  is of the form  $f(x) = x^\alpha$ , then  $\|f\|_{*Lip} = \alpha$ .*
- ii) *If  $f = \exp$  on  $[a, b]$  (where  $0 < a < b$ ), then  $\|f\|_{*Lip} = b$ .*
- iii) *Clearly,  $\|f\|_{*Lip} \leq 1$  for every non-decreasing functions  $f$  such that  $f(x)/x$  is non-increasing. For example, this is the case of the functions  $\sin$  and  $\sec$  on  $(0, \pi/2)$ .*

iv) If  $f$  and  $g$  are two multiplicatively Lipschitzian functions (defined on the same interval) and  $\alpha, \beta \in \mathbb{R}$ , then  $f^\alpha g^\beta$  is multiplicatively Lipschitzian too. Moreover,

$$\|f^\alpha g^\beta\|_{*Lip} \leq |\alpha| \cdot \|f\|_{*Lip} + |\beta| \cdot \|g\|_{*Lip}.$$

The following result represents the multiplicative counterpart of the classical Ostrowski inequality:

**Theorem 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a multiplicatively Lipschitz function with  $\|f\|_{*Lip} = L$ . Then

$$\max \left\{ \frac{f(x)}{M_*(f)}, \frac{M_*(f)}{f(x)} \right\} \leq \left( \frac{b}{a} \right)^{L(1/4 + \log^2(x/\sqrt{ab})/\log^2(b/a))}.$$

*Proof.* In fact,

$$\begin{aligned} \frac{M_*(f)}{f(x)} &= \exp \left( \frac{1}{\log b - \log a} \int_a^b \log \frac{f(t)}{f(x)} \frac{dt}{t} \right) \\ &\leq \exp \left( \frac{L}{\log(b/a)} \left( \int_a^x \log \frac{x}{t} \frac{dt}{t} + \int_x^b \log \frac{t}{x} \frac{dt}{t} \right) \right) \\ &= \exp \left( \frac{L}{\log(b/a)} \left( \log x (2 \log x - \log a - \log b) - \log^2 x + \frac{\log^2 a}{2} + \log^2 b \right) \right) \\ &= \exp \left( \frac{L}{\log b/a} \cdot \frac{(\log x - \log a)^2 + (\log b - \log x)^2}{2} \right) \\ &= \left( \frac{b}{a} \right)^{L(1/4 + \log^2(x/\sqrt{ab})/\log^2(b/a))} \end{aligned}$$

and a similar estimate is valid for  $f(x)/M_*(f)$ . ■

For  $f = \exp$  on  $[a, b]$  (where  $0 < a < b$ ), we have  $M_*(f) = \exp\left(\frac{b-a}{\log b - \log a}\right)$  and  $\|f\|_{*Lip} = b$ . By Theorem 3.4, we infer the inequalities

$$\exp \left| \frac{b-a}{\log b - \log a} - x \right| \leq \left( \frac{b}{a} \right)^{b(1/4 + \log^2(x/\sqrt{ab})/\log^2(b/a))}$$

i.e.,

$$\left| \frac{b-a}{\log b - \log a} - x \right| \leq b \left( 1/4 + \frac{\log^2(x/\sqrt{ab})}{(\log b - \log a)^2} \right) (\log b - \log a)$$

for every  $x \in [a, b]$ . Particularly,

$$\left| \frac{b-a}{\log b - \log a} - \sqrt{ab} \right| \leq \frac{b}{4} (\log b - \log a).$$

## 5. AN OPEN PROBLEM

The deviation of the values of a function from its mean value can be estimated via a variety of norms. For example, the Ostrowski inequality yields

$$(O') \quad \left\| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right\|_\infty \leq \frac{b-a}{2} \|f'\|_\infty$$

for every  $f \in C^1([a, b])$ .

In some instances, the  $L^p$ -norms are more suitable. An old result in this direction is the following inequality due to W. Stekloff [12], [13], [7],

$$(S) \quad \left( \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^2 dx \right)^{1/2} \leq \frac{b-a}{\pi} \left( \int_a^b |f'(x)|^2 dx \right)^{1/2}$$

that works also for every  $f \in C^1([a, b])$ . In terms of variance, (S) may be read as

$$\text{var}(f) \leq \frac{b-a}{\pi^2} \int_a^b |f'(t)|^2 dt,$$

so that, combined with the Schwarz inequality, it yields the following estimate for the covariance of two random variables (of class  $C^1$ ):

$$\begin{aligned} \text{cov}(f, g) &\leq \text{var}^{1/2}(f) \cdot \text{var}^{1/2}(g) \\ &\leq \frac{b-a}{\pi^2} \left( \int_a^b |f'(t)|^2 dt \right)^{1/2} \left( \int_a^b |g'(t)|^2 dt \right)^{1/2}. \end{aligned}$$

There is a large literature in this area, including deep results in the higher dimensional case. See [7].

A natural way to pack together (O') and (S) is as follows.

Consider a separable Banach lattice  $E$ . Then  $E$  contains quasi-interior points  $u > 0$  and admits strictly positive functionals  $x' \in E'$ . This means that

$$\lim_{n \rightarrow \infty} \|x - x \wedge nu\| = 0 \text{ for every } x \in E, x \geq 0,$$

and

$$x \geq 0, x'(x) = 0 \text{ implies } x = 0.$$

See H. H. Schaefer [11], for details.

To each such a pair  $(u, x')$  with  $x'(u) = 1$ , one can associate a positive linear projection,

$$M : E \rightarrow E, \quad M(x) = x'(x) \cdot u,$$

whose image is  $\mathbb{R} \cdot u$ . Clearly, this projection provides an analogue of the integral mean.

**Open Problem.** *Characterize all the pairs  $(u, x')$  as above, for which there exist a densely defined linear operator  $D : \text{dom } D \subset E \rightarrow E$  and a positive constant  $C = C(x', u)$  such that*

$$\|x - M(x)\| \leq C \|D(x)\| \text{ for every } x \in \text{dom } D.$$

In the examples at the beginning of this section,  $E$  is one of the spaces  $C([a, b])$  or  $L^2([a, b])$ ,  $u$  is the function identically 1,  $x'$  is the normalized Lebesgue measure and  $D$  is the differential. The same picture follows from Corollary 3 above. The problem is how general could be the existence of a differential like operator in the context of separable Banach lattices.

## REFERENCES

- [1] Lj. Dedić, M. Matić and J. Pečarić, *On some generalizations of Ostrowski inequality for Lipschitz functions and functions of bounded variations*, Math. Inequal. Appl., **3** (2000), 1-14.
- [2] Lj. Dedić, M. Matić and J. Pečarić, *On generalizations of Ostrowski inequality via some Euler-type identities*, Math. Inequal. Appl., **3** (2000), 337-353.
- [3] S. S. Dragomir, *On the Ostrowski inequality for mappings with bounded variation and applications*, RGMIA **2** (1999), No. 1.
- [4] S. S. Dragomir, *The discrete version of Ostrowski inequality in normed linear spaces*, Journal of Inequalities in Pure and Applied Mathematics, **3** (2002), no. 1, article 2.
- [5] A. M. Fink, *Bounds on the deviation of a function from its averages*, Czechoslovak Math. J., **42** (1992), 289-310.
- [6] K. S. K. Iyengar, *Note on an inequality*, Math. Student, **6** (1938), 75-76.
- [7] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, Kluwer Academic, Dordrecht, 1991.
- [8] C. P. Niculescu, *A multiplicative mean value and its applications*. In *Inequality Theory and Applications*, vol. 1, pp. 243-255, Nova Science Publishers, Huntington, New York, 2001, edited by Y. J. Cho, S. S. Dragomir and J. Kim.
- [9] C. P. Niculescu and F. Popovici, *A Note on the Denjoy-Bourbaki Theorem*, Real Analysis Exchange, submitted.
- [10] A. Ostrowski, *Über die Absolutabweichung einer differentiebaren Funktion von ihrem Integralmittelwert*, Comment. Math. Helv. **10** (1938), 226-227.
- [11] H. H. Schaefer, *Banach Lattices and Positive Operators*, Springer Verlag, Berlin, 1974.
- [12] W. Stekloff, *Sur le problème de refroidissement d'une barre hétérogène*, C. R. Sci. Paris **126** (1898), 215-218.
- [13] W. Stekloff, *Problème de refroidissement d'une barre hétérogène*, Ann. Fac. Sci. Toulouse **3** (2) (1901), 281-313.

UNIVERSITY OF CRAIOVA, DEPARTMENT OF MATHEMATICS, CRAIOVA 200585, ROMANIA  
E-mail address: cniculescu@central.ucv.ro  
URL: <http://inf.ucv.ro/~niculescu>

UNIVERSITY OF CRAIOVA, DEPARTMENT OF PHILOLOGY, CRAIOVA 200585, ROMANIA  
E-mail address: aurelia\_florea@yahoo.com