

A TYPE OF MEAN VALUES OF SEVERAL POSITIVE NUMBERS WITH TWO PARAMETERS

ZHEN-GANG XIAO, ZHI-HUA ZHANG, AND FENG QI

ABSTRACT. In this article, a type of mean values of several positive numbers with two parameters are defined by using the generalized Vandermonde determinant, some basic properties of them are given, and several inequalities for some mean values including the generalized symmetric means are established.

1. INTRODUCTION AND NOTATIONS

Let $\mathbb{R} = (-\infty, \infty)$, $\mathbb{R}_+ = (0, \infty)$, $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$, φ be a function defined in \mathbb{R}_+ , and

$$V(\mathbf{a}; \varphi) = \begin{vmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^{n-1} & \varphi(a_0) \\ 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} & \varphi(a_1) \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_n & a_n^2 & \cdots & a_n^{n-1} & \varphi(a_n) \end{vmatrix}. \quad (1.1)$$

Taking $\varphi(x) = x^{n+r}(\ln x)^k$, then

$$V(\mathbf{a}; \varphi) = \begin{vmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^{n-1} & a_0^{n+r}(\ln a_0)^k \\ 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} & a_1^{n+r}(\ln a_1)^k \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_n & a_n^2 & \cdots & a_n^{n-1} & a_n^{n+r}(\ln a_n)^k \end{vmatrix} \triangleq V(\mathbf{a}; r, k). \quad (1.2)$$

If letting $r = 0$ and $k = 0$ in (1.2), then

$$V(\mathbf{a}; 0, 0) = \sum_{i=0}^n (-1)^{n+i} a_i^n V_i(\mathbf{a}) = \prod_{0 \leq i < j \leq n} (a_j - a_i) \quad (1.3)$$

is the determinant of Vandermonde matrix of order $n + 1$, where

$$V_i(\mathbf{a}) = \begin{vmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^{n-1} \\ 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_{i-1} & a_{i-1}^2 & \cdots & a_{i-1}^{n-1} \\ 1 & a_{i+1} & a_{i+1}^2 & \cdots & a_{i+1}^{n-1} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_n & a_n^2 & \cdots & a_n^{n-1} \end{vmatrix} \quad (1.4)$$

2000 *Mathematics Subject Classification.* Primary 26D15.

Key words and phrases. mean values, Vandermonde determinant, inequality, monotonicity, basic property, generalized symmetric mean.

The third author was supported in part by the Science Foundation of Project for Fostering Innovation Talents at Universities of Henan Province, China.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

is the cofactor of the element a_i^n in $V(\mathbf{a}; 0, 0)$. So, we call $V(\mathbf{a}; \varphi)$ the generalized Vandermonde determinant.

Let a_0 and a_1 be two positive numbers. It is well known that the logarithmic mean $L(a_0, a_1)$ and the identric or exponential mean $I(a_0, a_1)$ of a_0 and a_1 are respectively defined by

$$L(a_0, a_1) = \begin{cases} \frac{a_1 - a_0}{\ln a_1 - \ln a_0}, & a_0 \neq a_1, \\ a_0, & a_0 = a_1 \end{cases} \quad (1.5)$$

and

$$I(a_0, a_1) = \begin{cases} \frac{1}{e} \left(\frac{a_1^{a_1}}{a_0^{a_0}} \right)^{\frac{1}{a_1 - a_0}}, & a_0 \neq a_1, \\ a_0, & a_0 = a_1. \end{cases} \quad (1.6)$$

Let $G(a_0, a_1) = \sqrt{a_0 a_1}$ and $A(a_0, a_1) = \frac{a_0 + a_1}{2}$ denote the geometric mean and the arithmetic mean of a_0 and a_1 , respectively. For $a_0 \neq a_1$, the following inequalities are proved in [4]:

$$G(a_0, a_1) < L(a_0, a_1) < I(a_0, a_1) < A(a_0, a_1). \quad (1.7)$$

In [28, 29], Zh.-H. Zhang and Zh.-G. Xiao studied the identric mean and two logarithmic means of $n + 1$ positive numbers a_0, a_1, \dots, a_n which are defined as

$$I(\mathbf{a}) = \exp \left[\frac{V(\mathbf{a}; 0, 1)}{V(\mathbf{a}; 0, 0)} - \sum_{k=1}^n \frac{1}{k} \right], \quad (1.8)$$

$$L(\mathbf{a}) = \frac{V(\mathbf{a}; 0, 0)}{nV(\mathbf{a}; -1, 1)}, \quad (1.9)$$

$$l(\mathbf{a}) = \frac{n!V_{\ln}(\mathbf{a}; 1, 0)}{V_{\ln}(\mathbf{a}; 0, n)}, \quad (1.10)$$

where

$$V_{\ln}(\mathbf{a}; r, k) \triangleq \begin{vmatrix} 1 & \ln a_0 & (\ln a_0)^2 & \cdots & (\ln a_0)^{n-1} & a_0^r (\ln a_0)^k \\ 1 & \ln a_1 & (\ln a_1)^2 & \cdots & (\ln a_1)^{n-1} & a_1^r (\ln a_1)^k \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & \ln a_n & (\ln a_n)^2 & \cdots & (\ln a_n)^{n-1} & a_n^r (\ln a_n)^k \end{vmatrix}. \quad (1.11)$$

They proved for \mathbf{a} with $a_i \neq a_j$ for all $i \neq j$ the following inequalities

$$G(\mathbf{a}) < L(\mathbf{a}) < I(\mathbf{a}) < A(\mathbf{a}) \quad (1.12)$$

and

$$G(\mathbf{a}) < l(\mathbf{a}) < I(\mathbf{a}) < A(\mathbf{a}), \quad (1.13)$$

where $G(\mathbf{a}) = \prod_{i=0}^n a_i^{1/(n+1)}$ and $A(\mathbf{a}) = \frac{1}{n+1} \sum_{i=0}^n a_i$ are the geometric mean and the arithmetic mean of $n + 1$ positive numbers $a_0, a_1, \dots, a_n \in \mathbb{R}_+$, respectively.

For $r \in \mathbb{R} \setminus \{-1, 0\}$, the extended logarithmic mean $S_r(a_0, a_1)$ and the generalized logarithmic mean $J_r(a_0, a_1)$ of two positive numbers a_0 and a_1 are defined (See [1, 2, 10, 13, 15, 24]) respectively by

$$S_r(a_0, a_1) = \begin{cases} \left(\frac{a_1^{r+1} - a_0^{r+1}}{(r+1)(a_1 - a_0)} \right)^{1/r}, & a_0 \neq a_1, \\ a_0, & a_0 = a_1 \end{cases} \quad (1.14)$$

and

$$J_r(a_0, a_1) = \begin{cases} \frac{r}{r+1} \cdot \frac{a_1^{r+1} - a_0^{r+1}}{a_1^r - a_0^r}, & a_0 \neq a_1, \\ a_0, & a_0 = a_1. \end{cases} \quad (1.15)$$

These two means and other mean values of two positive numbers are special cases of the extended mean values

$$E(p, q; a_0, a_1) = \begin{cases} \left[\frac{q(a_1^p - a_0^p)}{p(a_1^q - a_0^q)} \right]^{1/(p-q)} & \text{for } pq(p-q)(a_0 - a_1) \neq 0, \\ \left[\frac{a_1^p - a_0^p}{p(\ln a_1 - \ln a_0)} \right]^{1/p} & \text{for } p(a_0 - a_1) \neq 0 \text{ and } q = 0, \\ \exp \left\{ -\frac{1}{p} + \frac{a_1^p \ln a_1 - a_0^p \ln a_0}{a_1^p - a_0^p} \right\} & \text{for } p(a_0 - a_1) \neq 0 \text{ and } p = q, \\ \sqrt{a_0 a_1} & \text{for } a_0 - a_1 \neq 0 \text{ and } p = q = 0, \\ a_0 & \text{for } p = q \text{ and } a_0 = a_1, \end{cases} \quad (1.16)$$

$$= \begin{cases} \left[\frac{h(q; a_0, a_1)}{h(p; a_0, a_1)} \right]^{1/(q-p)}, & (p-q)(a_0 - a_1) \neq 0, \\ \exp \left[\frac{\partial h(p; a_0, a_1) / \partial p}{h(p; a_0, a_1)} \right], & a_0 - a_1 \neq 0, p = q, \end{cases} \quad (1.17)$$

where

$$h(t; a_0, a_1) = \begin{cases} \frac{a_1^t - a_0^t}{t}, & t \neq 0 \\ \ln a_1 - \ln a_0, & t = 0 \end{cases} = \int_{a_0}^{a_1} u^{t-1} du. \quad (1.18)$$

There have been a lot of literature about the extended mean values $E(p, q; a_0, a_1)$. For more information, please refer to [5, 6, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 25] and the references therein.

In [26, 27], Zh.-G. Xiao and Zh.-H. Zhang introduced

$$S_r(\mathbf{a}) = \begin{cases} \left[\frac{n!}{\prod_{k=1}^n (k+r)} \cdot \frac{V(\mathbf{a}; r, 0)}{V(\mathbf{a}; 0, 0)} \right]^{1/r}, & r \neq 0, -1, \dots, -n, \\ \exp \left[\frac{V(\mathbf{a}; 0, 1)}{V(\mathbf{a}; 0, 0)} - \sum_{k=1}^n \frac{1}{k} \right], & r = 0, \\ \left[\frac{n!V(\mathbf{a}; r, 1)}{(-1)^{r+1}(-r-1)!(n+r)!V(\mathbf{a}; 0, 0)} \right]^{1/r}, & r = -1, \dots, -n \end{cases} \quad (1.19)$$

and

$$J_r(\mathbf{a}) = \begin{cases} \frac{r}{n+r} \cdot \frac{V(\mathbf{a}; r, 0)}{V(\mathbf{a}; r-1, 0)}, & r \neq 0, -1, \dots, -n, \\ \frac{V(\mathbf{a}; 0, 0)}{nV(\mathbf{a}; -1, 1)}, & r = 0, \\ \frac{r}{n+r} \cdot \frac{V(\mathbf{a}; r, 1)}{V(\mathbf{a}; r-1, 1)}, & r = -1, \dots, -(n-1), \\ \frac{-nV(\mathbf{a}; 0, 1)}{V(\mathbf{a}; -n-1, 0)}, & r = -n \end{cases} \quad (1.20)$$

with $S_0(\mathbf{a}) = I(\mathbf{a})$ and $S_{-1}(\mathbf{a}) = J_0(\mathbf{a}) = L(\mathbf{a})$.

The main purpose of this article is to define a type of mean values of several positive numbers with two parameters, from which many mean values mentioned above can be deduced, by utilizing the generalized Vandermonde determinant, to research their basic properties, and to establish several inequalities.

2. LEMMAS

The following two lemmas are keys for us to research the basic properties of the mean values $E(p, q; \mathbf{a})$ defined by Definition 3.1 in next section.

Lemma 2.1. *Let $n \in \mathbb{N}$, $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$ and φ be a n -times differentiable function in \mathbb{R}_+ . Then*

$$V(\mathbf{a}; \varphi) = V(\mathbf{a}; 0, 0) \int_D \varphi^{(n)} \left(\sum_{i=0}^n a_i x_i \right) dx, \quad (2.1)$$

where $x_0 = 1 - \sum_{i=1}^n x_i$ and $dx = dx_1 dx_2 \cdots dx_n$ denotes the differential of the volume in

$$D = \left\{ (x_1, x_2, \dots, x_n) \mid \sum_{i=1}^n x_i \leq 1 \text{ with } x_i \geq 0 \text{ for } i = 1, 2, \dots, n \right\}. \quad (2.2)$$

Proof. Let $\bar{\mathbf{a}} = (a_n, a_1, \dots, a_{n-1}) \in \mathbb{R}_+^n$ and $\tilde{\mathbf{a}} = (a_0, a_1, \dots, a_{n-1}) \in \mathbb{R}_+^n$. From (1.3), one finds that

$$\begin{aligned} V(\mathbf{a}; 0, 0) &= (-1)^{n-1} \prod_{j=1}^n (a_j - a_0) V(\bar{\mathbf{a}}; 0, 0) \\ &= (a_n - a_0) \prod_{j=1}^{n-1} (a_n - a_j) V(\tilde{\mathbf{a}}; 0, 0), \end{aligned} \quad (2.3)$$

$$V_0(\mathbf{a}) = \prod_{j=1}^{n-1} (a_n - a_j) V_0(\tilde{\mathbf{a}}), \quad (2.4)$$

$$V_i(\mathbf{a}) = (-1)^n \prod_{j=1}^{n-1} (a_j - a_0) V_i(\bar{\mathbf{a}}) + \prod_{j=1}^{n-1} (a_n - a_j) V_i(\tilde{\mathbf{a}}), \quad 1 < i < n, \quad (2.5)$$

$$V_n(\mathbf{a}) = \prod_{j=1}^{n-1} (a_j - a_0) V_0(\bar{\mathbf{a}}). \quad (2.6)$$

The expansion of the generalized Vandermonde determinant (1.1) equals

$$V(\mathbf{a}; \varphi) = \sum_{i=0}^n (-1)^{n+i} \varphi(a_i) V_i(\mathbf{a}). \quad (2.7)$$

It is easy to see that

$$\begin{aligned} & \int_D \varphi^{(n)} \left(\sum_{i=0}^n a_i x_i \right) dx \\ &= \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-1} x_i} \varphi^{(n)} \left(a_0 + \sum_{i=1}^n (a_i - a_0) x_i \right) dx_1 dx_2 \cdots dx_n. \end{aligned} \quad (2.8)$$

For $n = 1$, formula (2.1) is true, since

$$(a_1 - a_0) \int_0^1 \varphi'(a_0 + (a_1 - a_0)x_1) dx_1 = \varphi(a_1) - \varphi(a_0) = V(\mathbf{a}; \varphi).$$

Assume that (2.1) holds for $n - 1$ with $n \geq 2$. Then

$$\begin{aligned} V(\bar{\mathbf{a}}; \varphi) &= (-1)^{n-1} \varphi(a_n) V_0(\bar{\mathbf{a}}) + \sum_{i=1}^{n-1} (-1)^{n-1+i} \varphi(a_i) V_i(\bar{\mathbf{a}}) \\ &= V(\bar{\mathbf{a}}; 0, 0) \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-2} x_i} \\ &\quad \varphi^{(n-1)} \left(a_n + \sum_{i=1}^{n-1} (a_i - a_n) x_i \right) dx_1 dx_2 \cdots dx_{n-1} \quad (2.9) \end{aligned}$$

and

$$\begin{aligned} V(\tilde{\mathbf{a}}; \varphi) &= \sum_{i=0}^{n-1} (-1)^{n-1+i} \varphi(a_i) V_i(\tilde{\mathbf{a}}) \\ &= V(\tilde{\mathbf{a}}; 0, 0) \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-2} x_i} \\ &\quad \varphi^{(n-1)} \left(a_0 + \sum_{i=1}^{n-1} (a_i - a_0) x_i \right) dx_1 dx_2 \cdots dx_{n-1}. \quad (2.10) \end{aligned}$$

Combination of (2.3) to (2.10) shows that

$$\begin{aligned} &V(\mathbf{a}; 0, 0) \int_D \varphi^{(n)} \left(\sum_{i=0}^n a_i x_i \right) dx \\ &= V(\mathbf{a}; 0, 0) \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-1} x_i} \\ &\quad \varphi^{(n)} \left(a_0 + \sum_{i=1}^n (a_i - a_0) x_i \right) dx_1 dx_2 \cdots dx_n \\ &= \frac{V(\mathbf{a}; 0, 0)}{a_n - a_0} \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-2} x_i} \left[\varphi^{(n-1)} \left(a_n + \sum_{i=1}^{n-1} (a_i - a_n) x_i \right) \right. \\ &\quad \left. - \varphi^{(n-1)} \left(a_0 + \sum_{i=1}^{n-1} (a_i - a_0) x_i \right) \right] dx_1 dx_2 \cdots dx_{n-1} \\ &= (-1)^{n-1} \prod_{j=1}^{n-1} (a_j - a_0) V(\bar{\mathbf{a}}; 0, 0) \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-2} x_i} \\ &\quad \varphi^{(n-1)} \left(a_n + \sum_{i=1}^{n-1} (a_i - a_n) x_i \right) dx_1 \cdots dx_{n-1} \\ &\quad - \prod_{j=1}^{n-1} (a_n - a_j) V(\tilde{\mathbf{a}}; 0, 0) \int_0^1 \int_0^{1-x_1} \cdots \int_0^{1-\sum_{i=1}^{n-2} x_i} \end{aligned}$$

$$\begin{aligned}
& \varphi^{(n-1)} \left(a_0 + \sum_{i=1}^{n-1} (a_i - a_0) x_i \right) dx_1 \cdots dx_{n-1} \\
&= (-1)^{n-1} \prod_{j=1}^{n-1} (a_j - a_0) \left[(-1)^{n-1} \varphi(a_n) V_0(\bar{\mathbf{a}}) + \sum_{i=1}^{n-1} (-1)^{n-1+i} \varphi(a_i) V_i(\bar{\mathbf{a}}) \right] \\
&\quad - \prod_{j=1}^{n-1} (a_n - a_j) \sum_{i=1}^{n-1} (-1)^{n-1+i} \varphi(a_i) V_i(\bar{\mathbf{a}}) \\
&= \varphi(a_n) \prod_{j=1}^{n-1} (a_j - a_0) V_0(\bar{\mathbf{a}}) - (-1)^{n-1} \varphi(a_0) \prod_{j=1}^{n-1} (a_n - a_j) V_0(\bar{\mathbf{a}}) \\
&\quad + \sum_{i=1}^{n-1} (-1)^{n-1+i} \varphi(a_i) \left[(-1)^{n-1} \prod_{j=1}^{n-1} (a_j - a_0) V_i(\bar{\mathbf{a}}) - \prod_{j=1}^{n-1} (a_n - a_j) V_i(\bar{\mathbf{a}}) \right] \\
&= \varphi(a_n) V_n(\mathbf{a}) + \sum_{i=1}^{n-1} (-1)^{n+i} \varphi(a_i) V_i(\mathbf{a}) + (-1)^n \varphi(a_0) V_0(\mathbf{a}) \\
&= V(\mathbf{a}; \varphi).
\end{aligned}$$

Thus, by induction, the required formula (2.1) is proved. \square

Lemma 2.2. *Let r be an integer and $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$. Then*

$$V(\mathbf{a}; r, 0) = \begin{cases} V(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\dots+i_n=r \\ i_0, i_1, \dots, i_n \geq 0 \text{ are integers}}} \prod_{k=0}^n a_k^{i_k} & \text{for } r > 0, \\ 0 & \text{for } r = 0, -1, \dots, -(n-1), \\ (-1)^n V(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\dots+i_n=-r \\ i_0, i_1, \dots, i_n \in \mathbb{N}}} \prod_{k=0}^n a_k^{-i_k} & \text{for } r \leq -n. \end{cases} \quad (2.11)$$

Proof. Taking $V_n(\mathbf{a}; r, 0) \triangleq V(\mathbf{a}; r, 0)$. It is obvious that, if $r = -1, \dots, -n$, then $V_n(\mathbf{a}; r, 0) = V(\mathbf{a}; r, 0) = 0$.

In the case of $n \in \mathbb{N}$ and $r \geq 0$, we will verify Lemma 2.2 by mathematical induction. It is clear that identity (2.11) holds trivially for $n = 1$, since

$$\begin{aligned}
V_2(\mathbf{a}; r, 0) &= a_1^{r+1} - a_0^{r+1} \\
&= (a_1 - a_0) (a_1^r + a_1^{r-1} a_0 + \cdots + a_0^r) \\
&= (a_1 - a_0) \sum_{\substack{i_0+i_1=r \\ i_0, i_1 \geq 0 \text{ are integers}}} a_0^{i_0} a_1^{i_1}.
\end{aligned}$$

Suppose identity (2.11) is true for $n-1$ and positive integers t . That is

$$V_{n-1}(\mathbf{a}; t, 0) = V_{n-1}(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\dots+i_{n-1}=t \\ i_0, i_1, \dots, i_{n-1} \geq 0 \text{ are integers}}} \prod_{k=0}^{n-1} a_k^{i_k}. \quad (2.12)$$

By (1.2) and (2.12), one reveals

$$\begin{aligned}
V_n(\mathbf{a}; r, 0) &= \begin{vmatrix} 1 & a_0 - a_n & \cdots & a_0^{n-1} - a_0^{n-2}a_n & a_0^{n+r} - a_0^{n-1}a_n^{r+1} \\ 1 & a_1 - a_n & \cdots & a_1^{n-1} - a_1^{n-2}a_n & a_1^{n+r} - a_1^{n-1}a_n^{r+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_{n-1} - a_n & \cdots & a_{n-1}^{n-1} - a_{n-1}^{n-2}a_n & a_{n-1}^{n+r} - a_{n-1}^{n-1}a_n^{r+1} \\ 1 & 0 & \cdots & 0 & 0 \end{vmatrix} \\
&= (-1)^{n+2} \begin{vmatrix} a_0 - a_n & \cdots & a_0^{n-1} - a_0^{n-2}a_n & a_0^{n+r} - a_0^{n-1}a_n^{r+1} \\ a_1 - a_n & \cdots & a_1^{n-1} - a_1^{n-2}a_n & a_1^{n+r} - a_1^{n-1}a_n^{r+1} \\ \cdots & \cdots & \cdots & \cdots \\ a_{n-1} - a_n & \cdots & a_{n-1}^{n-1} - a_{n-1}^{n-2}a_n & a_{n-1}^{n+r} - a_{n-1}^{n-1}a_n^{r+1} \end{vmatrix} \\
&= \prod_{i=0}^{n-1} (a_n - a_i) \begin{vmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^{n-1} & \sum_{t+i_n=r} a_0^{n-1+t} a_n^{i_n} \\ 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} & \sum_{t+i_n=r} a_1^{n-1+t} a_n^{i_n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_{n-1} & a_{n-1}^2 & \cdots & a_{n-1}^{n-1} & \sum_{t+i_n=r} a_{n-1}^{n-1+t} a_n^{i_n} \end{vmatrix} \\
&= \prod_{i=0}^{n-1} (a_n - a_i) \sum_{t+i_n=r} a_n^{i_n} \begin{vmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^{n-1} & a_0^{n-1+t} \\ 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} & a_1^{n-1+t} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_{n-1} & a_{n-1}^2 & \cdots & a_{n-1}^{n-1} & a_{n-1}^{n-1+t} \end{vmatrix} \\
&= \prod_{i=0}^{n-1} (a_n - a_i) \sum_{t+i_n=r} a_n^{i_n} V_{n-1}(\mathbf{a}; t, 0) \\
&= \prod_{i=0}^{n-1} (a_n - a_i) \cdot V_{n-1}(\mathbf{a}; 0, 0) \sum_{t+i_n=r} a_n^{i_n} \sum_{\substack{i_0+i_1+\cdots+i_{n-1}=t, \\ i_0, i_1, \dots, i_{n-1} \geq 0 \text{ are integers}}} \prod_{k=0}^{n-1} a_k^{i_k} \\
&= V(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\cdots+i_n=r, \\ i_0, i_1, \dots, i_n \geq 0 \text{ are integers}}} \prod_{k=0}^n a_k^{i_k}.
\end{aligned}$$

By induction, it is showed that (2.11) holds for n and positive integers r .

From (1.3), it is deduced easily that

$$\begin{aligned}
V(\mathbf{a}^{-1}; 0, 0) &= \sum_{i=0}^n (-1)^{n+i} a_i^{-n} V_i(\mathbf{a}^{-1}) \\
&= \prod_{0 \leq i < j \leq n} (a_j^{-1} - a_i^{-1}) \\
&= \prod_{0 \leq i \leq n} a_i^{-n} \prod_{0 \leq i < j \leq n} (a_i - a_j) \\
&= (-1)^{n(n+1)/2} \prod_{0 \leq i \leq n} a_i^{-n} V(\mathbf{a}; 0, 0),
\end{aligned} \tag{2.13}$$

where $\mathbf{a}^{-1} = (a_0^{-1}, a_1^{-1}, \dots, a_n^{-1}) \in \mathbb{R}_+^{n+1}$.

In the case of $r < -n$, we have $-(n+r) > 0$. From (1.2) and (2.13), one finds

$$\begin{aligned}
V(\mathbf{a}; r, 0) &= \prod_{0 \leq i \leq n} a_i^{n-1} \begin{vmatrix} a_0^{-(n-1)} & a_0^{-(n-1)} & \cdots & 1 & a_0^{r+1} \\ a_1^{-(n-1)} & a_1^{-(n-2)} & \cdots & 1 & a_1^{r+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{n-1}^{-(n-1)} & a_{n-1}^{-(n-2)} & \cdots & 1 & a_{n-1}^{r+1} \end{vmatrix} \\
&= (-1)^{n(n-1)/2} \prod_{0 \leq i \leq n} a_i^{n-1} \begin{vmatrix} 1 & a_0^{-1} & \cdots & a_0^{-(n-1)} & a_0^{-[n-(n+r+1)]} \\ 1 & a_1^{-1} & \cdots & a_1^{-(n-1)} & a_1^{-[n-(n+r+1)]} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & a_{n-1}^{-1} & \cdots & a_{n-1}^{-(n-1)} & a_{n-1}^{-[n-(n+r+1)]} \end{vmatrix} \\
&= (-1)^{n(n-1)/2} \prod_{0 \leq i \leq n} a_i^{n-1} V(\mathbf{a}^{-1}; 0, 0) \sum_{\substack{i_0+i_1+\cdots+i_n=-(n+r+1) \\ i_0, i_1, \dots, i_n \geq 0 \text{ are integers}}} \prod_{k=0}^n a_k^{-i_k} \\
&= (-1)^n \prod_{0 \leq i \leq n} a_i^{-1} V(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\cdots+i_n=-(n+r+1) \\ i_0, i_1, \dots, i_n \geq 0 \text{ are integers}}} \prod_{k=0}^n a_k^{-i_k} \\
&= (-1)^n V(\mathbf{a}; 0, 0) \sum_{\substack{i_0+i_1+\cdots+i_n=-r \\ i_0, i_1, \dots, i_n \geq 1 \text{ are integers}}} \prod_{k=0}^n a_k^{-i_k}.
\end{aligned}$$

The proof of Lemma 2.2 is completed. \square

Lemma 2.3 ([19, 21, 23]). *Let f be a continuous function and p a positive continuous weight on I . Then the weighted arithmetic mean of function f with weight p defined by*

$$F(x, y) = \begin{cases} \frac{\int_x^y p(t)f(t) dt}{\int_x^y p(t) dt}, & x \neq y, \\ f(x), & x = y \end{cases} \quad (2.14)$$

is increasing (decreasing) on I^2 if f is increasing (decreasing) on I .

3. DEFINITION AND BASIC PROPERTIES

Now we are in a position to give the definition of mean values of several positive numbers with two parameters.

Definition 3.1. Let $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$ with $a_i \neq a_j$ for $i \neq j$. The extended mean values $E(p, q; \mathbf{a})$ of $(n+1)$ -tuple \mathbf{a} with two parameters $p \in \mathbb{R}$ and

$q \in \mathbb{R}$ are defined as

$$E(p, q; \mathbf{a}) = \begin{cases} \left[\prod_{k=1}^n \frac{k+q}{k+p} \cdot \frac{V(\mathbf{a}; p, 0)}{V(\mathbf{a}; q, 0)} \right]^{1/(p-q)} \\ \text{for } (p-q) \prod_{k=1}^n [(k+p)(k+q)] \neq 0, \\ \left[\frac{(-1)^{q+1}(-q-1)!(q+n)!}{\prod_{k=1}^n (k+p)} \cdot \frac{V(\mathbf{a}; p, 0)}{V(\mathbf{a}; q, 1)} \right]^{1/(p-q)} \\ \text{for } p \neq q = -1, -2, \dots, -n, \\ \left[\frac{(-1)^{q-p}(-q-1)!(q+n)!}{(-p-1)!(p+n)!} \cdot \frac{V(\mathbf{a}; p, 1)}{V(\mathbf{a}; q, 1)} \right]^{1/(p-q)} \\ \text{for } p \neq q \text{ and } p, q = -1, -2, \dots, -n, \\ \exp \left[\frac{V(\mathbf{a}; p, 1)}{V(\mathbf{a}; p, 0)} - \sum_{k=1}^n \frac{1}{k+p} \right] \\ \text{for } p = q \neq -1, -2, \dots, -n, \\ \exp \left[\frac{V(\mathbf{a}; p, 2)}{2V(\mathbf{a}; p, 1)} - \sum_{\substack{k=1 \\ k \neq -p}}^n \frac{1}{k+p} \right] \\ \text{for } p = q = -1, -2, \dots, -n. \end{cases} \quad (3.1)$$

The mean values $E(p, q; \mathbf{a})$ have the same usual properties as many other means listed in [3].

Theorem 3.1. *The following basic properties hold:*

- (1) $E(p, q; \mathbf{a}) = E(q, p; \mathbf{a})$,
- (2) $\lim_{p \rightarrow \infty} E(p, q; \mathbf{a}) = \lim_{q \rightarrow \infty} E(p, q; \mathbf{a}) = \max\{a_0, a_1, \dots, a_n\}$,
- (3) $\lim_{p \rightarrow -\infty} E(p, q; \mathbf{a}) = \lim_{q \rightarrow -\infty} E(p, q; \mathbf{a}) = \min\{a_0, a_1, \dots, a_n\}$,
- (4) $\lim_{p \rightarrow q} E(p, q; \mathbf{a}) = E(q, q; \mathbf{a})$,
- (5) $\min\{a_0, a_1, \dots, a_n\} \leq E(p, q; \mathbf{a}) \leq \max\{a_0, a_1, \dots, a_n\}$,
- (6) $E(p, q; \mathbf{a}) = a_0$ if and only if $a_0 = a_1 = \dots = a_n$,
- (7) For $t > 0$, $E(p, q; t\mathbf{a}) = tE(p, q; \mathbf{a})$,
- (8) $[E(p, q; \mathbf{a})]^{p-q} = [E(p, t; \mathbf{a})]^{p-t} [E(t, q; \mathbf{a})]^{t-q}$,

where $t\mathbf{a} = (ta_0, ta_1, \dots, ta_n)$.

Proof. These follow straightforwardly from Definition 3.1 and Lemma 2.1 by standard arguments. \square

The theorem below gives an alternative expression of the mean values $E(p, q; \mathbf{a})$.

Theorem 3.2. *Let $n \in \mathbb{N}$, $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$, and $p, q \in \mathbb{R}$. Then*

$$E(p, q; \mathbf{a}) = \begin{cases} \left[\frac{\int_D \varphi_1^{(n)}(p, \sum_{i=0}^n a_i x_i) dx}{\int_D \varphi_1^{(n)}(q, \sum_{i=0}^n a_i x_i) dx} \right]^{1/(p-q)}, & p \neq q, \\ \exp \left\{ \frac{\int_D \varphi_2^{(n)}(p, \sum_{i=0}^n a_i x_i) dx}{\int_D \varphi_1^{(n)}(p, \sum_{i=0}^n a_i x_i) dx} \right\}, & p = q, \end{cases} \quad (3.2)$$

where D is the simplex defined by (2.2), $dx = dx_1 dx_2 \dots dx_n$ is the differential of the volume in D , $x_0 = 1 - \sum_{i=1}^n x_i$, $\varphi_1^{(n)}(p, t) = t^p$ and $\varphi_2^{(n)}(p, t) = t^p \ln t$.

Proof. This follows from substituting

$$\varphi_1(p, t) = \begin{cases} t^{n+p} \prod_{k=1}^n \frac{1}{k+p} & \text{for } p \neq -1, \dots, -n, \\ \frac{t^{n+p} \ln t}{(-1)^{p+1} (-p-1)! (n+p)!} & \text{for } p = -1, \dots, -n \end{cases} \quad (3.3)$$

and

$$\varphi_2(p, t) = \begin{cases} t^{n+p} \left(\ln t - \sum_{k=1}^n \frac{1}{k+p} \right) \prod_{k=1}^n \frac{1}{k+p} & \text{for } p \neq -1, \dots, -n, \\ \frac{t^{n+p} \ln t}{(-1)^{p+1} (-p-1)! (n+p)!} \left(\frac{\ln t}{2} - \sum_{\substack{k=1 \\ k \neq -p}}^n \frac{1}{k+p} \right) & \text{for } p = -1, \dots, -n \end{cases} \quad (3.4)$$

into Lemma 2.1. \square

Corollary 3.1. *Let $n \in \mathbb{N}$ and $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$. Then*

$$S_r(\mathbf{a}) = E(r, 0; \mathbf{a}) = \begin{cases} [n! \int_D \varphi_1^{(n)}(r, \sum_{i=0}^n a_i x_i) dx]^{1/r}, & r \neq 0, \\ \exp \{ n! \int_D \varphi_2^{(n)}(0, \sum_{i=0}^n a_i x_i) dx \}, & r = 0, \end{cases} \quad (3.5)$$

$$J_r(\mathbf{a}) = E(r, r-1; \mathbf{a}) = \frac{\int_D \varphi_1^{(n)}(r, \sum_{i=0}^n a_i x_i) dx}{\int_D \varphi_1^{(n)}(r-1, \sum_{i=0}^n a_i x_i) dx}, \quad (3.6)$$

$$E(p, q; \mathbf{a}) = \left\{ \frac{[S_p(\mathbf{a})]^p}{[S_q(\mathbf{a})]^q} \right\}^{1/(p-q)}, \quad (3.7)$$

where D is defined by (2.2), $dx = dx_1 dx_2 \cdots dx_n$ denotes the differential of the volume in D , and $x_0 = 1 - \sum_{i=1}^n x_i$.

Some mean values are special cases of the mean values $E(p, q; \mathbf{a})$ as Corollary 3.1 reveals. From Definition 3.1 and Lemma 2.2, we further obtain the following

Theorem 3.3. *Let $n \in \mathbb{N}$ and $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$. Then*

- (1) $E(1, 0; \mathbf{a}) = S_1(\mathbf{a}) = J_1(\mathbf{a}) = A(\mathbf{a})$,
- (2) $E(0, -(n+1); \mathbf{a}) = S_{-(n+1)}(\mathbf{a}) = G(\mathbf{a})$,
- (3) $E(-(n+1), -(n+2); \mathbf{a}) = J_{-(n+1)}(\mathbf{a}) = H(\mathbf{a})$,
- (4) $E(0, 0; \mathbf{a}) = S_0(\mathbf{a}) = I(\mathbf{a})$,
- (5) $E(0, -1; \mathbf{a}) = S_{-1}(\mathbf{a}) = J_0(\mathbf{a}) = L(\mathbf{a})$,

where $A(\mathbf{a})$, $G(\mathbf{a})$, $H(\mathbf{a})$, $I(\mathbf{a})$, and $L(\mathbf{a})$ denote respectively the arithmetic, geometric, harmonic, exponential, and logarithmic means of $n+1$ positive numbers a_0, a_1, \dots, a_n .

Proof. The proof is straightforward. \square

Some special cases of $E(p, q; \mathbf{a})$ have relationships with the generalized symmetric means $\sum_n^k(\mathbf{a})$ of order k introduced in [7] by D. M. DeTemple and J. M. Robertson.

Theorem 3.4. *Let r be a positive integer and $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$ for $n \in \mathbb{N}$. Then*

$$E(r, 0; \mathbf{a}) = S_r(\mathbf{a}) = \sum_{n+1}^r(\mathbf{a}), \quad (3.8)$$

$$E(r, r-1; \mathbf{a}) = J_r(\mathbf{a}) = \frac{\sum_{n+1}^r (\mathbf{a})}{\sum_{n+1}^{r-1} (\mathbf{a})}, \quad (3.9)$$

where

$$\sum_{n+1}^k (\mathbf{a}) = \binom{n+k}{n}^{-1} \sum_{\substack{i_0+i_1+\dots+i_n=k \\ i_0, i_1, \dots, i_n \geq 0}} a_0^{i_0} a_1^{i_1} \cdots a_n^{i_n} \quad (3.10)$$

denotes the k -th generalized symmetric mean of $n+1$ positive numbers a_0, a_1, \dots, a_n .

4. MONOTONICITY AND APPLICATIONS

Theorem 4.1. *The mean values $E(p, q; \mathbf{a})$ are increasing strictly with both $p \in \mathbb{R}$ and $q \in \mathbb{R}$.*

Proof. Let $g(x) = \sum_{i=0}^n a_i x_i$. If $p = q$, then

$$E(p, p; \mathbf{a}) = \exp \left\{ \frac{\int_D \varphi_2^{(n)}(p, g(x)) dx}{\int_D \varphi_1^{(n)}(p, g(x)) dx} \right\}$$

and

$$\begin{aligned} & \left[\int_D \varphi_1^{(n)}(p, g(x)) dx \right]^2 \frac{d}{dp} [\ln E(p, p; \mathbf{a})] \\ &= \int_D \varphi_2^{(n)}(p, g(x)) \varphi_2^{(n)}(0, g(x)) dx \int_D \varphi_1^{(n)}(p, g(x)) dx \\ & \quad - \left[\int_D \varphi_1^{(n)}(p, g(x)) \varphi_2^{(n)}(0, g(x)) dx \right]^2 \\ &= \int_D [g(x)]^p [\ln g(x)]^2 dx \int_D [g(x)]^p dx - \left[\int_D [g(x)]^p \ln[g(x)] dx \right]^2 \\ &> 0 \end{aligned}$$

by using the Cauchy-Schwartz-Buniakowski integral inequality. This implies that the mean values $E(p, p; \mathbf{a})$ are strictly increasing with $p \in \mathbb{R}$.

If $p \neq q$, then

$$E(p, q; \mathbf{a}) = \left[\frac{\int_D \varphi_1^{(n)}(p, g(x)) dx}{\int_D \varphi_1^{(n)}(q, g(x)) dx} \right]^{1/(p-q)}$$

and

$$\begin{aligned} \ln E(p, q; \mathbf{a}) &= \frac{1}{p-q} \int_q^p \frac{\int_D \varphi_2^{(n)}(\theta, g(x)) dx}{\int_D \varphi_1^{(n)}(\theta, g(x)) dx} d\theta \\ &= \frac{1}{p-q} \int_q^p \ln E(\theta, \theta; \mathbf{a}) d\theta. \end{aligned}$$

From Lemma 2.3 and the fact that $E(p, p; \mathbf{a})$ are strictly increasing with $p \in \mathbb{R}$, it is not difficult to see that the mean values $E(p, q; \mathbf{a})$ are increasing strictly with both p and q . The proof is complete. \square

As applications of the monotonicity of the mean values $E(p, q; \mathbf{a})$, the following inequalities about some mean values are obtained readily.

Corollary 4.1. *If $p_1 < p_2$ and $q_1 < q_2$, then*

$$E_{p_1, q_1}(\mathbf{a}) < E_{p_2, q_2}(\mathbf{a}). \quad (4.1)$$

If $q \geq 0$, then

$$S_q(\mathbf{a}) \leq E(p, q; \mathbf{a}). \quad (4.2)$$

If $q \geq p - 1$, then

$$J_p(\mathbf{a}) \leq E(p, q; \mathbf{a}). \quad (4.3)$$

If $p \leq q$, then

$$J_p(\mathbf{a}) \leq J_q(\mathbf{a}) \quad (4.4)$$

and

$$S_p(\mathbf{a}) \leq S_q(\mathbf{a}). \quad (4.5)$$

If $p \geq 1$, then

$$S_p(\mathbf{a}) \leq J_p(\mathbf{a}). \quad (4.6)$$

Corollary 4.2. *Let $G(\mathbf{a})$, $L(\mathbf{a})$, $I(\mathbf{a})$ and $A(\mathbf{a})$ denote the geometric, logarithmic, exponential and arithmetic means of $n + 1$ positive numbers respectively. Then*

$$G(\mathbf{a}) < L(\mathbf{a}) < I(\mathbf{a}) < A(\mathbf{a}). \quad (4.7)$$

Proof. From (4.5) in Corollary 4.1, it follows that

$$S_{-(n+1)}(\mathbf{a}) < S_{-1}(\mathbf{a}) < S_0(\mathbf{a}) < S_1(\mathbf{a}).$$

Thus, inequalities in (4.7) is obtained immediately from Theorem 3.3. \square

Corollary 4.3. *Let $r \in \mathbb{N}$, $n \in \mathbb{N}$, and $\mathbf{a} = (a_0, a_1, \dots, a_n) \in \mathbb{R}_+^{n+1}$. Then*

$$\left[\sum_{n+1}^{[r]}(\mathbf{a}) \right]^2 \leq \sum_{n+1}^{[r+1]}(\mathbf{a}) \sum_{n+1}^{[r-1]}(\mathbf{a}) \quad (4.8)$$

and

$$\left[\sum_{n+1}^{[r]}(\mathbf{a}) \right]^{\frac{1}{r}} \leq \left[\sum_{n+1}^{[r+1]}(\mathbf{a}) \right]^{\frac{1}{r+1}}. \quad (4.9)$$

Equalities in (4.8) and (4.9) are valid if and only if $a_0 = a_1 = \dots = a_n$.

Proof. By using Corollary 4.1, we find

$$\begin{aligned} J_r(\mathbf{a}) &\leq J_{r+1}(\mathbf{a}), \\ S_r(\mathbf{a}) &\leq S_{r+1}(\mathbf{a}). \end{aligned}$$

Further considering Theorem 3.4 proves the required results. \square

Corollary 4.4. *If $r \geq 1$, then*

$$S_r(\mathbf{a}) \leq M_r(\mathbf{a}), \quad (4.10)$$

where

$$M_r(\mathbf{a}) = \begin{cases} \left[\frac{1}{n+1} \sum_{i=0}^n a_i^r \right]^{1/r}, & r \neq 0, \\ G(\mathbf{a}), & r = 0. \end{cases} \quad (4.11)$$

Proof. Let $g(x) = \sum_{i=0}^n a_i x_i$. In the case of $r(r-1) > 0$, using the arithmetic-geometric mean inequality, we obtain

$$\begin{aligned} \varphi_1^{(n)}(r, g(x)) &= \left[a_0 + \sum_{i=1}^n (a_i - a_0)x_i \right]^r \\ &= \left[\left(1 - \sum_{i=1}^n x_i \right) a_0 + \sum_{i=1}^n x_i a_i \right]^r \\ &< \left(1 - \sum_{i=1}^n x_i \right) a_0^r + \sum_{i=1}^n x_i a_i^r \\ &= a_0^r + \sum_{i=1}^n (a_i^r - a_0^r)x_i, \end{aligned}$$

which is equivalent to

$$\begin{aligned} S_r^r(\mathbf{a}) &= n! \int_D \varphi_1^{(n)}(r, g(x)) dx \\ &< n! \int_D \left(a_0^r + \sum_{i=1}^n (a_i^r - a_0^r)x_i \right) dx \\ &= \frac{1}{n+1} \sum_{i=0}^n a_i^r \\ &= M_r^r(\mathbf{a}). \end{aligned}$$

Therefore, it is deduced that $S_r(\mathbf{a}) < M_r(\mathbf{a})$ for $r > 1$ and $S_r(\mathbf{a}) > M_r(\mathbf{a})$ for $r < 0$.

By the same argument as above, if $0 < r < 1$, then $S_r(\mathbf{a}) > M_r(\mathbf{a})$ and

$$S_0(\mathbf{a}) < S_{-(n+1)}(\mathbf{a}) = G(\mathbf{a}) = M_0(\mathbf{a}).$$

Thus, if $r < 1$, then $S_r(\mathbf{a}) > M_r(\mathbf{a})$. The proof is complete. \square

REFERENCES

- [1] H. Alzer, *Über eine einparametrische familie von mittewerten (On a one-parameter family of means)*, Bayer. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. **1987** (1988), 1–9. (German)
- [2] H. Alzer, *Über eine einparametrische familie von Mitlewerten (On a one-parameter family of mean values)*, II, Bayer. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. **1988** (1989), 23–29. (German)
- [3] P. S. Bullen, *Handbook of Means and Their Inequalities*, Mathematics and its Applications (Dordrecht) **560**, Kluwer Academic Publishers, Dordrecht, 2003.
- [4] B. C. Carlson, *Algorithms involving arithmetic and geometric means*, Amer. Math. Monthly, **78** (1971), 496–505.
- [5] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 2nd ed., Hunan Education Press, Changsha, China, 1993. (Chinese)
- [6] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 3rd ed., Shandong Science and Technology Press, Jinan City, Shandong Province, China, 2004. (Chinese)
- [7] D. W. DeTemple and J. M. Robertson, *On generalized symmetric means of two variables*, Univ. Beograd. Publ. Elek. Fak Ser. Mat. Fiz. **No. 634–677** (1979), 236–238.
- [8] X. Li and R. N. Mohapatra, *Extended means as weighted means*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **457** (2001), no. 2009, 1273–1275.
- [9] E. Neuman, *Stolarsky means of several variables*, J. Inequal. Pure Appl. Math. **6** (2005), no. 2, Art. 30. Available online at <http://jipam.vu.edu.au/article.php?sid=499>.

- [10] C. E. M. Pearce, J. Pečarić and V. Šimić, *On weighted generalized logarithmic means*, Houston J. Math. **24** (1998), 459–465.
- [11] J. Pečarić, F. Qi, V. Šimić and S.-L. Xu, *Refinements and extensions of an inequality, III*, J. Math. Anal. Appl. **227** (1998), no. 2, 439–448.
- [12] F. Qi, *Generalized abstracted mean values*, J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Art. 4. Available online at <http://jipam.vu.edu.au/article.php?sid=97>. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 4, 633–642. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [13] F. Qi, *Generalized weighted mean values with two parameters*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **454** (1998), no. 1978, 2723–2732.
- [14] F. Qi, *On a two-parameter family of nonhomogeneous mean values*, Tamkang J. Math. **29** (1998), no. 2, 155–163.
- [15] F. Qi, *The extended mean values: definition, properties, monotonicities, comparison, convexities, generalizations, and applications*, Cubo Mat. Educ. **5** (2003), no. 3, 63–90. RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 5. Available online at <http://rgmia.vu.edu.au/v5n1.html>.
- [16] F. Qi and Q.-M. Luo, *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), 356–359.
- [17] F. Qi and Q.-M. Luo, *Refinements and extensions of an inequality*, Math. Inform. Quart. **9** (1999), no. 1, 23–25.
- [18] F. Qi, J.-Q. Mei, D.-F. Xia, and S.-L. Xu, *New proofs of weighted power mean inequalities and monotonicity for generalized weighted mean values*, Math. Inequal. Appl. **3** (2000), no. 3, 377–383.
- [19] F. Qi, J. Sándor, S. S. Dragomir, and A. Sofo, *Notes on the Schur-convexity of the extended mean values*, Taiwanese J. Math. **9** (2005), no. 3, 411–420. RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 3, 19–27. Available online at <http://rgmia.vu.edu.au/v5n1.html>.
- [20] F. Qi and S.-L. Xu, *Refinements and extensions of an inequality, II*, J. Math. Anal. Appl. **211** (1997), no. 2, 616–620.
- [21] F. Qi, S.-L. Xu, and L. Debnath, *A new proof of monotonicity for extended mean values*, Internat. J. Math. Math. Sci. **22** (1999), no. 2, 415–420.
- [22] F. Qi and Sh.-Q. Zhang, *Note on monotonicity of generalized weighted mean values*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **455** (1999), no. 1989, 3259–3260.
- [23] H.-N. Shi, Sh.-H. Wu, and F. Qi, *An alternative note on the Schur-convexity of the extended mean values*, Math. Inequal. Appl. **9** (2006), accepted.
- [24] K. B. Stolarsky, *Generalizations of the logarithmic mean*, Math. Mag. **48** (1975), 87–92.
- [25] A. Witkowski, *Weighted extended mean values*, Colloq. Math. **100** (2004), no. 1, 111–117. RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 6. Available online at <http://rgmia.vu.edu.au/v7n1.html>.
- [26] Zh.-G. Xiao and Zh.-H. Zhang, *The Henleman mean of n positive numbers*, J. Yueyang Normal Univ. **14** (2001), no. 2, 1–5. (Chinese).
- [27] Zh.-G. Xiao and Zh.-H. Zhang, *The Stolarsky mean of n positive numbers*, J. Yueyang Normal Univ. **14** (2001), no. 4, 5–8. (Chinese).
- [28] Zh.-G. Xiao and Zh.-H. Zhang, *The inequalities $G \leq L \leq I \leq A$ in n variables*, J. Inequal. Pure Appl. Math. **4** (2003), no. 2, Art. 39. Available online at <http://jipam.vu.edu.au/article.php?sid=277>.
- [29] Zh.-H. Zhang, *Three classes of new means in $n+1$ variables and their applications*, J. Hunan Ed. Inst. **15** (1997), no. 5, 130–136. (Chinese)

(Zh.-G. Xiao) HUNAN INSTITUTE OF SCIENCE AND TECHNOLOGY, YUEYANG CITY, HUNAN PROVINCE, 414000, CHINA

E-mail address: xiaozg@163.com

(Zh.-H. Zhang) ZIXING EDUCATIONAL RESEARCH SECTION, CHENZHOU CITY, HUNAN PROVINCE, 423400, CHINA

E-mail address: zxzh1234@163.com

(F. Qi) RESEARCH INSTITUTE OF MATHEMATICAL INEQUALITY THEORY, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: qifeng@hpu.edu.cn, fengqi618@member.ams.org, qifeng618@hotmail.com, qifeng618@msn.com

URL: <http://rgmia.vu.edu.au/qi.html>